

Dental Health Problems among the Patients Attending in the Orthodontic Department of Dhaka Dental College & Hospital

MA Habib¹, RMG Rabbani², A Yousuf³, AK Saha⁴, HH Imon⁵, ASMR Haque⁶, MZ Hossain⁷

Abstract

Objective: To assess pattern of dental health problems among Bangladeshi dental patients.

Study design: Descriptive study.

Place of study: Department of Orthodontics & Dentofacial Orthopaedics of Dhaka Dental College & Hospital, Dhaka.

Period of study: From September '2011 to February '2012.

Sample selection: In this study, a total of 500 patients (250 males & 250 females) were selected from Orthodontic department of Dhaka Dental College & Hospital.

Results: This study was conducted among 500 patients attending at the department of Orthodontics and Dentofacial Orthopaedics, Dhaka Dental College and Hospital. Gingivitis was the commonest problem in Bangladeshi patients according to the present study. The second most dental problem was dental caries. The orthodontic problem was comparable to the observations made by others. Males and females were equally affected with slight variations in the nature of problems among genders.

Conclusion: Good oral health is essential to improve individual overall health & well-being. We urge to take this information & use it for program planning & advocating for the health of patients, specially for the patients who will receive orthodontic treatment. Therefore, current orthodontic students should receive more education & training before the management of malocclusion to improve the overall quality of care for orthodontic patients.

Key words: Dental health, caries, gingivitis, inflammation, periodontitis.

1. Dr. Md. Ahsan Habib BDS, FCPS, Department of Orthodontics and Dentofacial Orthopaedics, Dhaka Dental College and Hospital

2. Dr Rashed Md Golam Rabbani BDS, FCPS Department of Orthodontics and Dentofacial Orthopaedics, Dhaka Dental College and Hospital

3. Dr. Anwar Yousuf, BDS, PhD, Associate Professor & Head, Dental Unit, Somorita Medical College.

4. Dr. Arup Kumar Saha BDS, MPH, Assistant professor City Dental College and Hospital, Dhaka.

5. Dr. Hasibul Hasan Imon BDS, MS, Assistant Professor, Dept of Conservative Dentistry and Endodontics, Udayan Dental College, Rajshahi.

6. Dr. A.S.M Rafiul Haque BDS, Medical Officer, Dept. of Dentistry, BIRDEM, Dhaka

7. Prof. Dr Md. Zakir Hossain BDS, PhD, Professor & Head, Department of Orthodontics, Dhaka Dental College & Hospital

Address of correspondence:

Dr. Md. Ahsan Habib BDS, FCPS, Department of Orthodontics and Dentofacial Orthopaedics, Dhaka Dental College and Hospital

E-mail:ahabib@ymail.com

Introduction

Public Dental Health has been defined as "The science and art of preventing and controlling dental disease and promoting dental health through community effort" by the American Dental Association¹. The unique characteristic of dental diseases is that they are universally prevalent and do not undergo remission or termination if untreated and require technically demanding expertise and time consuming professional treatment. According to G. Dale if deciduous teeth are retained beyond time of exfoliation, they are known to cause delay in eruption of permanent teeth and thus lead to malocclusion and other orthodontic problems, which will then need expensive corrective measures².

More than 400 species of bacteria live in human mouth. Dr. Robert Genco points out that serious gum infection can release bacteria in the blood stream and can worsen the condition of patient suffering from heart disease, stroke and other similar ailments³. It is also known that periodontal diseases can even cause premature labour by release of prostaglandins by periodontal bacteria and also worsen conditions like diabetes and pneumonia.

This study was an attempt to assess the prevalence of four dental diseases-dental caries, gingivitis, retained deciduous teeth & fractured teeth among the patients attending at the Orthodontic department in Dhaka Dental College & Hospital, Bangladesh. Various reports^{4,5,6} have been published about the prevalence & incidence of malocclusion & treatment facilities of Dhaka Dental College & Hospital. However, no study so far had been made over prevalence of dental health problems among the people in our country. So, we did not know about the prevalence & pattern of dental health problems among the patients attending of the Orthodontic department in Dhaka Dental College & Hospital. In a developing country like Bangladesh preventive measures should be the most important concern to mitigate the occurrence of dental health problems so as to avoid complicated and expensive dental treatment. To provide oral health planners in Bangladesh with information on the prevalence & pattern of dental health problems among the patients attending in the Orthodontic department was an attempt to define the target population for orthodontic services in future. It may forecast to initiate a nationwide dental health status & establish nationwide treatment strategy of different dental health problems. Dental caries is the most common dental diseases with high occurrence in humans. It is crucial to control the disease process by assessing and rendering the treatment required along with spreading awareness regarding prevention. Several studies have been conducted and reported on different occasions on dental caries and the treatment needs in developing countries. However, much data is available on the occurrence of dental caries and the treatment needs in children. One of the main etiological factors of the appearance of periodontal diseases is the collection of microbe pad on periodontal edge, bed or recess. Resistance of periodontal webs to microbes depends on many local and general factors. Hyperergical reactions to bacteria belong to the immune mechanisms: anafilaxium and oversensitivity, reactions of immune complexes, and cytotoxic reactions.^{7, 8, 9}

Systematical factors are related to the general health of an organism, they influence periodontitis (e.g. endocrine factors, disorders of nutrition or its insufficiency, effects of medicine, psychological or emotional factors, inheritance, metabolic factors and hematological diseases)^{9,10,11}. Malocclusion is one of the commonest aesthetic problems in Bangladesh.

According to Hossain MZ,⁴ Class I malocclusion patient was 55.2%, Class II was 33.3% (Class II div 1 was 28.8 % and Class II div 2 was 4.5 %), Class III was 8.5% open bite 1.0% and other nonspecific case 2% who are attending Dhaka Dental college in the department of Orthodontics in the year of 1990 - 1991. According to Ahmed N,⁵ Class I malocclusion patient was 45.8% Class II was 40.0% (Class II div-1 was 32.7 % and Class II div 2 was 7.2 %), Class III patients was 14.3% who are attending Dhaka Dental College in the department of Orthodontics in the year of 1992 - 1994. As our general health status indicators are not upto satisfaction & most of the populations are not concerned about oral health, hardly any information was available on the dental morbidity pattern of adolescents in Dhaka. We believed hospital is the best place to contact a large number of teenagers together. It was therefore, decided to carry a hospital based study among 12 to 18 year olds in the Orthodontic department in Dhaka Dental College & Hospital, Bangladesh. The aim was to find out the dental morbidity pattern of adolescents for planning community dental services. This age group was chosen because it is the time of adolescent growth spurt and orthodontic problems are most amenable for correction in this age group¹². The information's was gathered throws light on the current needs in dental care among attending adolescents. Oral health is an essential component of health throughout life. However, millions of individuals suffer from dental caries and periodontal disease, resulting in unnecessary pain, difficulty in chewing, swallowing and speaking, and increased medical costs; hence, the present study was conducted to evaluate the oral health status of Orthodontic patient which would help us in planning and implementing preventive measures¹³.

Materials & Methods

Study design: Descriptive cross sectional study

Place of study: Department of Orthodontics & Dentofacial Orthopaedics of Dhaka Dental College & Hospital, Dhaka.

Sample selection: In this study, a total of 500 patients (250 males & 250 females) were selected from Orthodontic department of Dhaka Dental College & Hospital.

Selection criteria:

Inclusion criteria:

The patients and their parents were Bangladeshi in origin. They were between 12-18 years of age. They were free from any serious illness and have no history of trauma or surgery.

Exclusion criteria:

Non co-operative patients. Patients with systemic illnesses. History of previous orthodontic treatment.

Study procedure:

Each of the patients was selected in respect of inclusion and exclusion criteria. A data collection sheet was used for each patient.

Measurements:

Clinical examinations of each participant carried were followed:

Dental Caries:

The teeth showing discoloration, chalky appearance of enamel, softened enamel

or broken surface by visual examination or probing is defined as carious tooth¹⁴.

The caries Index DMF¹⁵ was developed by Klien, Polemar & Knutson. DMF =D indicates a decayed tooth, M indicates a missing tooth, F indicates a permanently filled tooth due to decay.

Methods:

All teeth except 8's are examined. There are no scoring patterns. In the provided boxes the decayed, missing, filled tooth or surfaces are marked & finally the total counts are made.

Rules:**DMFT-**

1. Primary caries of any surface of tooth comes under decay category.
2. Secondary caries under restoration comes under decayed category.
3. Tooth with temporary restoration comes under decay category.
4. Tooth exfoliated or extracted due to decay comes under missing category.
5. Tooth missed due to periodontal diseases or extraction due to orthodontic purpose doesn't come under missing category.
6. Any tooth with permanent restoration due to dental caries comes under filling category.
7. Root canal treated tooth because of pulpal involvement due to caries comes under filling category.
8. Normal exfoliation during mixed dentition does not come under missing category.

DMFS-

1. Here teeth surfaces are examined. The anterior teeth have 4 surfaces, and posterior has 5 surfaces.
2. Rules of decay category are as same as that of DMFT.
3. When a tooth is completely missing due to caries, while counting the missing surfaces, one surface less is calculated.
4. Filling category rules are as same as that of DMFT.

Gingivitis:

Bleeding from gum on visual examination or bleeding from sulcus on gentle probing & presence of deposits on teeth is diagnosed as gingivitis¹⁶.

The gingival index (GI) was developed by Loe and Silness¹⁷ to describe the clinical severity and location of gingival inflammation using a mouth mirror and periodontal probe, the mesial, distal, buccal and lingual surface of six index teeth examined: maxillary right first molar, maxillary right lateral incisor, maxillary first premolar, mandibular left first molar, mandibular left lateral incisor and mandibular right first premolar.

The scores were defined based on severity from 0-3.

0 = Normal gingival

1 = Mild inflammation-light change in colour, slight edema but no bleeding on probing (BOP).

2 = Moderate inflammation- redness, edema and glazing, and BOP.

3 = Severe inflammation- marked redness and edema, ulceration and tendency to spontaneous bleeding.

The average score of each tooth was calculated by dividing the score of each tooth surface by the number of surfaces examined. The final numerical score per person obtained using the following formula:

Score per person = sum of individual tooth scores / number of teeth examined.

Retained deciduous teeth:

Deciduous tooth is retained beyond the time of exfoliation is diagnosed as retained deciduous teeth¹⁸.

Fractured teeth:

Teeth with broken edges with no obvious evidence of caries are diagnosed as traumatic fractured teeth. This is confirmed by eliciting history of trauma after diagnosing fractured teeth¹⁹.

Types²⁰-

- Infraction of enamel
- Fracture of enamel
- Fracture of enamel & dentine
- Fracture of enamel & dentine with pulp exposure
- Crown-root fracture involving enamel, dentine, cementum & pulp
- Intra-alveolar root fracture
- Concussion
- Sub luxation
- Extrusive luxation
- Lateral luxation
- Intrusive luxation

Orthodontic problems:

Presence of all varieties of malocclusion due to any cause is diagnosed as orthodontic problem. Malocclusion is defined as any deviation from ideal occlusion²¹.

Malocclusions in this study were classified according to Angle classification.

Angle Class –I malocclusion.

Angle Class –II div1 malocclusion.

Angle Class –II div2 malocclusion.

Angle Class –III malocclusion.

Data collection and processing:

Dental examination for 50 patients, were carried out per week. All the patients of the respective age groups were gathered and given introductory health talk. Methods of healthy tooth brushing, and general oral hygiene were taught to all. A small brochure on dental hygiene was given to all. The patients were examined by a single examiner for dental caries, gingivitis, retained deciduous teeth, fractured teeth and orthodontic problems to avoid inter examiner variations. After collection of data the obtained data was checked, verified & edited. These were entered in a personal computer using the SPSS (statistical package for social science) software. Entered data were cleaned, edited and appropriate statistical tests were done depending on the distribution of data.

2. Data analysis:

All data analyzed through standard statistical methods by using SPSS / STATA 10 software.

2. Ethical measures:

The purpose of this study was to know the prevalence & pattern of dental health problems among Bangladeshi people. There were no physical risk of the patients throughout the study period. All patients in the study signed a written informed consent form. No information has been withheld from the patient. No experimental drug of placebo was used.


Result:

Table 1, graph 1:

This was a descriptive study conducted among 500 patients in the department of Orthodontics and Dentofacial Orthopaedics, Dhaka Dental College and Hospital.

Table-1 Distribution the age and sex of surveyed population

Age in years	12yrs	13yrs	14years	15yrs	16yrs	17yrs	18yrs	Total
Male	61	27	10	32	15	34	71	250(50.0)
Female	57	20	31	33	29	23	57	250(50.0)
Total	118(23.6)	47(9.4)	41(8.2)	65(13.0)	44(8.8)	57(11.4)	128(25.6)	500(100)




Graph-1: Distribution of age and sex of patients

Table 2 & 3 and Graph 2 & 3 showed 42.4% had caries and caries rate slightly more in male (42.0%) than female(42.0%) but not significant. Filling 8.6 % , more in female (12.0%) than male (5.0%), Gingivitis 48.4%, more in male (59.2%) than Female (37.6%), retained deciduous 6%, fractured teeth 16.8% , more in male (22.0%) than Female (11.6%), multiple problem 84.6% and others 6.4%. It was observed that the gingivitis was most prevalent condition. This was followed by dental caries, fractured teeth, restored teeth and retained deciduous teeth.

Table-2 Gender wise distribution of dental morbidities

*Dental condition	Total Affected (n=500)	%	Male (n=250)	%	Female (n=250)	%	P-value**
Dental Caries	212	42.4	107	42.8	105	42.0	>0.05
Filling for Caries	43	8.6	13	5.0	30	12.8	<0.01
Gingivitis	242	48.4	148	59.2	94	37.6	<0.001
Retained deciduous teeth	30	6.0	15	6.0	15	6.0	NS
Fractured teeth	84	16.8	55	22.0	29	11.6	<0.001
Multiple problems	423	84.6	225	90.0	198	79.2	<0.001
Other problems	32	6.4	17	6.8	15	6.0	>0.05


*Multiple response
**P < 0.05 is statistically significant



Graph-2 Age Wise distribution of different dental morbidities

Table-3 Showing the Commonest Dental Morbidity Conditions

Disease Conditions	Total no. examined	No. suffering from each condition	Proportion %
Dental caries	500	212	42.4
Gingivitis	500	242	48.4
Filling for Caries	500	43	8.6
Fractured teeth	500	84	16.8



Graph-3 Gender wise distribution of different dental morbidities


Table 4 showed malocclusion (Orthodontic problem). Result showed Class I malocclusion was the most (61.8%) and Class I malocclusion was more in Male (66.8%) than female (56.8%). Class II div-1 was 23% and Class II div-1 malocclusion was more in Female (26.4%) than Male (19.6%). Class II div-2 was 2.2% and Class II div-2 malocclusion was more in Female (2.8%) than Male (1.6%). Class III was 13% and Class III malocclusion was more in Female (14.0%) than Male (12.0%).

Table-4 Type of Malocclusion

Angle's classification	Total	%	Male	%	Female	%	P-value
Class-1	309	61.8	167	66.8	142	56.8	<0.05
Class-II div 1	115	23.0	49	16.6	66	26.4	<0.05
Class-II div 2	11	2.2	4	1.6	7	2.8	>0.05
Class-III	65	13.0	30	12.0	35	14.0	>0.05

Total 500 100.0 250 100.0 250 100.0

*P < 0.05 is statistically significant



Graph-4: Type of Malocclusion

Discussion

Gingivitis was the commonest dental problem encountered. The gingivitis observed in this study was 48.4% and higher in males (59.2%) than in female (37.6%). This finding is in disagreement with Jose²² who found only 15% of gingivitis with higher in female (56%) than in male (44%). Sutcliff survey also shows high prevalence among females.²³ The dental caries observed in this study was 42.4%. Both males and females were almost equally affected by caries with slightly higher rate among males. A study conducted by Jose A and Joseph MR²⁴ in rural Kerala reports 54.3% prevalence of dental caries and both males and females were almost equally affected by caries with slightly higher among males. A study conducted by Sogi G²⁵ in Davangare using DMFT/DMFS score reports higher rate in females that was statistically significant. A study conducted by Cand.Odont. Severre Aukland and Cand. Odont. Johny Bjelkaroe report 45.8% of dental caries. Findings of this present study are in agreement with Jose A and Johny Bjelkaroe. The rate of fractured tooth observed in this study was 16.8%, and higher in male (22.0%) than in female (11.8%). Finn also reports higher incidence of fractured teeth among boys. This may well be explained by the more aggressive eating habits of the boys²⁶. Also chances of fights and falls are commoner among them which also contribute to fractured teeth.¹⁴ Among 500 male and female subjects only 8.6 had dental filling and among of all those subjects more females (12.8%) have dental fillings than males (5.2%). Jose A²² also reports 3.18% dental fillings and higher in males which disagreed with the present study. As age advances retained deciduous teeth become less but still 6% Subjects had retained deciduous teeth males and females were equally affected. Jose A found 7% retained deciduous teeth.³¹ Present study found Class I malocclusion the most (61.8%) and it was more in Male (66.8%) than female (56.8%). Class II div-1 was 23% and Class II div-1 malocclusion was more in Female (26.4%) than Male (19.6%). Class II div-2 was 2.2% and Class II div-2 malocclusion was more in Female (2.8%) than Male (1.6%). Class III was 13% and Class III malocclusion was more in Female (14.0%) than Male (12.0%).

Limitation of study:

1. The study group was selected from Dhaka Dental College & Hospital. So the findings might be specific area, which may not represent the whole national situation.
2. The size of the sample of the study was limited in relation to the great number of populations in Bangladesh to represent the situation prevailing nationality.

Conclusion

Good oral health is essential to improve individual overall health & well-being. We urge to take this information & use it for program planning & advocating for the health of patients. It is only through working together that we can make excellent oral health a reality for these patients. Since the number of orthodontists available to treat patients in Bangladesh is limited, there is a high demand on each practitioner. Therefore current orthodontic students should receive more education & training in the management of malocclusion to improve the overall quality of care for patients.

Recommendation

As the size of the sample of this study is limited in relation to the number of people in Bangladesh, recommendation is put forward for future researcher to do additional depth research consisting of large sample group for more acceptability of the study.

References:

- Peter S.: Essentials of Preventive and Community Dentistry. 1st ed. New Delhi; Arya Medi publications 1999; 8.
- G. Dale. Guidance of Occlusion: Serial Extraction. In Graber T.M., Swain.B.F. eds. Orthodontics Current Principles and Technique. St. Louis: C. V. Mosby Company, 1985; 284 - 95.
- Genco R., Offenbacher S., Beck. J.: Periodontal disease and cardiovascular disease. Journal of American Dental Association 2002; 133: 14s - 21s.
- Hossain MZ, Haque S, Yasmin S, et al. Prevalance of malocclusion and treatment facilities at Dhaka Dental College and Hospital. J of Oral Health. 1994; 1:4-6.
- Niaz A, Kamran C. Prevalance of malocclusion and its aetiological factors. J of Oral Health 1996; 2:12-16.
- Study of malocclusion in Dhaka Dental College; An epidemiological overview by Sattar MH, Khaleque KA, Haq ME; 7th Bangladesh National & 3rd SADAF Dental Conference, 1995; Presentation B-7; Session 7.
- Sheiham A. In the chemical prevention of gingivitis necessary to prevent severe periodontitis. Periodontol 2000. 1997; 15:15-24.
- D'Almeida HB, Kagami N, Maki Y, Takaesu Y. Self-reported oral hygiene habits, health knowledge, and sources of oral health information in a group of Japanese junior high school students. Bull-Tokyo-Dent-Coll 1997; 38(2): 123-31.
- Ashley FP, Usinskin LA, Wilson RF. The relationship between irregularity of the teeth, plaque, and gingivitis. Eur J Orthod 1998; 20(1): 65-72.
- Tae Ju-Oh, Robert Eber, Hom Lay-Wang. Periodontal diseases in the child and adolescent // J Clin Periodontol 2002; 29 (5): 400-10.
- Milčiuvienė S, Jasulaitytė L. Prevention of stomatological diseases; Kaunas 1999. p. 31-71.
- Bhalajhi S. I., Orthodontics - The Art and Science. 1st ed. New Delhi; Arya (Medi) Publishing house 1999: 10.
- Das UM, Beena JP, Azher U. Oral health status of 6- and 12-year-old school going children in Bangalore city: an epidemiological study. J Indian Soc Pedod Prev Dent. 2009 Jan-Mar;27(1):6-8.
- Lundensen T. F., Roberson T. M., Cariology, the lesion etiology, prevention and control. In Sturdevant. C.M., Roberson T.M., Heymann.H.O., Sturdevant J.R. eds.: The Art and Science of Operative Dentistry Singapore: Harcourt Brace & Company Asia PTE Ltd, 1997; 102 - 103.
- Debnath T., Ashok's Public Health & Preventive Dentistry, 2nd Edition: 2007; P-58-59.
- Carranza F. A.: Clinical Features of Gingivitis. In Carranza, Newman eds Clinical Periodontology Bangalore: Prism Books PVT LTD., 1996; 223-224.
- Loe H, Silness J. Periodontal disease in pregnancy. I. prevalence and severity. Acte Odontol Scand 1963; 21: 533-51.
- Tencate A. R.: Shedding of Deciduous Teeth. In Bhaskar S.N. Eds. Orban's Oral Histology and Embryology, 10th ed. St. Louis. Toronto. Princeton: C.V. Mosby Company, 1986; 392 - 393.
- McDonald R.E., Avery D.R., Lynch T.R.: Management of Trauma to the Teeth and Supporting Tissues. In McDonald R.E., Avery V.R. eds. Dentistry for the Child and Adolescent New Delhi; Laxmanchand Arya, 1998; 512.
- Pitt Ford T.R., Endodontics in Clinical Practice. 5th Edition. 2009; P197.
- Profit W. R., Ackerman J. L. Diagnosis and Treatment Planning in .Orthodontia. In Graber T. M., Swain. B. F. eds. Orthodontia. In Graber T. M., Swain B. F. eds. Orthodontics Current Principles and Techniques ST. Louis: C.V. Mosby Company, 1985; 5.

22. JOSE A and JOSEPH M R. Prevalence of dental health problems among school going children in rural Kerala.
23. McDonald R. E., Avery D. R., Weddell J. A.: Gingivitis and Periodontal Disease. In McDonald R. E, Avery V. R. eds.:Dentistry for the Child and Adolescent, 5th ed. New Delhi:Laxmanchand Arya, 1998; 478.
24. Sogi G., Bhaskar D.J. Dental Caries and Oral Hygiene Status of 13 to 14 year Old School Children of Davangere. J Indian Soc Pedo Prev Dent 2001; 21: 113 - 116.
25. Cand.Odont. Severre Aukland and Cand.Odont. Johny Bjelkaroey. The Dental Health of School Children in Betul District Madhya Pradesh. J Indian Dent Assoc 1982; 54: 367 -369.
26. Finn. Clinical Pedodontology. 4th ed. New Delhi; AITBS 1998: 225.