

Article

Effect of seed rate on biomass production of Dhaincha (*Sesbania aculeata*)

Md. Hamidur Rahaman¹, Sultan Ahmed¹, Swadesh Chandra Samanta¹ and S.M. Ahsan²

¹Department of Agronomy, Faculty of Agriculture, Patuakhali Science and Technology University, Patuakhali, 8602, Bangladesh

²Department of Agriculture, Faculty of Life Science, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj 8100, Bangladesh

*Corresponding author: S.M. Ahsan, Department of Agriculture, Faculty of Life Science, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj 8100, Bangladesh. Phone: +8801763818889; E-mail: smvahsan@gmail.com

Received: 07 August 2016/Accepted: 24 August 2016/ Published: 31 August 2016

Abstract: A field experiment was conducted at the Agronomy field, Patuakhali Science and Technology University (PSTU), Patuakhali during the period of May 2013 to August 2013. The experiment was conducted to find out the effect of seed rate on nodulation of dhaincha (*Sesbania aculeata*) under non-saline agro-ecosystem. The experiment comprised of seven different seed rates viz. 30, 35, 40, 45, 50, 55 and 60 kg ha⁻¹. The experiment was laid out in randomized complete block design with three replications. The seed rates had significant influence on different parameters of dhaincha plant. It was found that (30 kg ha⁻¹) recorded the tallest plant (4.11 m) followed by (45 kg ha⁻¹) while (60 kg ha⁻¹) gave the smallest plant (3.66 m). Recorded maximum plant diameter was (0.92 cm) at harvest while the lowest plant diameter was obtained from (55 kg seed ha⁻¹) (0.79). The highest plant population (336.0) was found in (60 kg ha⁻¹) while (30 kg ha⁻¹) showed the lower plant population (114.0). The maximum number of branches (23.93 plant⁻¹) at harvest while (60 kg ha⁻¹) produced the lowest number of branches plant⁻¹ (7.93 plant⁻¹). It was recorded that (60 kg ha⁻¹) produced the highest fresh weight m⁻² (4.10 kg) and was statistically different from other treatments. On the other hand, (30 kg ha⁻¹) produced significantly the lowest fresh weight of plants m⁻² (1.93 kg) which was also statistically different from other treatments. The higher below ground biomass production m⁻² (1.00 kg) was found in (30 kg ha⁻¹) followed by (35 kg ha⁻¹), (40 kg ha⁻¹) and (45 kg ha⁻¹) while (50 kg ha⁻¹) observed the lower below ground biomass production m⁻² (0.19 kg) at (60 kg ha⁻¹). Production of pods branch⁻¹ was maximum (22.00) at (30 kg ha⁻¹) followed by (35 kg ha⁻¹) while the lowest number of pods branch⁻¹ (17.07) at (60 kg ha⁻¹). It was found that (30 kg ha⁻¹) produced the maximum number of seeds pod⁻¹ (32.73) which was significantly different from other treatments. On the other hand, (60 kg ha⁻¹) recorded the significantly minimum number of seeds pod⁻¹ (25.40) which was also statistically similar with (60 kg ha⁻¹).

Keywords: seed rate; biomass; dhaincha; *Sesbania aculeata*

1. Introduction

Dhaincha (*Sesbania* spp.) belongs to the family Leguminosae (sub-family Papilionoideae), and is well known for its diversified use in Bangladesh. Three species of *Sesbania* viz. *S. sesban* (L.) Merr., *S. bispinosa* (Jacq.) Wight and *S. cannabina* (Retz.) Poir., are commonly known as dhaincha in Bangladesh (Prain, 1903; Ahmed *et al.*, 2009). It is an ideal green manure crop as it is quick-growing, succulent, easily decomposable with low moisture requirements, and add maximum amounts of organic matter and nitrogen in the soil (Palaniappan and Siddeswaran, 2001). Long time (10/12 yrs) cultivation of dhaincha would combat desertification of marginal lands, e.g., char land, saline area, etc. and rehabilitate degraded lands into productive crop lands for intensive food crop agriculture (Carroll and Somerville, 2009). Dhaincha also showed its potentiality as a raw material for

paper pulp (Jahan *et al.*, 2009). Researchers, recently, have found that the leaves of dhaincha are a good source of Pinitol, an anti-diabetic compound (Misra and Siddiqui, 2004). Nevertheless, it would not bring desirable benefit to many of the less favored areas where adequate fertilizers are not available and soils are very deficient in nutrients, especially N and P (Basu and Kabi, 1987). As a legume crop, dhaincha (*Sesbania aculeata*) has the unique ability of fixing and utilizing atmospheric nitrogen for its growth and enriching soil fertility as well as increasing the status of organic matter content of the soil. Such fixation of atmospheric nitrogen takes place through a biological process *i.e.* symbiotic association between *Sesbania* root and rhizobium bacteria. Consequently, the use of bio fertilizers has increasingly been gaining global attention as one of the practices to restore and maintain soil fertility (Bhuiya *et al.*, 1995). In third world countries, fertilizer nitrogen is applied for cultivation of only a few cash crops and main dietary crops like rice and wheat. The nitrogen fixation potential of *Sesbania aculeata* has already been established and estimated as 176 kg N ha⁻¹ in a period of 56 days (Furoc *et al.*, 1985). In addition, several workers (Tiwari *et al.*, 1980; Furoc *et al.*, 1985) have documented increased nodulation and high dry matter accumulation due to rhizobium inoculation. However, research work on the seed inoculation of *Sesbania* for effective nodulation, growth and biomass production and N₂-fixation received less attention in our country and nevertheless, it is necessary to identify the effective strains in terms of nodulation and nitrogen fixation ability of the host crop. Biological nitrogen fixation by tree legumes is important in relation to global environment. In the context of afforestation of marginal land, and nodulated nitrogen fixing tree legumes have a special advantage over other tree species. Wide ranges of leguminous N₂-fixing trees are utilized in forestry, agroforestry and land reclamation (Dommergues, 1993). If the planting density is lower than its optimum value then total production will be lower and weeds will be higher (Allard, 1999). In addition, increasing planting density will also increase intra-specific competition among the plants, which affects the vegetative and reproductive growth of plants (Zhang *et al.*, 2006). Higher planting density also increases the relative humidity within canopy and leaf wetness by reducing air movement and sunlight penetration within canopy (Burdon and Chilvers, 1982; Tu, 1997). Thus, plant density could have significant impact on plant disease incidence (Burdon and Chilvers, 1982; Copes and Scherm, 2005). Reduction in seeds yield may be the result of lower number of pods, lower seeds weight or a combination of these components. In dense populations, many seeds may not develop. Jettner *et al.* (1999) stated that increasing yields at high sowing rate could be directly attributed to large plant population. There was strong relationship between economic optimum plant density and seed yield potential (Regan *et al.*, 2003). It is, therefore, necessary to determine the optimum density of plant population per unit area for obtaining maximum yields, for which it is important to know the effect of increasing planting densities on plant yield and biomass production. Keeping the above points in view, the present study was undertaken with the objective is to assess the optimum seed rate and the effect of seed rate on nodulation of dhaincha (*Sesbania aculeata*).

2. Materials and Methods

2.1. Description of the experimental site

2.1.1. Site selection

The experiment was conducted at the Patuakhali Science and Technology University farm during kharif season. The maximum area is covered by Gangatic tidal floodplains and falls under Agro ecological Zone "AEZ 13". The area lies at 0.9 to 2.1 meter above mean sea level. This region occupies a vast area of tidal floodplain land in the north-west part of Patuakhali district.

2.1.2. Soil

Soil characteristics of the Patuakhali Science and Technology University farm are silty loams or alluvium. However, the soil of the experimental field was silty clay loam having pH value of 6.8. The characteristics of experimental soil was examined by the soil resource development institute (SRDI), regional laboratory, Barisal. The organic carbon content found 0.93% in most cases. Deficiency of nitrogen is acute and widespread. Status of exchangeable potassium is almost satisfactory. Phosphorus, Sulphur and other characteristics of soil status are also satisfactory.

2.2. Experimental crop

The crop used in the study was dhaincha (*Sesbania aculeata*). The seeds were obtained from the agronomy Field Laboratory, department of agronomy, Patuakhali Science and Technology University. The seeds were healthy, vigorous and well matured. The germination percentage of seeds was 90.

2.3. Lay out of the experiment

The experiment was laid out in a randomized complete block design with three replications. Each replication represents a block. Each block was subdivided into seven unit plots. The treatments were randomly distributed to the unit plots in each block. Total numbers of plots were 21 and the unit plot size was $4\text{m} \times 2.5\text{m} = 10\text{m}^2$. Plot to plot distance was 0.5 m and block-to-block distance was 1.00 m.

2.4. Treatments

There were seven treatments with different seed rates. Treatments were as follows:

1. T₁: 30 kg seed ha⁻¹
2. T₂: 35 kg seed ha⁻¹
3. T₃: 40 kg seed ha⁻¹
4. T₄: 45 kg seed ha⁻¹
5. T₅: 50 kg seed ha⁻¹
6. T₆: 55 kg seed ha⁻¹
7. T₇: 60 kg seed ha⁻¹

2.5. Experimental details

2.5.1. Land preparation

The land was prepared by ploughing with the help of a power tiller. Laddering was done properly after each ploughing for breaking the clods and leveling the land. Weeds, stubbles and crop residues were removed from the field before final ploughing and leveling.

2.5.2. Sowing of seeds

Seeds were broadcasted on 22 May 2013 before 8 PM and were covered by soils soon after seeding.

2.5.3. Germination of seeds

Germination of seeds started from 3 DAS and continued up to 10 DAS. High percentage of germination was more than 80 and on the 11th day, nearly all plants came out of the soil.

2.5.4. Intercultural operations

2.5.4.1. Weeding

The experimental plots were infested by some weeds which were uprooted from the field twice at 15 and 30 days after sowing.

2.5.4.2. Application of fertilizers and insecticides

During land preparation TSP, MoP and DAP were applied at the rate of 45 kg ha⁻¹, 30 kg ha⁻¹, 35 kg ha⁻¹ which mixed with soil properly to improve soil properties. The crop was infested with insect which was controlled by applying Sevin 60 EC insecticide once.

2.5.5. Data collection

Five plants were randomly selected from each plot to record nodulation, shoot growth, root growth and total growth character at 30, 45 and 60 days of sowing. Plant samples were collected from the plots before maturation of seed. From each plot, five plants were carefully uprooted with soil, so that no nodule was left in the soil. The collected nodulated plants were washed in a slow stream of tap water and were finally cleaned with a soft camel hairbrush to ensure the removal of soil particles adhering to the root surface. The nodules from the root system of each plant were separately collected and counted. The shoot portion of each plant was then separated from the root and the length of each plant was recorded. The root, shoot and nodules were air-dried and then oven dried at 65°C for 72 hours. The oven dry weight of roots, shoots and nodules were recorded and the shoot and root samples were stored for analysis. Shoot growth, root growth and total growth obtained from each plot after final harvest were dried and weighed carefully.

2.6. Harvesting

Harvesting was started when 80% of the plant population of each plot reached to the seed maturity and showed physiological maturity symptoms.

2.7. Statistical analysis

Data obtained from the field experiment were analyzed statistically by the method of analysis of variance of Gomez and Gomez (1984). The treatment means of different characters were compared following duncan's new multiple range test (DMRT).

3. Results and Discussion

3.1. Effect of seed rate

3.1.1. Plant height

Different seed rates had significant effect on plant height of dhaincha at harvest. It was found that T₁ (30 kg ha⁻¹) recorded the tallest plant (4.11 m) followed by T₄ (45 kg ha⁻¹) while T₇ (60 kg ha⁻¹) gave the smallest plant (3.66 m) (Figure 1). These results revealed that 30 kg ha⁻¹ was more efficient than other seed rate regarding plant height. Due to lower plant population which got more nutrient, spacing and sunlight. Similar result was also obtained by Zhang *et al.* (2006) in case of soybeans. Figure followed by same letter(s) are statistically similar as per DMRT at 5%.




Figure 1. Effect of seed rate on plant height ($S_{\bar{X}} = 0.063$).

3.1.2. Plant diameter

Effect of different seed rate significantly influenced plant diameter at harvest where T₁ (30 kg ha⁻¹) produced significantly the maximum plant diameter (0.92 cm) at harvest while the lowest plant diameter was obtained from T₆ (55 kg seed ha⁻¹) (0.79) (Figure 2). Similar study was also done by Venkanna (2014) who reported similar result in case of dhaincha. Figure followed by same letter(s) are statistically similar as per DMRT at 5%.




Figure 2. Effect of seed rate on plant diameter ($S_{\bar{X}} = 0.037$).

3.1.3. Plant population

Plant population showed significant variation due to different seed rates (Table 1) where significantly the highest plant population (336.0) was found in T₇ (60 kg ha⁻¹) while T₁ (30 kg ha⁻¹) showed the lower plant population (114.0). This result indicated that higher seed rate produced the maximum plant population than lower seed rate which ultimately resulted in higher production of biomass. Similarly, Morrisson *et al.* (1990) reported that significantly the highest plant population was obtained in higher seed rate in summer rape.

Table 1. Effect of seed rate on plant population and number of branches plant⁻¹

Treatments	Plant population (m ⁻²)	Number of branches plant ⁻¹
T ₁	114.0 g	23.93 a
T ₂	128.0 f	11.93 b
T ₃	153.0e	11.33 bc
T ₄	185.7 d	11.13 bc
T ₅	228.3 c	10.87 cd
T ₆	278.0 b	10.07 d
T ₇	336.0 a	7.933 e
$S_{\bar{x}}$	2.143	0.303
CV (%)	1.83	4.21
Level of significance **		**

** denotes significant at 1% level of significance, significant at 5% level of significance and non-significant, respectively. Figures followed by same letter(s) are statistically similar as per DMRT at 5%.

3.1.4. Number of branches plant⁻¹

Effect of different seed rates significantly influenced number of branches plant⁻¹ at harvest where T₁ (30 kg ha⁻¹) produced significantly the maximum number of branches (23.93 plant⁻¹) at harvest while T₇ (60 kg ha⁻¹) produced the lowest number of branches plant⁻¹ (7.93 plant⁻¹) (Table 1). These results indicated that, number of branches plant⁻¹ was gradually decreased due to the decrease in seed rate. Similar study was obtained by Regan *et al.* (2003) who found that the lower crop density performed better than higher crop density regarding number of branches plant⁻¹ in chickpea.

3.1.5. Fresh weight of plants m⁻²

Effect of different seed rate was significant at 1 % level of significance for fresh weight of plants m⁻² (Table 2). Treatment T₇ (60 kg ha⁻¹) produced the highest fresh weight m⁻² (4.10 kg) and was statistically different from other treatments. On the other hand, T₁ (30 kg ha⁻¹) produced significantly the lowest fresh weight of plants m⁻² (1.93 kg) which was also statistically different from other treatments. These results revealed that higher seed rate produced higher fresh weight than that of lower seed rate. This might be due to the fact that higher plant population enhanced fresh weight production. Similar study was obtained by Regan *et al.* (2003) who found that the lower crop density performed better than higher crop density regarding fresh weight in chickpea. Tabib and mortuza (2012) reported that *Sesbania rostrata* produced 10.69 to 28.25 t ha⁻¹ of 30, 35, 40, 45, 50, 55 and 60 kg ha⁻¹ planting density. The finding of the present study lies within the range presented above.

3.1.6. Below ground biomass production m⁻²

Below ground biomass production m⁻² showed significant variation due to different seed rates at 5 % level of significance (Table 2) where significantly the higher below ground biomass production m⁻² (1.00 kg) was found in T₁ (30 kg ha⁻¹) followed by T₂ (35 kg ha⁻¹), T₃ (40 kg ha⁻¹) and T₄ (45 kg ha⁻¹) while T₅ (50 kg ha⁻¹) observed the lower below ground biomass production m⁻² (0.19 kg) at T₇ (60 kg ha⁻¹). This result indicated that 30-50 kg ha⁻¹ produced the maximum below ground biomass production m⁻² than other seed rate which ultimately resulted in higher production of crop. Venkanna *et al.* (2013) found the same result in case of dhaincha.

Table 2. Effect of seed rate on fresh weight m⁻², dry weight of nodule plant⁻¹ and below biomass ground m⁻².

Treatments	Fresh weight of plant (kg m ⁻²)	Dry weight of nodules plant ⁻¹ (g)	Below ground biomass (kg m ⁻²)
T ₁	1.93 f	5.59 a	0.19 d
T ₂	2.31 e	5.263b	0.33 cd
T ₃	2.68 d	5.18 b	0.57 bcd
T ₄	3.00 c	4.48 c	0.58 abcd
T ₅	3.17 c	3.93 d	0.68 abc
T ₆	3.50 b	3.79 de	0.78 ab
T ₇	4.10 a	3.61 e	1.00 a
$S_{\bar{x}}$	0.077	0.080	0.128
CV (%) 4.52 3.01 37.62	CV (%) 4.52 3.01 37.62	CV (%) 4.52 3.01 37.62	CV (%) 4.52 3.01 37.62
Level of significance	**	**	**

** denotes significant at 1% level of significance, significant at 5% level of significance and non-significant, respectively.

3.1.7. Number of pods branch⁻¹

Number of pods branch⁻¹ was significantly affected by the effect of different seed rates at harvest. Production of pods branch⁻¹ was maximum (22.00) at T₁ (30 kg ha⁻¹) followed by T₂ (35 kg ha⁻¹) while the lowest number of pods branch⁻¹ (17.07) at T₇ (60 kg ha⁻¹) (Figure 3). Similarly, Regan *et al.* (2003) found the significant differences among the seed rates in chickpea. Figures followed by same letter(s) are statistically similar as per DMRT at 5%.



Figure 3. Effect of seed rate on number of pods branch⁻¹ ($S_{\bar{x}} = 0.717$).

3.1.8. Number of seeds pod⁻¹

A highly significant variation was found due to the effect of seed rate in respect of number of seeds pod⁻¹ at harvest. It was recorded that T₁ (30 kg ha⁻¹) produced the maximum number of seeds pod⁻¹ (32.73) which was significantly different from other treatments (Figure 4). On the other hand, T₇ (60 kg ha⁻¹) recorded the significantly minimum number of seeds pod⁻¹ (25.40) which was also statistically similar with T₆ (60 kg ha⁻¹). This result indicated that T₁ (30 kg ha⁻¹) was more efficient to produce more seeds. Similarly, Jettner *et al.* (1999) found the significant differences among the seed rate in desi chickpea. Figures followed by same letter(s) are statistically similar as per DMRT at 5%.




Figure 4. Effect of seed rate on number of seeds pod⁻¹ ($S_{\bar{x}} = 0.418$).

4. Conclusions

The research work was conducted at the Agronomy field, Patuakhali Science and Technology University, to investigate the effect of different seed rates on growth, nodulation, and yield, and yield performance of dhaincha during the period from May 2013 to August 2013. The research work consisted of seven treatments viz. 30, 35, 40, 45, 50, 55, 60 kg ha⁻¹. The experiment was laid out in randomized complete block design with three replications and the MSTAT-C package program did analyses whereas means were adjudged by DMRT (Duncan's new multiple range test) at 5% level of probability. Data were recorded on various growth and yield attributing traits. Seed rate had significant influence on growth, yield and yield contributing characters. However, seed rate 30kg ha⁻¹ recorded significantly the tallest plant (4.11 m). Among the other characters higher below ground biomass production m⁻² (1.00 kg) was found at the seed rate of 30 kgha⁻¹ followed by seed rate 35kgha⁻¹(0.78), seed rate 40kgha⁻¹(0.68) and seed rate 45 kgha⁻¹ (0.58). Maximum number of pods branch⁻¹

¹(22.00), and 30 seeds pod⁻¹ (32.73) and branches plant⁻¹ (23.95) and plant population (114) was found at the seed rate of 30 kg⁻¹. From the above result investigation, it is clear that under agro-ecological region of Patuakhali (AEZ 13), seed rate influenced the growth and yield attributes of dhaincha. 30 kg ha⁻¹ performed the best over other treatments in respect of seed production. Therefore, seed rate 30 kg ha⁻¹ would be appropriate for the seed production. However, further study is needed to confirm the present result.

Conflict of interest

None to declare.

References

- Ahmed ZU, MA Hassan, ZNT Begum, M Khondker, SMH Kabir, M Ahmad, and ATA Ahmed, 2009. Encyclopedia of Flora and Fauna of Bangladesh, Angiosperms: Dicotyledons (Fabaceae–Lythraceae). Asiatic Soc. Bangladesh, 8: 1-474.
- Allard RW, 1999. History of plant population genetics. Annual of Review of Plant Genetics, 33: 1-27.
- Basu PK and MC Kabi, 1987. Effect of application of bio fertilizers on the growth and nodulation of seven forest legumes. Indian-Forester. Silvicultural Div. (South), Midnapore, West Bengal, India, 113: 249- 259.
- Bhuiya, MSA, MA Kashem, SMA Hossain and MS UIalukder, 1995. Effect of green manuring with *S. rostrata* in presence of urea-N on transplanted amanrice. Bangladesh J. Agri. Sci., 2: 677-679.
- Burdon JJ and GA Chilvers 1982. Host density as a factor in plant diseased ecology. Annual Review of Phytopathology, 20: 143-166.
- Carroll A and C Somerville, 2009. Cellulosic biofuels. Ann. Rev. Plant Biol., 60: 165-182.
- Copes WE and H Scherm, 2005. Plant spacing effects on microclimate and *rhizoctonia* web blight development in container-grown Azalea. Horti. Sci., 40: 1408-1412.
- Dommergues Y, G Rinaudo and B Dreygus, 1993. *S. rostrata* green manure and the nitrogen content of rice crop and soil. Soil Biol. Biochem., 15: 111-113.
- Furoc RE, MA Dizon, RA Morris and EP Marqueses, 1985. Effect of flooding regimes and planting dates to N accumulation of three *Sesbania* species and consequently to transplanted rice. Paper presented the 16th annual sci. convention of the crop sci. soci. of Philippines, 10 May 1985, central Luzon State uni. Munoz, Nueva Ecija, Philippines.
- Jahan MS, S Rawshan, T Banu, DNA Chowdhury, A Noori and A Al-Maruf, 2009. Effect of harvesting age on the chemical and morphological properties of dhaincha (*Sesbania aculeata*) and its pulpability and bleachability. BioResources, 4: 471-481.
- Jettner RJ, KH Siddique, SP Loss and RJ French, 1999. Optimum plant density of desi chickpea (*Cicer arietinum* L.) increase with increasing yield potential in south western Australia. Australian J. Agri. Research, 50: 1017-1026.
- Misra LN and SA Siddiqui, 2004. Dhaincha (*Sesbania bispinosa*) leaves: A good source of antidiabetic (+)-pinitol. Curr. Sci., 87: 10.
- Morrisson J, EBE Mcvetty and R Scarth, 1990. Effect of altering plant density on growth characteristics of summer rape. Canadian J. Plant Sci., 70: 13-49.
- Palaniappan SP and K Siddeswaran, 2001. Regional overview on green manure in rice-based cropping systems. In: Gowda CLL, A Ramakrishna, OP Rupela and SP Wanieds. Legumes in rice based cropping systems in Tropical Asia: constraints and opportunities. Int. Crop Res. Inst. Semi-Arid Trop., Patancheru (India), pp. 126-135.
- Prain D, 1903. Bengal Plants, Indian Rep., 1981, B. Singh and M.P. Singh, India.
- Regan KL, KH Siddique and LD Martin, 2003. Response of Kabuli chickpea (*Cicer artietinum*) to sowing rate in Mediterranean – type environments of southwestern, Australia. J. Expt. Agri., 43: 87-97.
- Tabib FAI and MGG Mortuza, 2012. Growth and biomass production of relayed *S. rostrata* as influenced by its planting geometry and age of boro rice seedlings. J. Expt. Biosci., 3: 27-32.
- Tiwari KN, AN Pathak and H Ram, 1980. Green manuring in combination with fertilizer N on rice under double cropping system in an alluvial soil. J. Indian Soc. Soil Sci., 28: 162-169.
- Tu JC, 1997. An integrated control of white mold (*Sclerotinia sclerotiorum*) of beans, with emphasis on recent advances in control. J. Bot., 38: 73-76.
- Venkanna B, 2014. Influence of row spacing and phosphorus levels on seed production of dhaincha (*S. aculeata*). 2nd Int. conference on agri. and hortic. sci. February, 03-05, 2014 Hyderabad, India.
- Zhang W, H Zhang and HWang, 2006. Effects of spacings and planting densities on agronomic traits and yield in high oil soybeans. Soybean Sci., 25: 283- 287.