

PHYTOCHEMICAL COMPOUNDS OF *LILIUM LEDEBOURII* BLOSS USING BULB EXPLANTS

SINA GHANBARI^{1*}, BARAT ALI FAKHERI AND NAFISEH MAHDINEZHAD

*Department of Plant Breeding and Biotechnology, Faculty of Agriculture,
University of Zabol, Zabol, Iran*

Keywords: *Lilium ledebourii* Bloss, Superoxide dismutase, Peroxidase, Flavonoid, Antioxidant, DPPH, Total Phenol.

Abstract

Lilium ledebourii Bloss is a wild species of *Lilium*, which grows naturally in Damash Mountains of Roodbar, Iran. Secondary metabolite synthesis in plants is one of the most important defensive mechanisms applied against various pathogens, and the quality and quantity of those materials depend on the species diversity, organs, habitat conditions, and different extraction methods. In this study, phytochemical compounds were measured quantitatively and qualitatively. Results showed the existence of secondary metabolites in the regenerated plants. Phenolic content was 5.198 ± 0.17 mg in each extract. Radical trapping of the diphenyl-picryl hydrazyl test was 3.24 ± 0.52 mg/ml. Antioxidant activity based on reduced iron was 25.88 ± 1.47 mmol Fe^{+2} /g DW. Flavonoid content of *L. ledebourii* Bloss was 0.78 ± 0.07 mgRE/G. Peroxidase and superoxide dismutase's enzymes activity in *L. ledebourii* Bloss was low. Also, in this study, the combination of saponin, cardiac glycosides, Steroids, Alkaloids and Terpenoid was seen qualitative in the *L. ledebourii* Bloss.

Introduction

Nature is a rich source of medical metabolites and some of them are found in plants. Extracts, which are rich in flavonoid metabolites, through decreasing oxidative stress, protect cells (Maber and Hanneken 2005). Phenols and polyphenols are found in many foods and have great antioxidant effects (Bouayed *et al.* 2011). Extracts, which are rich in flavonoid metabolites, through decreasing oxidative stress, protect cells (Maber and Hanneken 2005). Phenols and polyphenols are found in many foods and have great antioxidant effects (Bouayed *et al.* 2011). In developing countries, due to the side effects of chemical medicines, use of herbal plants in treating various diseases and use of natural antioxidants in food industries have been received much attention, and studies are increasingly being conducted on the extraction of active biological metabolites of plants (Asghari and Mazaheri-tehrani 2010). Plant value is most important because of the production and performance of its secondary metabolites (Akinmoladun *et al.* 2007), and some of the most important metabolites are alkaloids, flavonoids, tannins, saponins, terpenoid and phenolic compounds (Edeoga *et al.* 2005). As plants are important source of antioxidants, the studies in this field have been increased gradually. Those plants which are rich in antioxidant metabolites can protect cells from oxidative dangers (Kumaran and Karunakaran 2006). Since the different metabolites act based on the different mechanisms, it is obvious that only one method cannot provide a comprehensive prediction of the effects of all involved issues of antioxidant characteristics (Pellegrini *et al.* 2000).

Some of the well-known methods of measuring antioxidative features, are radical entrapping of 1,1-diphenyl-2-picrylhydralise (DPPH, based on the hydrogen endowing ability of extract) (Miliauskas *et al.* 2004), the study of inhibition of lipoic acid oxidation (BCB), and the evaluation

*Author for correspondence: <sina_qanbari@yahoo.com> or <sina.qanbari@gmail.com>. ¹Plant Bank, Iranian Biological Resource Center (IBRC), ACECR, Tehran, Iran

of antioxidant activity of extracts is based on the reduction ability of divalent iron (Benzie and Strain 1999). In fact it shows an estimation of antioxidative characteristics of soluble metabolites in water. FRAP method evaluates the antioxidative activity of extracts based on iron reduction ability (Deepa *et al.* (2007). It also shows an estimation of antioxidative characteristics of soluble metabolites in water. Gardner *et al.* (2000) showed that ascorbic acid content has a direct relationship with antioxidant metabolites of extract in FRAP method. The refining activity of existing free radicals in the extract is studied by DPPH method, and Velioglu *et al.* (1998) reported that it had a linear relationship with existing phenolic and flavonoid metabolites in the plant. Phenolic metabolites are a group of compounds which exist in all plants such as fruits, vegetables, and cereals. These compounds are secondary metabolites of plants (Taiz and Zeiger 2002). Naturally, about 8000 different phenolic compounds with the characteristics such as involvement in making NDF, the defensive mechanism of the plant, and fruit characteristics like color, flavor, taste, exist in plants. In addition, phenolic compounds are considered as indexes for identifying physiological stages during fruit growth (Macheix *et al.*1990). The content of plant secondary metabolites such as, pure flavonoids, and their antioxidative characteristics depends on the different factors such as weather, species, extraction method, and antioxidant measuring method (Kumar and Neelam 2006). Flavonoids and other phenolic compounds are widespread in plants and previously various biological activities of these compounds such as antioxidative, antibacterial, and anti-inflammatory characteristics have been studied by (Jamshid *et al.* (2010).

Environmental tensions through increasing production of different types of active oxygen and protein-DNA, induce some harms to vital bio molecules such as lipids. In order to decrease negative effects of various active oxygens in cells, plant's cells apply different antioxidative systems which include an enzyme and non-enzyme factors such as superoxide dismutases (SOD), catalase (CAT), and peroxidase (POD) (Singh *et al.* 1994). Enzyme activities are the most important factors in plant resistance to environmental conditions. Superoxide dismutases as the key antioxidant mainly in the cytosol, chloroplast, mitochondria, and peroxisome attracted much attention of the scientists as it catalyzes radical superoxidase to hydrogen peroxidase (Singh *et al.* 1994). Therefore, SOD is considered as one of the important components of the defensive mechanism.

Lilium ledebourii Bioss is a species of *Lilium* family and it is an herbaceous and permanent plant. This plant has uncovered bulbs of 50 to 150 cm height. It has linear and dagger-like leaves which are 10 to 24 cm long and 1 to 2 cm wide. It has white flowers with long peduncles which are 13 cm long, and there are 2 to 25 of them in each plant (Zhou *et al.* 2008). This plant which is the rarest species of *Lilium* grows in few areas of Iran such as Ardebil, Mazandaran, and Gilan provinces. In addition to Iran, it only grows in Lankaran in the Azerbaijan Republic (Rechinger 1989, Zhou *et al.* 2008). Although few studies have been done on *L. ledebourii* Bioss (Azadi 2003, Padasht 2005, Azadi and Khosh-Khui 2007), bioactive compounds of this plant are unknown. So, the aim of this study was to evaluate the phytochemical compounds of this native plant.

Materials and Methods

Lilium ledebourii Bioss bulbs were collected from Damash Mountains and were sent to the phytochemical laboratory of Iranian Biological Resource Center. For the preparation of Hydroalcoholic extracts 2 g of plant samples were dried by oven dry and freeze dry methods (Bernard *et al.* 2014) and shaken for 24 hrs at 30°C with 150 rpm in hydroalcoholic solution, and then filtrated. Finally, extracts were prepared for non-enzyme tests.

During Enzyme extraction in order to obtain cellular extract to study superoxide dismutase (SOD), peroxidase and catalase enzymes, about 1 g of fresh plant bulb were put into porcelain mortar and after adding liquid nitrogen, it was ground completely. Ground sample 0.5 gr was weighted and mixed with the buffer. Based on studied enzyme types, the enzyme extraction buffer was prepared. To produce extract one ml of intended buffer was added in the micro tube and was centrifuged in 4°C with 125000 rpm for 20 min.

To extract superoxide dismutase SOD at 4°C, 50 mM of potassium phosphate buffer with 7 pH and 0.5 mM EDTA was used. SOD activity was measured using spectrophotometry. One ml of reaction solution contains 0.1 mM EDTA, 50 mM phosphate buffer, 13 mM methionine, 75 µM NBT, and 0.21 ml molar riboflavin and enzyme supernatant. Covets containing reaction solution while were softly shaken for 15 min were put in front of inflorescence light (2 light bulbs with 250-watt inflorescence) and then sample absorption was read at 560 nm wavelength. In order to study enzyme activity, along with sample blank covets, control covets were used which contained blank cell, and similar control of sample cell. It should be mentioned that control cell along with samples was putted at the light and its wavelength was read as 560 nm, and blank cell putted in darkness. The activity of each SOD includes an amount of enzyme which is necessary for 50% reduction inhibition of phytochemical NBT under measurement condition (Dhindsa *et al.* 1980).

Extraction buffer for peroxidase (*POD*) enzyme was 50 mM potassium phosphate buffer with 7 pH including 0.5 EDTA and was held at 4°C environments. *POD* activity was studied for one min in a reactive solution (1 ml) including 475 µl guaiacol 475 µl of 100 mM hydrogen peroxidase, and 50 µl of extracted enzymatic supernatant and was read against blank cell including all materials except enzyme sample. *POD* activity was measurable through following absorption changes in 470 nm wavelength, which is the result of guaiacol oxidation (Polle *et al.* 1994)

To extract catalase (*CAT*) at 4°C from 50 mM potassium phosphate, the buffer including 1 mM EDTA and 2% PVPP was used. Catalase enzyme activity after decreasing H₂O₂ absorption determined with 39.4 mM/cm extinction coefficient for 2 min at 240 nm. It should be noted that some minor changes were applied in the mentioned method. One mM of reactive solution for determining *CAT* activity includes 10 mM hydrogen peroxidase, 25 mM phosphate buffer, 0.1 mM EDTA with 7 pH. Every level of catalase enzyme is definable in form of the content of enzyme, which decomposes one mM H₂O₂ in one min in testing condition (Rigo and Rotilio 1977).

The basis of reduction power (FRAP) method is on converting Fe^{3+} -TPTZ to Fe^{2+} -TPTZ at the presence of an antioxidant. FRAP solution contains 0.3 mM buffer acetate (3.6 PH), TPTZ solution in 40 mM HCL and 20 mM solution of iron chloride (III) with 1:1:10 sequence. FRAP should be prepared freshly. In this method 100 µl of diluted extract with 1.4 ml of FRAP was mixed and after 15 min, the absorption in 593 nm wavelength was read (Benzie and Strain 1999).

Total phenolic content in all samples was measured in the presence of folin-ciocalteu and then a standard graph with different concentrations of galic acid (0.01-0.05 mg/l) was drawn (Fig. 1). Some different concentrations which were different for various species were applied in order to obtain a linear range of each sample. For this purpose 500 µl of diluted extract were mixed with 500 µl of folin-ciocalteu, and after 3 min, 500 µl of 0.01 sodium carbonate were added. Finally, samples were put in darkness for two hrs at room temperature, and then the absorbance was read at 765 nm wavelengths. Results were presented in the form of mg GAE/g DW (Fig. 1). One important point in measuring the total phenolic content is the production of air bubbles in the presence of folin which interferes in absorbance, so before reading absorbance levels, air bubbles should be removed by using vortex (Spanos and Wrolstad 1990).




Fig. 1. Calibration curve for gallic acid standards to measure total phenol.

Total flavonoid content was measured using aluminum chloride. The solution contains 300 μ l of diluted solution, 1.6 ml water, and 100 μ l of 0.05 sodium nitrate. After 6 min, 150 μ l of 10% aluminum chloride was added. Next, after 5 min, 0.6 ml of 1 M sodium hydroxide and 250 μ l water were added and mixed completely by vortex. Then, the absorbance at 510 nm wavelength was read immediately. Standard graph with different routine concentrations (10-100 mg/l) was drawn and obtained results were presented in form of mg RE/g DW (Hosu *et al.* 2014).

In Inhibitory activity of DPPH free radicals method used for cleansing activity of DPPH radical was studied. The method is based on decreasing DPPH solution at the presence of hydrogen reducing antioxidants. The 300 μ l of 1 mM DPPH were mixed into 100 μ l of diluted sample and reached to their final volume of 2 ml by the use of methanol. After putting for 30 min in darkness, absorption at 517 nm wavelength was read (Sanchez-Moreno *et al.* 1998)

Methods used for the quality analysis of secondary compounds namely, tannin, saponin, cardiac glycosides, terpenoid, alkaloids and steroids are as follows:

Tannin: 0.5 g of powdered plant was poured into 20 ml of diluted water and warmed indirectly. Then the solution was filtrated and finally, some drops of ferric chloride III (0.1 %) were added to the extract. Development of brownish green and dark blue colors showed the presence of catechin tannin and pyrogallol tannin (Krishnaiah *et al.* 2009).

Saponin: Two grams of plant sample were boiled in 30 ml of diluted water in Bon Mary method (Rouhi-Boroujeni *et al.* 2016) and then it was filtrated. Filtrated samples (10 ml) were added to 5 ml of diluted water in a test tube and then the tube was shaken rapidly for about 1 min until stable foam produced. Then the produced foam was combined with three drops of olive oil. Emulsion (oil in water) and stable foam are representatives of saponin presence in plant extract (Krishniah *et al.* 2009).

Cardiac glycosides: One ml of dense sulfuric acid was poured into a test tube and set aside. Then 5 ml of diluted watery extract and 2 ml of acetic acid and one drop of ferric chloride were mixed together in another test tube. Then the content of the second tube was added to the first in such a way that dense sulfuric acid remains at the bottom. If the plant sample includes cardiac glycosides, then brown loops will appear (Krishniah *et al.* 2009).

Terpenoid: Five ml of diluted watery extract was combined with 2 ml of chloroform in a test tube and then they were added to 3 ml of dense sulfuric acid to be mixed. Appearance of reddish brown layer shows the presence of terpenoid compounds in plant extract (Krishniah *et al.* 2009).

Alkaloids: Two g of the powdered plant was extracted with 8 ml of chloroform and 10 ml of ammonia chloroform solution was added. Then the solution was diluted and was transferred to a test tube. Sulfuric acid amounting 0.5 ml (2 normal) was added to test tube and was shaken well. Finally, 3 drops of Mayer reagent were added. Milk like sediment shows the alkaloids in plant sample (Parekh and Chanda 2007)

Steroids: 200 mg of plant samples were extracted with 10 mg of chloroform. The solution was filtrated and 2 ml of dense sulfuric acid sample was slowly added to 2 ml of acetum anhydrate and 2 ml of dense sulfuric acid. Blue-green loop shows steroids in the extract (Parekh and Chanda 2007).

Results and Discussion

Analysis of Phytochemical metabolites in *L. ledebourii* Bloss showed the activity of superoxide dismutases, peroxidase, antioxidants, phenolic compounds and flavonoid but no catalase (Table 1).

Table 1. Results of phytochemical metabolites in *L. ledebourii* Bloss.

Plant	Superoxide dismutases	Peroxidase	Catalase	Antioxidants	Phenolic compounds	Flavonoid
<i>L. ledebourii</i>	*	*		*	*	*

*= Plant secondary compounds

The activities of superoxide dismutase and peroxidase enzymes were 13.82 ± 0.74 and 0.033 ± 0.002 $\mu\text{M}/\text{min}$, respectively (Table 2). Which shows that the activity of these enzymes in the bulb of *L. ledebourii* Bloss was low.

Table 2. Analysis of enzyme tests in *L. ledebourii* Bloss.

Explant	SOD ($\mu\text{M}/\text{min}$)	POD ($\mu\text{M}/\text{min}$)	CAT ($\mu\text{M}/\text{min}$)
Bulb	13.82 ± 0.74	0.033 ± 0.002	-

SOD = Superoxide dismutases, POD = Peroxidase, CAT = Catalase.

Superoxide dismutase (SOD) plays a protective role in plants. SOD is antioxidative which catalases more active superoxide anions and converts them to oxygen and less active types of hydrogen peroxidase. SOD causes stability of cell membrane of plants during drought (Jose *et al.* 1999). Plant peroxidases, which are widely spread enzymes in organic plants, play important role in antioxidative defensive system of the cell and detoxificate reactive oxygen species. PODs protect the cell against toxic amounts of H_2O_2 (Parida and Das 2005). No activity was observed for catalase in this plants (Table 2). Catalase enzyme protects cells against hydrogen peroxidase. and is necessary for some cells under the natural conditions. It plays an important role in resistance to oxidative stress in adoptive reactions of cells (Fridovich 1995). One important method for losing H_2O_2 is peroxidase. This enzyme which is found in all over the cell has more combining affinity to H_2O_2 than catalase (Jimnez *et al.* 1997). Plants can identify environmental

stresses and react to them. Like to other organisms, they activate radicals in reaction to biotic and abiotic stresses. In plants, producing reactive oxygen species (ROS) is for their adaptability and tolerance to different biotic and abiotic stresses. In this point of view, the activity of enzymes such as catalase (CAT), peroxidase (POD), and superoxide dismutases (SOD) thereby neutralize the activity of produced ROS in cells. Also, producing ROS in plant cells causes stimulations and increased activity of referred enzymes (Dat *et al.* 2000).

Flavonoid antioxidant activity in *L. ledebourii* Bioss based on freeze and oven dry methods was 0.78 ± 0.07 and 0.42 ± 0.02 mg RE/g DW, respectively (Table 3). The value of 0.78 ± 0.07 mg RE/g is less than Flavonoid compounds ($41.34 \text{ mgRE.g}^{-1}\text{DW}$) reported by Jin *et al.* 2012. The best method for drying non-enzymatic tests is freeze dry method. Flavonoids are a big group of a phenol compound with low molecular weight whose basic skeleton is formed by three benzene chains and a hydroxyl group. Mint (*Mentha*) extract has many phenol and flavonoid compounds and shows a good antioxidative activity (Sweetie *et al.* 2007).

FRAP method which studies the anti oxidative activity of extract based on iron rehabilitation ability, in fact, shows an estimation of antioxidative characteristics of soluble metabolites in water (Deepa *et al.* 2007). Enzymes are a group of proteins, which plays an activating role in the main metabolic pathway of plants. FARP antioxidant activity in *L. ledebourii* Bioss explants, based on freeze dry and oven dry methods was 25.88 ± 1.48 and 7.79 ± 0.48 $\mu\text{M Fe}^{2+}/\text{g DW}$, respectively (Table 3). FRAP method which studies the anti oxidative activity of extract based on iron rehabilitation ability, in fact, shows an estimation of antioxidative characteristics of soluble metabolites in water (Deepa *et al.* 2007). Enzymes are a group of proteins, which play activating role in the main metabolic pathway of plants.

The antioxidative defensive system is active in all growth steps of plants. The acts of these antioxidants differ, and their change depends on some factors such as maturity, weather, used parts of plants, harvest conditions (Mejia *et al.* 1988). During plant growth, phytochemical changes affect their antioxidative activity and influence the quality of different fruits and vegetables in particular periods of time (Conforti *et al.* 2007). Strong ant oxidative activity of extracts is related to the high levels of existing phenols and flavonoids in extracts (Shukla *et al.* 2009, Sun *et al.* 2011).

DPPH is a free radical with nitrogen as its central atom, which changes its color from purple to yellow when it is rehabilitee and is converted to DPPH-H stable molecule. DPPH radical own 515 to 528 nm absorption level, but as rehabilitee by an antioxidant, its absorption lessens. The antioxidative activity of samples appears in the disappearance of purple and final emerging yellow. This reaction is performed rapidly (Khalili and Ebrahimzadeh 2015). Trapping activity of DPPH free radicals based on freeze dry method was 3.24 ± 0.52 mg/ml while trapping activity of DPPH free radical in *L. concolor* was 4.531 TE $\mu\text{M/g}$ (Jin *et al.* 2012). Thus, trapping activity of DPPH free radical in *L. ledebourii* Bioss is lower than *L. concolor* and this is due to the low amount of phenolic compounds in *L. ledebourii* Bioss than *L. concolor*. DPPH is a stable free radical with nitrogen as the central atom which changes its color through reduction and converting 2, 2 diphenyl-1-picrylhydrazyl color from purple to yellow. DPPH in *L. ledebourii* Bioss explants using freeze and oven dry methods was 3.24 ± 0.52 and 11.27 ± 0.97 mg/ml, respectively (Table 3). Inhibitory activity of DPPH free radicals in plant extracts depends on the density and as density increases, inhibitory effect exceeds too (Shukla *et al.* 2009). Compositions in these extracts are able to give electrons to active free radicals.

Total phenol content that measured using Folin-Ciocalteu method (Sponos and Wrolstad 1990) and the linear equation of the standard curve was 1 mg of extract. The result of total

phenolic content was calculated from the regression equation of the standard plot ($Y = 18.82x - 0.0844$, $R^2 = 0.9983$) (Fig. 1).

The total content of phenol for *L. ledebourii* Bloss extract obtained by the use of Freeze Dry and Oven Dry methods was 5.198 ± 0.17 and 1.814 ± 0.05 mg GAE/g DW of extract, respectively (Table 3). Increase in the density of phenolic compounds directly increases the capability of different extracts in trapping free radicals (Yong *et al.* 2009). The total content of phenolic compounds is not an exact and stable criterion to prove the high antioxidative power of a sample, but its nature, diversity, and the amount of phenol and flavonoid are main indexes to determine high anti oxidative power. *L. ledebourii* Bloss bulb, a native flower in Iran, has considerable amounts of phenol compounds in freeze dry method (5.198 mg/g of extract). This is much lower than that in *L. concolor* (389.76 mg in extract) reported by Jin *et al.* (2012). This indicates that phenolic compounds in *L. ledebourii* Bloss bulbs have very little activity. In all plants, antioxidative activity has a direct relationship to phenol and flavonoid levels. Mint (*Mentha*) extract has many phenol and flavonoid compounds and shows a good antioxidative activity (Sweetie *et al.* 2007). Rosemary (*Rosmarinus officinalis* L.) extract has high anti oxidative activity and this activity is in a close relationship with plants phenol content (Elmasta *et al.* 2006).

Table 3. Results of the non-enzyme test in *L. ledebourii* Bloss bulb explants.

Drying method	Flavonoid mg RE/g DW	FRAP $\mu\text{M Fe}^2/\text{g DW}$	DPPH (IC ₅₀) mg/ml	Total Phenol mg GAE/g DW
Oven Dry	0.02 ± 0.42	0.48 ± 7.79	0.97 ± 11.27	0.05 ± 1.814
Freeze Dry	0.07 ± 0.78	1.47 ± 25.88	0.52 ± 3.24	0.17 ± 5.198

FRAP = ferric reducing/antioxidant power and DPPH = 2,2-diphenyl-1-picrylhydrazyl

Iron III rehabilitee method is applied as a criterion for electron endowing ability of phenolic compositions. It forms an important mechanism in oxidation process of phenol composition. Electron endowing capacity (rehabilitating power) is related to the anti oxidative activity of biological compounds (Arabshani-Delouee and Urooj 2007).

Existence of some secondary metabolites namely cardiac glycosides, tannin, saponin, steroids, alkaloids and terpenoid in the bulb of *L. ledebourii* Bloss. was observed. Presence of tannin activity was not observed (Table 4).

Table 4. Analysis of phytochemical compounds in *L. ledebourii* Bloss.

Plant	Cardiac glycosides	Tannin	Saponin	Steroids	Alkaloids	Terpenoid
<i>L. ledebourii</i>	*		*	*	*	*

*= Plant secondary compounds.

Considering the results, the importance of this research and the importance of this domestic plant these kinds of investigations are crucial and valuable. Although these tests are the introductory identification of chemical components, but open the path to further researches where new techniques could be applied.

Acknowledgments

The Authors are thankful to the staffs and managers of Iranian Biological Resource Center for providing necessary facilities to conduct this study.

References

- Akinmoladun AC, Ibukun EO, Afor E, Obuotor EM and Farombi EO 2007. Phytochemical constituent and antioxidant activity of extract from the leaves of *Ocimum gratissimum*. *Sci. Res. Essay* **2**: 163-166.
- Arabshahi-delouee S and Urooj A 2007. Antioxidant properties of various solvent extracts of mulberry (*Morus indica* L.) leaves. *Food Chem.* **102**: 1233-1240.
- Asghari J and Mazaheri-tehrani M 2010. Extraction of tannin from *Eucalyptus camaldulensis* Dehnh and trimyristin from *Myristica fragrans* Houtt by using microwave irradiation. *J. Medicinal Aromatic Plants* **26**(2): 185-195.
- Azadi P 2003. Effect of growth regulators, sucrose concentration and scale pieces on micropropagation of Chelcheragh lily (*Lilium ledebourii*) in spring season. *Proceedings of the 2nd Sci. and Applied Seminar on Ornament. Plants.* p. 43
- Azadi P and Khosh-Khui M 2007. Micropropagation of *Lilium ledebourii* (Baker) Bioss as affected by plant growth regulator, sucrose concentration, harvesting season and cold treatments. *Electronic J. Biotech.* **10**(4): 582-591.
- Benzie IFF and Strain JJ 1999. Ferric reducing/antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous of total antioxidant power and ascorbic acid concentration. *Methods in Enzymol.* **299**: 15-27.
- Bernard D, Kwabena AI, Osei OD, Achel Daniel G, Achoribo Elom S and Sandra A 2014. The Effect of Different Drying Methods on the Phytochemicals and Radical Scavenging Activity of Ceylon Cinnamon (*Cinnamomum zeylanicum*) Plant Parts. *European J. Medicinal Plants* **4**(11): 1324-1335.
- Bouayed J, Hoffmann L and Bohn T 2011. Total phenolics, flavonoids, anthocyanins and antioxidant activity following simulated gastro-intestinal digestion and dialysis of apple varieties: Bioaccessibility and potential uptake. *Food Chem.* **128**: 14-21.
- Conforti F, Statti GA and Menichini F 2007. Chemical and biological variability of hot pepper fruits (*Capsicum annuum* var. *acuminatum* L.) in relation to maturity stage. *Food Chem.* **102**: 1096-1104.
- Dat J, Vandenabeele S, Vranová E, Van montagu M, Inzé D and Van breusegem F 2000. Dual action of active oxygen species during plant stress responses. *Cell Mole Life Sci.* **57**: 779-795.
- Deepa N, Kaura CH, Georgea B, Singhb B and Kapoor HC 2007. Antioxidant constituents in some sweet pepper (*Capsicum annuum* L.) genotypes during maturity. *LWT Food Sci. Tech.* **40**: 121-129.
- Dhindsa RS, Dhindsa PP and Thorpe TA 1980. Leaf senescence correlated with increased levels of membrane permeability and lipid-peroxidation and decreased levels of superoxide dismutase and catalase. *J. Exp. Bot.* **32**: 93-101.
- Edeoga HO, Okwu DE and Mbaebie BO 2005. Phytochemical Constituents of some Nigerian Medicinal Plants. *African J. Biotech.* **4**(7): 685-688.
- Elmastas M, Gülçin IIL DAK, Küfrevio LU, Bao lu K and Aboul-enein HY 2006. Radical scavenging activity and antioxidant capacity of Bay leaf extracts. *J. Iran Chem.* **3**(3): 1258-266.
- Fridovich L 1995. Superoxide radical and superoxide dismutases. *Ann. Rev. Biochem.* **64**: 97-112.
- Gardner PT, White TAC, Mcphail DB and Duthie GC 2000. The relative contributions of vitamin C, carotenoids and phenolics to the antioxidant potential of fruit juices. *Food Chem.* **68**: 471-474.
- Hosu A, Cristea VM and Cimpoi 2014. Analysis of total phenolic, flavonoids, anthocyanins and tannins content in Romanian red wines: Prediction of antioxidant activities and classification of wines using artificial neural networks. *Food Chem.* **150**: 113-118.
- Jamshid M, Ahmadi HR, Rezazadeh SH, Fathi F and Mazanderani M 2010. Study on phenolic and antioxidant activity of some selected plant of Mazandaran province. *Medic Plant* **9**(34): 177-183.
- Jimenez A, Hernandez JA, Del rio LA and Sevilla F 1997. Evidence for the presence of the ascorbate-glutathione cycle in mitochondria and peroxisomes of pea leaves. *Plant Physiol.* **114**: 275-284.
- Jin L, Zhang Y, Yan L, Guo Y and Niu L 2012. Phenolic compounds and antioxidant activity of bulb extracts of six *Lilium* species native to China. *Mole J.* **17**: 9361-9378.

- Jose M, Eristinaperez M and Gomez 1999. Antioxidant enzymes and human disease chemical biochemistry. *Chemical Biochem.* **32**(8): 595-603.
- Khalili MA and Ebrahimzadeh MA 2015. A review on antioxidants and some of their common evaluation methods. *J. Mazandaran Univ. Med. Sci.* **24**(120): 188-208.
- Krishnaiah D, Devi T, Bono A and Sarbatly R 2009. Studies on phytochemical constituents of six Malaysian medicinal plants. *J. Medicinal Plants Res.* **3**(2):067-072.
- Kumar Gupta A and Neelam M 2006. Hepatoprotective activity of aqueous ethanolic extract of chamomile capitula in paracetamol intoxicated albino rats. *Amer. J. Pharmacol and Toxicol.* **1**(1): 17-20.
- Kumaran A and Karunakaran RJ 2006. Antioxidant and free radical scavenging activity of an aqueous extract of *Coleus aromaticus*. *Food Chem.* **97**: 109-114.
- Maber P and Hanneken A 2005. Flavonoids protect retinal ganglion cells from oxidative stress-induced death. *Investigative Ophthalmol Visual Sci.* **46**(12): 4796-4803.
- Macheix JJ, Fleurent A and Billot J 1990. Phenolic compounds in fruit processing. *In: Fruit Phenolic*. CRC Press, Boca Raton, FL. pp. 295-342.
- Mejia LA, Hudson E, Gonzalez, DE, Mejia E and Vasquez F 1988. Carotenoid content and vitamin A activity of some common cultivars of Mexican peppers (*Capsicum annuum*) as determined by HPLC. *J. Food Sci.* **53**: 1448-1451.
- Miliauskas G, Venskutonis PR and Vnbeek, TA 2004. Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chem.* **85**:231-237.
- Mohsen SM and Ammar ASM 2009. Total phenolic contents and antioxidant activity of corn tassel extracts. *Food Chem.* **112**: 595-598.
- Padasht dahkaei MN 2005. The investigation of different methods for culturing and propagation of Chelcheragh lily (*Lilium ledebourii*), native of Iran, and its introduction possibility as a new floricultural crop. Ph. D. Thesis, Islamic Azad University, Science and Research Unit, Tehran, Iran. pp. 120.
- Parekh J and Chanda SV 2007. *In vitro* Antimicrobial activity and phytochemical analysis of some Indian medicinal plants. *Turk. J. Biol.* **31**: 53-58.
- Parida, AK and Das AB 2005. Salt tolerance and salinity effects on plants: A review. *Ecotoxicol and Environmental Safety* **60**: 324-349.
- Pellegrini N, Simonetti P, Gardana C, Brenna O, Brighenti F and Pietta PG 2000. Polyphenol content and total antioxidant activity of vini novelli (young red wines). *J. Agri. Food Chem.* **48**: 732-735.
- Polle A, Otter T and Seifert F 1994. Apoplastic peroxidases and lignification in needles of Norway spruce (*Picea abies* L.). *Plant Physiol.* **106**: 53-60.
- Rechinger KH. 1989. *Flora Iranica*, Liliaceae. Akademische Druck-u, Verlagsantalt, Graz, Austria. pp. 58-59.
- Rigo A and Rotilio G 1977. Simultaneous determination of superoxide dismutase and catalase in biological materials by polarography. *Ann. Rev. Biochem.* **81**: 157-166.
- Rohi-Boroujeni H, Heidarian S, Rohi-Boroujeni H and Rafieian-Kopaei M. Phytochemical constituents of *Moringa peregrina* seeds, *Der Pharma Chemica*, **8**(5):80-86
- Sanchez-Moreno C, Larrauri JA and Saura-Calixto F 1998. A procedure to measure the antiradical efficiency of polyphenols. *J. Sci. Food Agric.* **76**: 270-276.
- Shukla SH, Mehta A, Bajpai VK and Shukla S 2009. *In vitro* antioxidant activity and total phenolic content of ethanolic leaf extract of *Stevia rebaudiana* Bert. *Food Chem. Toxicol.* **47**: 2338-43.
- Singh KBR, Malhotra S, Halila MH, Kinghts EJ and Verma MM 1994. Current status and future strategy in breeding chickpea for resistance to biotic and abiotic stresses. *Euphytica* **73**: 137-149.
- Sponos GA and Worlsted RE 1990. Influence of processing and storage on phenolic composition of Thompson seedless grape juice. *J. Agric Food Chem.* **38**: 1565-1571.
- Sun L, Zhang J, Lu X, Zhang L and Zhang Y. 2002. Evaluation to the antioxidant activity of total flavonoids extract from persimmon (*Diospyros kaki* L.) leaves. *Food Chem. Toxicol.* **49**: 2689-2696.
- Sweetie R, Ramesh CH and Arun SH 2007. Antioxidant potential of mint (*Menthaspicata* L.) in radiation processed lamb meat. *Food Chem.* **100**(2): 451-458.

- Taiz L and Zeiger E. 2002. Plant Physiology, 3rd edition. pp. 690.
- Velioglu YS, Mazza G, Gao L and Oomah BD 1998. Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. J. Agri. Food Chem. **46**: 4113-4117.
- Young kil H, Seong ES, Ghimire BK, Chung IM, Kwon SS, Goh EJ, Heo K, Kim MJ, Lim JD, Lee D and Yu CY. 2009. Antioxidant and antimicrobial activities of crude sorghum extract. Food Chem. **115**: 1234-1239.
- Zhou S, Ramanna, MS, Visser RGF and Van tuyl JM 2008. Analysis of the meiosis in the F1 hybrids of *Longiflorum* × Asiatic (LA) of lilies (*Lilium*) using genomic in situ hybridization. J. Genetics and Genom. **35**: 687-695.

(Manuscript received on 12 April, 2018; revised on 2 June, 2018)