

EVALUATION OF ANTIMICROBIAL AND CYTOTOXIC EFFECTS OF FOUR TURKISH SPECIES OF *ERYNGIUM* L.

BERNA ÖZBEK CELİK*, EMEL MATARACI KARA, CAGLA BOZKURT GUZEL,
GÜLAY ECEVIT GENÇ¹, ILKER GENÇ¹, SEZİN KURTOĞLU ANIL² AND GULAY MELİKOĞLU²

Department of Pharmaceutical Microbiology, Faculty of Pharmacy,
Istanbul University, Beyazıt, 34116, Istanbul, Turkey

Keywords: *Eryngium*, Antibacterial, Antifungal, Cytotoxic, Herbal therapy

Abstract

The antimicrobial and cytotoxic effects of various endemic and non-endemic *Eryngium* species from Turkey were investigated. Nine endemic and two non-endemic *Eryngium* species were analyzed using a microdilution assay for their antibacterial and antifungal activities. Cytotoxic activities of *E. pseudothoriifolium*, *E. thoriifolium*, *E. davisii*, and *E. falcatum*, on prostate carcinoma and endometrial cancer cells were analyzed by cytotoxic activity assay. It was observed that 16 out of 22 extracts (aerial or root part) showed antibacterial activities. Fifteen out of 22 extracts showed antifungal activity with the lowest MIC (156 mg/l) value against *Candida albicans* ATCC 10231 and *C. tropicalis* ATCC 750. Aerial parts of *E. pseudothoriifolium*, *E. thoriifolium*, *E. davisii*, and *E. falcatum* exhibited cytotoxic effects on endometrial cancer cells. This evaluation of tested endemic *Eryngium* species' antimicrobial and cytotoxic activities is the first of its kind.

Introduction

The genus *Eryngium* L., belonging to the subfamily Saniculoideae of Apiaceae, is the largest and arguably the most taxonomically complex genus of the family Apiaceae. The genus has 250 species (317 taxa) in the world, which make it the most species rich genus of the *Apiaceae* (Pimenov 1993). The *Eryngium* species contain acetylenes, flavonoids, coumarins, and triterpene saponins. *Eryngium* taxa have a long tradition as a popular culinary and medicinal plant in the world (Paul *et al.* 2011). Some *Eryngium* species are used as folk medicine or ornamental plant and vegetables. Similarly, in Turkish folk medicine various species of the plant are used for a wide range of ailments fresh (Gümüş 1994, Sezik *et al.* 1997, Ertug 2000, Sezik 2001, Özgökçe and Özçelik 2004, Ozturk 2005, Tuzlacı 2005, Ecevit 2006). In particular, roots are used against various inflammatory disorders, sinusitis, colds, goiters, respiratory diseases, digestive diseases, and snake or scorpion bites as well as a diuretic and antitussive (Kupeli 2006, Mekhora 2012). Their leaves are used for infertility and as herbs for wound healing and treating kidney stones as well as food while fresh.

The authors aimed to investigate the antimicrobial effects of both aerial and root parts of nine endemic *Eryngium* species (*E. isauricum* Contandr. & Quézel, *E. kotschyi* Boiss., *E. trisectum* Wörz & H. Duman, *E. bithynicum* Boiss., *E. davisii* Kit Tan & Yıldız, *E. babadaghensis* G.Ecevit-Genç, E.Akalin & A.Wörz, *E. polycephalum* Hausskn. ex H.Wolff, *E. thoriifolium* Boiss., *E. pseudothoriifolium* Contandr. & Quézel) and two non-endemic *Eryngium* species (*E. glomeratum* Lam. and *E. falcatum* F. Delaroché) from Turkey were investigated. One of the non-endemic strains, *E. glomeratum*, is an east Mediterranean species, with a main distribution range in south

*Author for correspondence: <berna.ozbek@istanbul.edu.tr>, <bernaozbek@hotmail.com>. ¹Department of Pharmaceutical Botany, Faculty of Pharmacy, Istanbul University, Beyazıt, 34116, Istanbul, Turkey. ²Department of Pharmacognosy, Faculty of Pharmacy, Istanbul University, Beyazıt, 34116, Istanbul, Turkey.

Turkey, Lebanon, Palestine, and Israel. Other strain, *Eryngium falcatum*, is a widespread species on coastal sand dunes and usually common along the Balkan coasts. Additionally, cytotoxic activities of *E. pseudothoriifolium*, *E. thoriifolium*, *E. davisii*, and *E. falcatum*, which have more antimicrobial activities, on prostate carcinoma (PC-3) and endometrial cancer (ECC-1) cell lines at 12.5, 25, 100, 250, 500 g/ml were analyzed by cytotoxic activity assay.

Materials and Methods

Plant materials were collected from different locations in Turkey. The species were collected and identified by G. Ecevit Genc and I. Genc, which were deposited in the Herbarium of the Faculty of Pharmacy, Istanbul University (ISTE), Turkey. The plant materials used in this study are listed in Table 1. The root and the aerial parts of the plants were separated. The dried plant materials were percolated with methanol (95%) at room temperature. The methanolic extracts (MEs) were evaporated to dryness under pressure and controlled temperature (400 - 500°C) in a rotary evaporator. All the extracts were kept at -20°C and then lyophilized. Crude methanolic extracts were obtained.

Table 1. The plant materials used in this study.

Species	Collection data	Collector number
<i>E. falcatum</i>	Muğla, Fethiye Gidrak oition, 29. viii 2014, G. Ecevit Genc & I.Genc	IG 2282
<i>E. thoriifolium</i>	Muğla, Fethiye-Korkuteli road, Bayır village 29 viii 2014, G. Ecevit Genc & I.Genc	IG 2283
<i>E. polycephalum</i>	Konya: Bozkir-Hadim road, Korualan, 28. viii 2014, G. Ecevit Genc & I.Genc	IG 2272
<i>E. babadaghensis</i>	Muğla, Fethiye, Babadag, Esek Bayiltan position, 29. viii 2014, G. Ecevit Genc & I.Genc	IG 2281
<i>E. davisii</i>	Konya: Bozkir-Hadim road, Korualan, Akdam plateau, 28. viii 2014, G. Ecevit Genc & I.Genc	IG 2273
<i>E. bithynicum</i>	Konya-Seydişehir road, 27. viii 2014, G. Ecevit Genc & I.Genc	IG 2270
<i>E. kotschyi</i>	Konya: Bozkir-Hadim road, Korualan, Akdam plateau, rocky slopes, 29. viii 2014, G. Ecevit Genc & I.Genc	IG 2274
<i>E. trisectum</i>	Konya: Derebucak, Çamlık town, Kızıldağ., 27. viii 2014, G. Ecevit Genc & I.Genc	IG 2271
<i>E. isauricum</i>	Konya, Ermenek, Tepebaşı-Güzelyurt, 29 viii 2014, G. Ecevit Genc & I.Genc	IG 2277
<i>E. pseudothoriifolium</i>	Muğla, Fethiye-Korkuteli road, Avlan village road, 29 viii 2014, G. Ecevit Genc & I.Genc	IG 2285
<i>E. glomeratum</i>	Muğla, Fethiye, Babadağ road, 29. viii 2014, G. Ecevit Genc & I.Genc	IG 2279

Antimicrobial activity of the crude methanolic extracts against *Staphylococcus aureus* ATCC 29213, *Streptococcus epidermidis* ATCC 12228, *Escherichia coli* ATCC 25922, *Klebsiella pneumoniae* ATCC 4352, *Pseudomonas aeruginosa* ATCC 27853, *Proteus mirabilis* ATCC 14153, *Candida albicans* ATCC 10231 and *Candida tropicalis* ATCC 750 were determined by

the microbroth dilution technique using the Clinical and Laboratory Standards Institute (CLSI) recommendations (CLSI 2000a, CLSI 2000b). For the bacteria Mueller-Hinton broth, for yeast strains RPMI-1640 medium were used as the test medium. From 10000 mg/l crude extract in Mueller-Hinton broth or RPMI-1640 medium, serial two-folds dilution (5000 to 4.9 mg/l) were prepared. The inoculum was prepared using a 4 - 6 hrs broth culture of each bacterium and 24 hrs culture of yeast strains adjusted to a turbidity equivalent to a 0.5 McFarland standard, then diluted in broth media to give a final concentration of 5×10^5 cfu/ml for bacteria and 0.5×10^3 to 2.5×10^3 cfu/ml for yeast in the test tray. Well series with cefuroxime or ceftazidime antibiotics and an antifungal agent (clotrimazole) were also included to serve as positive controls. Antibiotic-free negative controls were also used. The trays were covered and placed into plastic bags to prevent evaporation. The trays containing Mueller-Hinton broth were incubated at 35°C for 18 - 20 hrs and the trays containing RPMI-1640 medium were incubated at 35°C for 46 - 50 hrs. The MIC was defined as the lowest concentration of compound giving complete inhibition of visible growth.

The cell culture medium for PC-3 and ECC-1 cell lines were Dulbecco's Modified Eagle's Medium (DMEM) (Winsent, St-Bruno, QC, Canada) supplemented with 10% heated fetal bovine serum (FBS) (Biochrome, Berlin, Germany) and 1% penicillin-streptomycin solution (Winsent, St-Bruno, QC, Canada). Prepared media were filtered by 0.22 μ m sterile filter (Minisort, Sartorius, Germany). The PC-3 and ECC-1 cells were cultured at 37°C under a humidified atmosphere containing 5% CO₂. According to protocol (Park 1994, McDaid 1999, Freshney 2000), cells were first cultured in 96-well plates at a density of 2×10^4 cells/well for 24 hrs. After incubation, the medium was changed, and the cells were treated with different concentrations of the plant extracts (12.5, 25, 100, 250, 500 g/l dilutions of the lyophilized extracts in dimethylsulphoxide [DMSO]). Following incubation for 48 hrs, cytotoxicity assays were carried out.

Cytotoxicity in the cells was monitored using MTT assay. Cells in their log growth phase were resuspended with trypsin 0.25%-EDTA 10 mM. For the microtitration assay, 2.5×10^4 cells in 0.2 ml of DMEM were seeded in each of the 96-well microtitre plates and incubated in a humidified atmosphere at 37°C for 24 hrs. At that time, when the cells were in the exponential phase of growth, different dilutions of the methanolic plant extracts, prepared in medium, were added (4 wells were used for each plant extract concentration to give quadruplicate determinations within each experiment). The 96-well microtitre plates were further incubated for 48 hour. Cellular growth control was performed using medium alone or with DMSO instead of plant extract. The DMSO concentration for the tested dilutions was not higher than 1%, as in the solvent-control wells. Fresh medium then replaced the medium from all the wells, and the plate was further incubated for 24 hrs. Cell proliferation was evaluated with MTS/PMS (Cell Titer 96 Aqueous Non-Radioactive Cell Proliferation Assay, Cat.G5421-Promega). The medium of all the plates was removed and 100 μ l of fresh GM with 20 μ l of MTS/PMS (20 : 1) was added to each well. The plates, wrapped in aluminum foil, were incubated for 3.5 hrs in a humidified atmosphere at 37°C. Absorbance at 490 nm was recorded in a microplate reader (Bio-Rad, USA). The wells with medium and MTS/PMS without cells were utilized as blanks for the plate reader. The IC₅₀ concentration was determined to be the plant extract concentration required to reduce the absorbance (490 nm) to half the control value.

Results and Discussion

According to previous studies, *in vitro* bioactivities such as cytotoxicity, against anti-inflammatory, antimicrobial, antioxidant, and antihyperglycemic effects were demonstrated with *Eryngium* extracts. Ndip *et al.* (2007) showed that the methanolic extract from *E. foetidum* leaves showed moderate antibacterial activity against *Helicobacter pylori* (Ndip 2007). Another investigation on the leaf hydromethanolic extract of *E. maritimum* showed that using the

microdilution method against food-borne pathogens and clinical isolates, exhibited antimicrobial activity (Meot-Duros 2008). Similarly, the antimicrobial activity of three other species belonging to genus *Eryngium* (*E. creticum*, *E. campestre* and *E. thoriifolium*) were studied with the disc diffusion method against nine clinical strains of methicillin-resistant *S. aureus* (MRSA) (Celik 2011) and the essential oil obtained from *E. thoriifolium*, which caused an inhibition zone ranging from 13 to 19 mm (similar to that exhibited by vancomycin), was demonstrated to be the most active species. According to the present results, it was observed that 16 of the 22 extracts (aerial or root parts) showed antibacterial activity with the lowest MIC (78.1 mg/l) value against tested Gram-positive bacteria (*S. aureus* and *S. epidermidis*). Thirteen of the 22 extracts showed antifungal activity with the lowest MIC (625 mg/l) value against *C. albicans* and *C. tropicalis*. The MICs ranged from 0.22 to 5000 mg/l. The microbial sensitivity to the different extracts represented by the mean MIC values ranged from 78 to 1250 mg/l (Table 2). *E. falcatum* was the most sensitive species (lowest MIC = 625 mg/l), which has 8 activities against Gram-positive or Gram-negative bacteria. It was followed by *E. thoriifolium* and *E. pseudothoriifolium* (both aerial and root parts), each of which had 6 antibacterial activities against Gram-positive or Gram-negative bacteria (lowest MIC = 78 mg/l). Overall, in the present study the Gram-positive bacteria were more sensitive to the extracts than the Gram-negative bacteria. The cell walls of Gram-positive bacteria compared with Gram-negative bacteria, are more sensitive to antimicrobial chemical compounds and even many herbal drugs. This may be because of inherent tolerance of Gram-negatives and the nature and composition of herbs. Having a complex cell wall, which acts as an effective permeability barrier to restrict the penetration of compounds and lipopolysaccharides layer and periplasmic space of Gram-negative bacteria are the reasons of relative resistance of Gram-negative bacteria (Sharifa 2008). Furthermore a set of multidrug resistance pumps in their cell wall extrudes toxins across the outer membrane (Tegos 2002). Similarly, Thiem *et al.* (2010) studied the antimicrobial activity of ethanolic extracts from leaves and roots of 3 *Eryngium* genera (*E. planum*, *E. campestre*, and *E. maritimum*) native to Poland, and these plants were tested by the method of series dilutions against different Gram-positive bacteria (2 strains) and fungi (5 species). The results have shown that the ethanolic extracts inhibit the growth of *S. aureus* and all tested fungi (Thiem 2010). The present results confirm that *E. isauricum* (root part), *E. davisii* (both aerial and root parts), *E. falcatum* (both aerial and root parts), and *E. thoriifolium* (both aerial and root parts) exhibit a moderate antibacterial activity on Gram-positive strains of *S. aureus* and *S. epidermidis*. A weak antibacterial effect on *P. aeruginosa* was highlighted for *E. davisii* (root part), *E. falcatum* (root part), *E. thoriifolium* (aerial part) and *E. pseudothoriifolium* (aerial part). Furthermore, the methanolic extracts from *E. pseudothoriifolium* (aerial parts and roots), *E. triseicum* (aerial part), and *E. babadaghensis* (root part) showed activity towards *P. mirabilis* with MIC values in the range of 1250 µg/ml. The extract from aerial parts and roots of *E. falcatum* and the aerial part of *E. thoriifolium* showed activity toward *K. pneumoniae* with MIC values in the range of 1250 µg/ml. However, no activity was found with tested plants against *E. coli*.

Candida species with *C. albicans* -as the most common- are causal agents of opportunistic human infections. In addition, hospital-acquired infections by *C. albicans* have become a cause of major health concerns (Ferreira 2013). It was previously reported that many *Eryngium* species, like *E. maritimum* exerted antifungal activity against *C. albicans* and other strains (Abou-Jawdah 2002). Also, in an in vitro antimycotic activity screening against eight phytopathogenic fungi, *E. creticum* showed more than 95% inhibition of spore germination in at least two fungi (Yusuf 2002). According to the present results, aerial and root parts of *E. kotschyi*, *E. glomeratum*, *E. davisii*, *E. thoriifolium*, and *E. pseudothoriifolium* having minimum MIC of 625 mg/l or 1250 mg/l were considered active against *C. albicans* and are summarized in Table 2. Also, *E.*

babadaghensis (root part) and *E. polycephalum* (aerial part) had antifungal activity against *C. albicans*. On the other hand, the antifungal activity of the aerial and root extract of *E. kotschy* and *E. pseudothorifolium* were 156.25 µg/ml against *C. tropicalis*. Additionally, the root part of *E. isauricum* and aerial part of *E. bithynicum* had moderate antimicrobial activity for *C. tropicalis* at 312.5 mg/ml. More plants showed considerable antifungal activity against *C. albicans* than *C. tropicalis*.

Table 2. Antibacterial and antifungal effects of various endemic *Eryngium* species.

Extracts	Sa	Se	Ec	Ef	Kp	Pa	Pm	Ca	Ct
	ATCC 29213	ATCC 12228	ATCC 25922	ATCC 29212	ATCC 4352	ATCC 27853	ATCC 14153	ATCC 10231	ATCC 750
<i>E. isauricum</i> aerial	-	-	-	-	-	-	-	-	156.25
<i>E. isauricum</i> root	1250	1250	-	-	-	-	-	1250	-
<i>E. kotschy</i> aerial	-	-	-	-	-	-	-	1250	156.25
<i>E. kotschy</i> root	-	1250	-	-	-	-	-	1250	156.25
<i>E. glomeratum</i> aerial	-	-	-	-	-	-	-	1250	-
<i>E. glomeratum</i> root	-	-	-	-	-	-	-	1250	-
<i>E. trisectum</i> aerial	1250	-	-	-	-	-	1250	-	-
<i>E. trisectum</i> root	-	-	-	-	-	-	-	-	-
<i>E. bithynicum</i> aerial	-	-	-	-	-	-	-	-	-
<i>E. bithynicum</i> root	-	-	-	-	-	-	-	-	312.5
<i>E. davisii</i> aerial	625	625	-	-	-	-	-	1250	312.5
<i>E. davisii</i> root	625	1250	-	-	-	625	-	1250	-
<i>E. falcatum</i> aerial	1250	1250	-	-	1250	-	-	-	-
<i>E. falcatum</i> root	1250	625	-	625	1250	625	-	-	-
<i>E. babadaghensis</i> aerial	-	-	-	-	-	-	-	-	-
<i>E. babadaghensis</i> root	-	-	-	-	-	-	1250	1250	-
<i>E. polycephalum</i> aerial	-	-	-	-	-	-	-	1250	-
<i>E. polycephalum</i> root	-	-	-	-	-	-	-	-	-
<i>E. thorifolium</i> aerial	78.12	312.5	-	-	1250	625	-	625	-
<i>E. thorifolium</i> root	312.5	625	-	-	-	-	-	1250	-
<i>E. pseudothorifolium</i> aerial	-	625	-	-	-	625	1250	1250	156.25
<i>E. pseudothorifolium</i> root	-	312	-	-	-	625	1250	1250	156.25
Positive controls MIC (mg/l)	CXM 1.2	CXM 9.8	CXM 4.9	CXM 4.9	CXM 4.9	CAZ 2.4	CXM 2.4	CLT 4.9	CLT 4.9

Staphylococcus aureus American Type Culture Collection (ATCC) 29213 (Sa), *Streptococcus epidermidis* ATCC 12228 (Se), *Escherichia coli* ATCC 25922 (Ec), *Klebsiella pneumoniae* ATCC 4352 (Kp), *Pseudomonas aeruginosa* ATCC 27853 (Pa), *Proteus mirabilis* ATCC 14153 (Pm), *Candida albicans* ATCC 10231 (Ca), and *Candida tropicalis* ATCC 750 (Ct). CXM: Cefuroxime; CAZ: Ceftazidime; CLT: Clotrimazole.

There are few studies on the cytotoxic effects of *Eryngium* species but none for *E. pseudothorifolium*, *E. thorifolium*, *E. davisii* and *E. falcatum*. According to the previous studies on *Eryngium* species, ethanol extracts of *E. planum* displayed cytotoxic activities to some leukemia cell lines (Bogucka-Kocka 2008). In another study by Zhang et al. (2008), 3 eryngiosides (eryngioside J, eryngioside L, and saniculasapoinin III) isolated from *Eryngium yuccifolium* were found to be mostly effective on human cancer cell line A-549 and normal cell line MRC-5, and they markedly inhibited the growth of pancreas cancer cell line PANC-1. Also, compounds isolated from the roots of *Eryngium campestre* showed weak cytotoxic activity against human tumor cell lines HCT 116 and HT-2923 (Kartal et al. 2005). Another study by Yurdakok

and Baydan (2013) revealed that, aerial and root parts of *E. maritimum* and the endemic *E. kotschyi* induced cytotoxicity on Hep2, U138-MG, HepG2 and Vero cell lines in a dose-dependent manner with both LDH and MTT assays (Yurdakok and Baydan 2013). Similarly, in the present study, the anti-IC50 values for aerial parts of *E. pseudothoriifolium*, *E. thoriifolium*, *E. davisii*, and *E. falcatum* on ECC-1 cells (23.14, 10.41, 13.51, and 22.72 mg/ml, respectively) by MTT assay were found to be lower than the US National Cancer Institute recommendations (IC50 < 30 mg/l) to define the activity against cancer cells (Table 3). Following this fact, *E. pseudothoriifolium*, *E. thoriifolium*, *E. davisii*, and *E. falcatum* aerial extracts examined in this study with observed IC50 values lower than 30 mg/l were considered to have significant activity on ECC-1 cells. However, it was also found that these extracts (aerial parts of *E. pseudothoriifolium*, *E. thoriifolium*, *E. davisii*, and *E. falcatum*) had no cytotoxic activity on PC-2 at tested concentrations.

Table 3. IC50 values for aerial parts of the tested *Eryngium* spp. (mg/ml)

	<i>E. pseudothoriifolium</i>	<i>E. thoriifolium</i>	<i>E. davisii</i>	<i>E. falcatum</i>
*ECC-1 cells	23.14	10.41	13.51	22.72
*PC-2 cell	> 30	> 30	> 30	> 30

*Endometrial cancer (ECC-1); prostate carcinoma (PC-3).

The present study showed that all endemic plants (*E. isauricum* Contandr. & Quézel, *E. kotschyi* Boiss., *E. trisectum* Wörz & H. Duman, *E. bithynicum* Boiss., *E. davisii* Kit Tan & Yıldız, *E. babadaghensis* G. Ecevit-Genç, E. Akalın, & A. Wörz, *E. polycephalum* Hausskn. ex H. Wolff, *E. thoriifolium* Boiss., *E. pseudothoriifolium* Contandr. & Quézel) could be a potential source for inhibitory substances for human pathogenic fungi and bacteria. Furthermore, aerial parts of *E. pseudothoriifolium*, *E. thoriifolium*, *E. davisii* and *E. falcatum* induced cytotoxicity on ECC-1 cells with MTT assays.

Acknowledgments

This work was supported by Scientific Research Projects Coordination Unit of Istanbul University. Project number: 43682.

References

- Abou-Jawdah Y, Sobh H and Salameh A 2002. Antimycotic activities of selected plant flora, growing wild in Lebanon, against phytopathogenic fungi. *J. Agric. Food Chem.* **50**(11): 3208-3213.
- Bogucka-Kocka A, Smolarz HD and Kocki J 2008. Ethanol extracts of *E. planum* displayed sitotoxic activities to some leukemia cell lines. *Fitoterapia* **79**(7-8): 487-97.
- Celik A, Aydinlik N and Arslan I 2011. Phytochemical constituents and inhibitory activity towards methicillin-resistant *Staphylococcus aureus* strains of *Eryngium* species (Apiaceae). *Chem. Biodivers* **8**: 454-9.
- Clinical and Laboratory Standards Institute 2006. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically, 7th ed. approved standard M7-A7: CLSI, Wayne, PA.
- Ecevit Genç G and Özhatay N 2006. An ethnobotanical study in Çatalca (European part of İstanbul) II. *Turk. J. Pharm. Sci.* **3**(2): 73-89.
- Ertuğ F 2000. An ethnobotanical study in central anatolia (Turkey). *Econ. Bot.* **54**(2): 155-182.

- Ferreira AV, Prado CG, Carvalho-Dias KST and Dias ALT 2013. *Candida albicans* and non-*C. albicans* species: comparison of biofilm production and metabolic activity of biofilms, and putative properties of isolates from hospital environments and infections. *Mycopathol.* **175**: 265-72.
- Freshney RI 2000. Cytotoxicity. *In: Culture of animal cells. A manual of basic technique*, 4th ed, pp. 329-345. Wiley-Liss, USA.
- Gumus I 1994. Ağrı yöresinde yetişen bazı faydalı bitkilerin yerel adları ve kullanılışları. *Turk. J. Bot.* **18**: 107-112.
- Kartal M, Mitaine-Offer AC, Abu-Asaker M, Miyamoto T, Calis I, Wagner H, and Lacaille-Dubois MA 2005. Two new triterpene saponins from *Eryngium campestre*. *Chem Pharm Bull (Tokyo)*. **53**: 1318-20.
- Kupeli E, Kartal M, Aslan S and Yesilada E 2006. Comparative evaluation of the anti-inflammatory and antinociceptive activity of Turkish *Eryngium* species. *J. Ethnopharmacol.* **107**(3): 32-37.
- McDaid HM and Johnston PG 1999. Synergistic interaction between paclitaxel and 8-chloro-adenosine 3',5'-monophosphate in human ovarian carcinoma cell line. *Clin. Cancer Res.* **5**: 215-220.
- Mekhora C, Muangnoi C, Chingsuwanrote P, Dawilai S, Svasti S and Chasri K (Eds) 2012. *Eryngium foetidum* suppresses inflammatory mediators produced by macrophages, *Asian Pac. J. Cancer Prev.* **13**: 653-664.
- Meot-Duros L, Le Floch G and Magné C 2008. Radical scavenging, antioxidant and antimicrobial activities of halophytic species. *J Ethnopharmacol.* **116**(2): 258-262
- Ndip RN, Tarkang AEM, Mbulah SM, Luma HN, Malongue A and Ndip LM (Eds) 2007. *In vitro* anti-*Helicobacter pylori* activity of extracts of selected medicinal plants from North West Cameroon. *J. Ethnopharmacol.* **114**(3): 452-457.
- Özgökçe F and Özçelik H 2004. Ethnobotanical aspects of some taxa in east Anatolia, Turkey I. *Econ Bot.* **58**(4): 697-704.
- Öztürk N and Dinç M 2005. Nizip (Aksaray) bölgesinin etnobotanik özellikleri. *Ot Sistematik Botanik Dergisi.* **12**(1): 93-102.
- Park JG, Lee SK, Hong IG, Kim HS, Lim KH and Choe KJ (Eds) 1994. MDR1 gene expression: Its effect on drug resistance to doxorubicin in human hepatocellular carcinoma cell line. *JNCI: Journal of the National Cancer Institute.* **86**: 700-5.
- Paul JH, Seaforth CE and Tikasingh T 2011. *Eryngium foetidum* L.: A review. *Fitoterapia* **82**: 302-8.
- Pimenov MG and Leonov MV 1993. The genera of the Umbelliferae: A nomenclator. 1th ed. Royal Botanic Gardens: Kew.
- Sezik E, Yeşilada E, Honda G, Takaishi Y, Takeda O, and Tanaka T 2001. Traditional medicine in Turkey X. folk medicine in central anatolia. *J. Ethnopharmacol.* **75**: 95-115.
- Sezik E, Yesilada E, Tabata M, Honda G, Takaishi Y and Fujita T (Eds) 1997. *Econ. Bot.* **51**(3): 195-211.
- Sharifa AA, Neoh YL, Iswadi MI, Khairul O, Abdul Halim M and Jamalubin M (Eds) 2008. Effects of methanol, ethanol and aqueous extract of plantago majoron gram positive bacteria, gram negative bacteria and yeast. *Annals of Microscopy* **8**: 42-44.
- Tegos G, Stermitz FR, Lomovskaya O and Lewis K 2002. Multidrug pump inhibitors uncover remarkable activity of plant antimicrobials, *Antimicrob. Agents Chemother* **46**: 3133-3141.
- Thiem B, Goślińska O, Kikowska M, and Budzianowski J 2010. Antimicrobial activity of three *Eryngium* L. species (Apiaceae). *Herba polonica* **56**(4): 52-8.
- Tuzlacı E 2005. Geçmişten günümüze Bodrum'da bitkiler ve yaşam. 1th edition, pp. 1-251. Güzel Sanatlar Matbaası, İstanbul.
- Yusuf, AJ, Hana S and Abdu S 2002. Antimycotic activities of selected plant flora, growing wild in Lebanon, against phytopathogenic fungi. *J. Agric. Food Chem.* **50**(11): 3208-3213.
- Zhang Z, Li S, Ownby S, Wang P, Yuan W and Zhang W (Eds) 2008. Phenolic compounds and rare polyhydroxylated triterpenoid saponins from *Eryngium yuccifolium*. *Phytochemistry* **69**(10): 2070-80.