

## ANALYSIS ON THE ACCUMULATION OF ORIDONIN IN DIFFERENT PORTIONS OF *ISODON RUBESCENS* (HEMSLEY) H. HARA

JIAN ZAIYOU\*, XU GUIFANG AND DAI LEI

*Henan Institute of Science and Technology, Xinxiang, China, 453003*  
*Henan Collaborative Innovation Center in Modern Biological Breeding, 453003, China*

*Keywords: Isodon rubescens, Oridonin, Accumulation, RP-HPLC*

### Abstract

Analysis on the accumulation of oridonin in different parts of *Isodon rubescens* was studied. The contents of oridonin in all parts of *I. rubescens* were analyzed with RP-HPLC. The results showed that the differences between the contents of oridonin in the different parts of *I. rubescens* are extremely significant. The content of oridonin in the leaf of *I. rubescens* is higher than that of other parts. The content of oridonin in the stem was close to that of leaf. The root of *I. rubescens* had lowest content of oridonin in the three portions. But the content of oridonin in the root obtained was 0.0811 mg/g. The root in long run should be utilized to avoid wasting the *I. rubescens* resources.

### Introduction

It is set in the Chinese Pharmacopoeia (2015 Ed.) that *Rabdosiae Rubescentis Herba* (traditional Chinese medicine) is the dry aerial portions of *Isodon rubescens* (Hemsl.) H. Hara (Chinese Pharmacopoeia 2015 Ed.). *I. rubescens* belongs to Labiatae family (Flora of China 1979). There are much resources of *I. rubescens* in the Taihang Mountain in China. *Rabdosiae Rubescentis Herba* are used as traditional Chinese medicine for the treatment of sore throat, inflammation and gastrointestinal problems ((Sun *et al.* 2006). Oridonin is a bioactive chemical component in *I. Rubescens*, and has potent *in vitro* and *in vivo* activity against human cancer cells (Ikezoe *et al.* 2003, Leung *et al.* 2005 and Bai *et al.* 2010). Growers of medicinal plants usually harvest the aerial portions of *I. rubescens* and then dry it in the sun before sale as *Rabdosiae Rubescentis Herba*. *I. rubescens* is subshrub plant with 3 - 4 years of life. The root of *I. rubescens* is very flourishing. The weight of *I. rubescens* root is usually larger than that of its aerial parts. The reports about medicinal compositions in the root of *I. rubescens* are scanty. Therefore, the value of *I. rubescens* root is not clear. In addition, there are differences between the contents of medicinal compositions in *I. rubescens* stem and those in its leaf. In this study, the difference between the accumulation of oridonin in different portions of *I. rubescens* was analysed to fully exploit *I. rubescens* resources and reasonably utilize *Rabdosiae Rubescentis Herba*.

### Materials and Methods

Shimadzu HPLC-2010 instrument, shimadzu (C18 reverse-phase column, 5  $\mu$ m, 250  $\times$  4.6 mm), lectronic analytic balance (precision: 0.0001), ultrasonator and rotary evaporator were used. Methanol (AR), ethanol (AR) and acetonitrile (HPLC grade) were used as reagents. Standard oridonin (99.8 %) were purchased from Sichuan Weikeqi Biotechnology Co. Ltd. in China in June 2017.

Thirty plants of *I. rubescens* were randomly dug in Guanshan of Xinxiang city in Henan province China in July, 2018. The leaves, stems and roots of these plants were separated and dried to get constant weight at 40°C.

The dry leaf, stem and root of *I. rubescens* were respectively crushed and sieved with 80 meshes sieve. Each material was weighed for 2 g and extracted with 25 ml ethanol solvent (75%)

\*Author for correspondence: <jian19732004@126.com>.

in the ultrasonic bath for 30 min. The mixture was filtered with filter paper. The residue was extracted with the same solvent (25 ml of 75% ethanol) and filtered once again. This filtrate was merged and added to 50 ml. The extract was filtered with 0.22  $\mu\text{m}$  membrane filter. The extraction of each kind material was repeated three times. Standard oridonin solutions were prepared at 0.001, 0.005, 0.025, 0.05 and 0.15, mg/ml respectively.

The Diamonsil C18 reverse-phase column (5  $\mu\text{m}$ , 250  $\times$  4.6 mm) was used as HPLC column. The temperature in HPLC column was 35°C. The volume of extract injected was 10  $\mu\text{l}$ . The gradient mobile phase consists of acetonitrile and water. The content (v/v) of acetonitrile in the gradient mobile phase varied from 25 to 29 % in 0 - 10 min, 29 % in 10 - 15 min and 29 to 30 % in 15 - 20 min. The flow rate of mobile phase was 0.8 ml/min. A variable wavelength recorder was set at 238 nm to detect ingredients eluted from the column.

These standard solutions and prepared extracts were respectively analyzed according to the above HPLC method. Chromatography peak areas of oridonin in each chromatogram were respectively recorded. These contents of oridonin in extracts were analyzed according to their chromatography peak areas and the standard curves (relating these peak areas to their contents). All of the data were analyzed with SPSS (Statistical Product and Service Solutions).

### Results and Discussion

The HPLC chromatogram of standard oridonin is presented in Fig. 1. The retention time of oridonin is 14.304 min.

The standard curve of oridonin is set up according to the contents and their corresponding peak areas (Table 1 and Fig. 2). The adopted standard curve of oridonin is  $y = 10020360.7899 \times + 685.1315$  (x: Concentration, y: Peak area,  $R^2 = 0.9999$ ).




Fig. 1. HPLC chromatograms of standard oridonin.

The peaks of oridonin in extract chromatograms were identified according to their retention time in HPLC (Fig. 3). The concentrations of oridonin in extracts were analyzed according to their peak areas and standard curves (Table 2). The contents of oridonin in *I. rubescens* materials were analyzed according the methods of preparation extract.

**Table 1. HPLC analysis results of standard oridonin.**

Concentration (mg/ml)	Peak area (Retention time 14.304 min)
0.001	10950
0.005	50983
0.025	251273
0.05	500956
0.15	1503967





Fig. 2. Standard curves of oridonin.

Fig. 3. HPLC chromatograms of *I. rubescens* materials.

The results showed that there was large difference between the contents of oridonin in the different portions of *I. rubescens*. Then the variance and multiple comparisons on these contents of oridonin in the different portions of *I. rubescens* were analysed (Table 2).

The differences between the contents of oridonin in different portions of *I. rubescens* were extremely significant ( $p < 0.01$ ). The content of oridonin in the leaf of *I. rubescens* is higher than that of other portions. The stem of *I. rubescens* owns close content of oridonin to that of leaf. The root of *I. rubescens* has lowest content of oridonin in the three portions. But the content of oridonin in the root achieve 0.0811 mg/g.

**Table 2. Contents of oridonin in *I. rubescens* materials.**

Part	Peak area	Concentration (mg/ml)	Content (mg/g)	Multiple comparisons*
Leaf	1332583	0.132918953	3.322974	
Leaf	1115786	0.111283315	2.782083	3.106 <sup>a</sup>
Leaf	1288220	0.128491669	3.212292	
Stem	502622	0.050091534	1.252288	
Stem	571380	0.05695336	1.423834	1.385 <sup>b</sup>
Stem	593876	0.059198388	1.47996	
Root	35001	0.003424473	0.085612	
Root	30327	0.002958023	0.073951	0.0811 <sup>c</sup>
Root	34271	0.003351622	0.083791	

\*The mean difference is significant at the 0.01 level. The different letters indicate, there is obvious difference between these means, The same letters indicate there is not obvious difference between these means.

The dry aerial portions of *Isodon rubescens* is used as *Rabdosiae Rubescentis Herba* which is specified in the Chinese Pharmacopoeia (Chinese Pharmacopoeia 2015 Ed.). There are very a few published reports about the accumulation of oridonin in different portions of *Isodon rubescens* (SU *et al.* 2009). The result of present study and other published reports similarly showed that the contents of oridonin in the aerial portions (especially leaf) of *Isodon rubescens* are all higher than that of root (SU *et al.* 2009). The content of oridonin in the root achieve 0.0811 mg/g. Therefore, the root of *I. rubescens* has considerable medicinal value. The root would be exsiccated in 3- 4 years if it be harvested for *I. rubescens* possesses 3 - 4 years of life. The root in long age should be utilized to avoid wasting the *I. rubescens* resources.

### Acknowledgement

This study was supported by Science and Technology Research Project of Henan Province (No. 182102310634 & No. 152300410093).

### References

- Bai NS, He K, Zhou Z, Tsai ML, Zhang L, Quan Z, Shao X, Pan MH and Ho CT 2010. Ent-Kaurane diterpenoids from *Rabdosia rubescens* and their cytotoxic effects on human cancer cell lines. *Planta Med.* **76**(2): 140-145.
- Chinese Pharmacopoeia Committee 2015. Pharmacopoeia of the People's Republic of China, 2015 ed, Part I. Chemical Industry Press, Beijing, 106-107.
- Flora of China Committee 1979. Flora of China (volume 66). Science Press, Beijing, 457-458.
- Ikezoe T, Chen SS, Tong XJ, Heber D, Taguchi H and Koeffler HP 2003. Oridonin induces growth inhibition and apoptosis of a variety of human cancer cells. *Int. J. Oncol.* **23**(4): 1187-1193.
- Leung CH, Grill SP, Lam W, Han QB, Sun HD and YC Cheng 2005. Novel mechanism of inhibition of nuclear factor-kappa B DNA-binding activity by diterpenoids isolated from *Isodon rubescens*. *Mol. Pharm.* **68**: 286-297.

- Sun HD, Huang SX and Han QB 2006. Diterpenoids from *Isodon* species and their biological activities. Nat. Prod. Rep. **23**: 673-698.
- SU Xiuhong, LI Xueju, DONG Chengming and Wang Weili 2009. Study on the dynamics of spatial accumulation of oridonin and rosmarinic acid of *rabdosia rubesens* (Henmsl.) Hara. China Medical Herald **6**(21): 5-6.

*(Manuscript received on 16 October, 2018; revised on 18 February, 2019)*