

Homozygous mutation of OCLN gene in Pseudo-TORCH Syndrome Type I : A Case Report

Naznin Akter¹, Iffat Ara Shamsad²

Abstract:

Clinical similarities of TORCH syndrome but serological negative findings for an infectious agent named it Pseudo-TORCH syndrome, which follows autosomal recessive inheritance but it has genetic heterogeneity. Here we reported a case of 11 month old girl of consanguineous parent with psychomotor retardation, microcephaly and intracranial calcifications but she was serologically negative for TORCH infection and positive for homozygous mutation in the gene encoding occluding (OCLN) on chromosome 5.

DOI: <https://doi.org/10.3329/bjch.v46i2.72125>

Introduction:

Pseudo- TORCH syndrome is a rare disorder of autosomal recessive inheritance where the patient presents with variable manifestations that resemble congenital intrauterine infection.¹ Intracranial calcification and microcephaly are the important clue to detect TORCH syndrome (toxoplasmosis, rubella, cytomegalovirus, herpes, syphilis) here pseudo - TORCH syndrome mimic the clinical and neuroradiologic features of TORCH infection in the absence of bacteriological or virological evidence of infection.² The name of this disease was first described by Baraister et al. who reported two boys of same family presented with spastic tetraplegia, microcephaly, seizure and intracranial calcifications.³

Case study:

Baby Afifa a 13-month old girl of consanguineous parents presented with the complaints of recurrent seizure since 3 months of age, in the form of clonic movement of limbs and blinking of eye involving the left half of body occurring 3-4 times in a day. Mother also complained that her child was unable to stand alone, able to recognize her mother. She was delivered normally at hospital and cried immediately after birth. Her developmental milestone was delayed, she achieved her neck control at 6 months of age

and started to sit unsupported since 10 months of age, and she can speak only baba and dada. Her parents were maternal cousins and family history of epilepsy was positive among the family members. Her one maternal uncle and one paternal aunt have been suffering from epilepsy since their childhood, both of them were well controlled with antiepileptic drugs. Baby Afifa has been treated with antiepileptic drug phenobarbital for the last 6 months without remission of seizure. On examination OFC 41 cm, which lies below 3rd centile, no facial dysmorphism except convergent squint on both eyes and has no neurocutaneous markers on skin survey. Vitals were stable during examination. Anthropometrical examination reveals her weight is 8 kg, weight for age lies just above 15th centile, supine length is 65 cm, length for age just on 10th centile. Hypertonia present on four limbs. All deep tendon reflexes are brisk without clonus. Her motor developmental age corresponds 6 months, speech and cognition corresponds 9 months of age. Her vision and hearing were intact.

TORCH screening was negative, S. ammonia, Blood lactate, CBG, S electrolytes, ABG all reports were within normal levels, IMD panel was normal. EEG showed focal epileptiform discharge from the left frontoparietal region of the brain with poverty of normal activity in background. Hearing assessment refers to no hearing impairment. Eye evaluation reveals normal anterior and posterior chambers. CT scan of the brain reveals intracranial calcification with cerebral atrophy. Clinical exome sequencing shows:

1. Assistant Professor, Department of Pediatric Neurology, Dhaka Medical College, Dhaka.

2. Professor of Pediatrics, Dhaka Medical College, Dhaka.

Correspondence: Dr. Naznin Akter, Assistant Professor, Department of Pediatric Neurology, Dhaka Medical College, Dhaka. Cell no: 01712552740, email: nazninruby73@gmail.com

Received: 16/02/2022

Accepted: 19/05/2022

Gene	Location	Variant	Zygoty	Disease(OMIM)	Inheritance	Classification
OCLN(+) ENST00000355237.6	Exon 3	c.90del p.Gly31Glu ufsTer34	Homozygous	Pseudo-TORCH syndrome - 1	Autosomal Recessive	Pathogenic
F SCN3A(-) ENST00000283254.12	Exon23	c.4184A>C p.Asn139 5Thr	Heterozygous	Developmental and epileptic encephalopathy-62	Autosomal Dominant	Uncertain Significance
PAK1(-) ENST00000278568.8	Exon 4	c.398C>T p.Ser133 Leu	Heterozygous	Intellectual developmental disorder with macrocephaly, seizures and speech delay	Autosomal Dominant	Uncertain Significance



According to the results of next generation sequencing patient was finally diagnosed as a case of Epileptic encephalopathy due to Pseudo-TORCH Syndrome-1, we managed the patient by multidisciplinary approach, for seizure oral form of levetiracetam was prescribed and at the same time speech therapy, developmental and occupational therapy was started and now patient is under our supervision with good seizure control and there is progressive improvement of neurocognitive behavior.

Discussion:

Pseudo-TORCH syndrome is one of the rare clinical syndromes that mimics the presentation of congenital TORCH infection without sero positivity and it follows the autosomal recessive inheritance.³ The common presentation of pseudo-TORCH syndrome are congenital microcephaly, developmental delay, recurrent

seizure, neuroradiology, intracranial calcification, simplified gyration and polymicrogyria.⁴ Here our case presented with recurrent seizures and delayed developmental milestones with family history of epilepsy. Our patient had no history of perinatal asphyxia. One study reported two patients with Pseudo-TORCH syndrome from a consanguineous family with complex heterogeneous presentation like early onset seizure, severe psychomotor development, visual impairment, chronic renal dysfunction, calcifications in thalami, basal ganglia and subcortical white matter, sanger sequencing identified a homozygous genomic rearrangement involving the OCLN gene.⁵ O’Driscoll et al.⁴ also reported their patient with some renal impairment Reardon et al.⁶ have reported patients with hepatosplenomegaly, platelet dysfunction and purpura along with seizure and microcephaly which mimics the presentation of TORCH infection, but in our case we

didn't find any renal, hematological or hepatobiliary abnormality. Eye manifestations like nystagmus, cortical visual impairment, corneal clouding, congenital cataract and microphthalmia was documented from different studies, we have only found convergent squint as eye manifestations.⁷⁻⁹ Facial dysmorphism in the form of hypertelorism, microphthalmia, micrognathia, high arched palate, low set ears and sloping forehead was observed by Monastiri et al.⁹ and Abdel-Salam et al.¹⁰ Different cases presents with different manifestations with clinic-radiological findings similar to congenital CMV infections but was negative for serology. One cases of pseudo-TORCH syndrome presented as Dandy-Walker syndrome which is a hind brain deformity with large posterior fossa cyst, this presented was explained in that way that extensive calcium debris from pseudo-TORCH syndrome in cerebrospinal fluid lead to obstruction at fourth ventricle outlet that leads to hydrocephalus and enlargement of posterior fossa cyst with cerebellar compression.³ So the main difference of DWM from pseudo-TORCH syndrome is the presence of features of raised ICP in the DWM. Aicardi-Goutieres syndrome (AGS) is another inherited form of microcephaly which may mimic the presentation of pseudo-TORCH syndrome, in most of the cases it follows autosomal recessive inheritance.¹¹ Spastic quadriplegia, progressive microcephaly, profound psychomotor retardation with bilateral symmetrical extensive calcifications in the basal ganglion is the prominent features of AGS.^{12,13} The identified gene mutation differs from pseudo-TORCH syndrome; the most common gene mutations identified in AGS are TREX1 gene, RNASEH2A, RNASEH2B and RNASEH2C.¹⁴ Whereas the most common identified genetic mutations are the homozygous or compound heterozygous mutations in the OCLN gene located on chromosome 5q13 in pseudo-TORCH syndrome 1, USP18 gene mutation on chromosome 22q11 associated with pseudo-TORCH syndrome 2 and STAT2 gene on chromosome 12q13 mutation is associated with pseudo-TORCH syndrome 3.^{2,4,10} Along with homozygous mutation of OCLN gene there are another two heterozygous mutations occurring in our cases for that reason our patient presented with recurrent seizure. Though the patients showed clinical and radiologic features of congenital CMV infection but she was negative for serology and due to presence of marital consanguinity we went for genetic study and finally the patient was diagnosed as a case of pseudo-TORCH syndrome. We managed the patient symptomatically by multidisciplinary approach and counseled them about the course of the disease.

References:

1. Watts's P., Kumar N., Ganesh A. et al. Choriorretinal dysplasia, hydranencephaly, and intracranial calcifications: pseudo-TORCH or a new syndrome? *Eye*, 2008; 22:730-33
2. Briggs TA, Wolf NI, D'Arrigo S, Ebinger F, Harting I, Dobyns WB, et al. Band like calcification with simplified gyration and polymicrogyria: A distinct "pseudo-TORCH" phenotype. *Am J Med Genet A*. 2008;146A: 3173-80.
3. Barister M, Brett EM, Piesowicz AT. Microcephaly and intracranial calcification in two brothers. *J Med Genet*, 1983;20:210-12.
4. Patnaik A, Mishra SS, Das S. Extensive intracranial calcification of pseudo-TORCH syndrome with features of Dandy-Walker malformation. *Asian J Neurosurg*. 2017;12: 541-43.
5. O'Driscoll MC, Daly SB, Urquhart JE, Black GCM, Pilz DT, Brockmann K et al. Recessive mutations in the gene encoding the tight junction protein occluding cause band-like calcification with simplified gyration and polymicrogyria. *Am J Hum Genet*. 2010;87: 345-64.
6. LeBlanc MA, Penny LS, Gaston D, Shi Y, Aberg E, Nightingale M et al. A novel rearrangement of occluding causes brain calcification and renal dysfunction. *Hum Genet*, 2013;132:1223-34.
7. Reardon W, Hockey A, Silberstein P, Kendall B, Farag TI, Swash M et al. Autosomal recessive congenital intrauterine infection like syndrome of microcephaly, intrauterine calcification and CNS disease. *Am J Med Genet*. 1994;52: 58-65.
8. Burn J, Wickramasinghe HT, Harding B, Baraitser M. A syndrome with intracranial calcification and microcephaly in two sibs, resembling intrauterine infection. *Clin. Genet*. 1986;30:112-16.
9. Monastiri K, Salem N, Korbi S, Snoussi N. Microcephaly and intracranial calcification: two new cases. *Clin Genet*. 1997;51:142-3.
10. Slee J, Lam G, Walpole I. Syndrome of microcephaly, microphthalmia, cataracts and intracranial calcification. *Am J Med. Genet*. 1999;84:330-33.
11. Abdel-Salam GMH, Zaki MS, Saleem SN, Gaber KR. Microcephaly, malformation of brain development and intracranial calcification in sibs: pseudo-TORCH or a new syndrome. *Am J Med Genet*. 2008;146:2929-36.
12. Aicardi J, Goutieres F. A progressive familial encephalopathy in infancy, with calcifications of the basal ganglia, and chronic cerebrospinal fluid lymphocytosis. *Ann Neurol*. 1984;15:49-54.
13. Giroud M, Gouyon JB, Chaumet F, Cinquin AM, Alison M, Dumas R. A case of progressive familial encephalopathy in infancy with calcification of the basal ganglia and chronic cerebrospinal fluid lymphocytosis. *Childs Nerv Syst*. 1986;2: 47-48.
14. Kumar D, Rittey C, Cameron AH, Variend S. Recognizable inherited syndrome of progressive central nervous system degeneration and generalized intracranial calcification with overlapping phenotype of the syndrome of Aicardi and Goutieres. *Am J Med Genet*. 1998;75: 508-15.
15. Rice G, Patrick T, Parmar R, Taylor CF, Abey A, Aicardi J, Artuch R et al. Clinical and molecular phenotype of Aicardi-Goutieres syndrome. *Am J Hum Genet*. 2007;81: 713-25.