


# The development of speed and speed-strength qualities in short-distance runners in complicated training conditions

Polevoy G.G. 

## ABSTRACT

### Background

The problem of developing the speed of movement and speed-strength qualities of sprinters can be solved by applying special exercises in artificially difficult training conditions.

### The purpose of the study

is to explore the possibilities of using special training tools, in particular running in artificially difficult conditions at the stage of special physical training of sprinters of mass categories.

### Materials and methods

The study involved young men aged 17-19 years old in the number of 28 people who were engaged in sprint running in the athletics section at the sports school of the Olympic reserve of the city of Kirov (Russia). The pedagogical experiment began on November 5, 2022 and ended on May 24, 2023. By the method of random sampling, all athletes were differentiated into a control group that practiced according to the standard methodology for training sprinters in natural conditions of difficulty and an experimental group in which sprinters practiced in artificially created conditions of difficulty. When processing the results of the study, the Student's T-test was used.

### Results

After the end of the pedagogical experiment, the indicators in both groups improved. The sprinters from the control group had higher scores from 1% to 3% in all tests ( $p>0.05$ ). In the athletes from the experimental group, from the beginning to the end of the study, the indicators improved from 4.5% to 5.9% on all tests ( $p<0.05$ ). This proves the effectiveness of using a set of exercises and load components in working with athletes in artificially created difficult exercise conditions.

### Conclusion

If you use a special set of exercises in artificially created difficult conditions when working with sprinters aged 17-19, then the indicators of speed and strength abilities and speed of movement will significantly improve in comparison with athletes who train in natural conditions of difficulty.

### Keywords

long-term training; young athletes; speed qualities; sprinters; training.

## INTRODUCTION

Currently, the system of training athletes has a multifactorial influence, that is, tasks, means and methods to achieve the highest possible results<sup>1,2</sup>.

Training in children is a great incentive to achieve success and achieve goals<sup>3,4</sup>. Systematic physical education and sports are of great importance for the development of physical and moral qualities of children<sup>5,6</sup>.

Sports training is a long-term plan in the implementation of the set goals. During the year, the athlete strengthens his health, solves motor tasks, develops psychologically, develops tactical skills and knowledge in the field of physical education<sup>7,8</sup>.

In planning, it is important to analyze and identify the weaknesses and strengths of athletes and competitive work. It is important to present a clear running model in competitive work. This is important in order to obtain adequate means and methods of training<sup>9,10</sup>.

Based on the data available in the scientific and methodological literature, the construction of a one-year training cycle for sprinters has the following form<sup>2,11</sup>:

1. The general preparatory stage (November-February) is 11-13 weeks (improvement of general physical fitness and development of

## Correspondence

Polevoy G.G. Department of Physical Education, Moscow Polytechnic University, Moscow Russia.  
E-mail: [g.g.polevoy@gmail.com](mailto:g.g.polevoy@gmail.com).

basic physical qualities).

2. Special preparatory stage (March-May) 11-12 weeks (improvement of special physical training, improvement of equipment and development of elements of tactics, improvement of speed-strength and special running training).

3. The competition period (June-July) is 6-8 weeks (achieving a high level of special training and its implementation in the most responsible competitions).

4. The transition period (August-October) is 11-12 weeks (load reduction, active rest, recovery, it is necessary to create conditions for maintaining physical fitness in order to start a new large cycle from a level exceeding the previous one)

According to experts and coaches, the intensity of work should sometimes exceed the intensity required at competitions. For example, in a sprint, this task can be solved with the help of exercises performed in difficult, ordinary and lightweight conditions, combined in a certain sequence<sup>12</sup>.

The ways to complicate the conditions for performing a competitive exercise can be divided into two groups:

1) natural conditions of difficulty – running on sand, running uphill, running on snow, on water, running on stairs;

2) artificially created conditions of difficulty – running with negative traction (with rubber shock absorbers or rubber resistance fixed on the lower part of the shin, thigh; with a parachute; with a tire; with a winch – technical devices that impede movement, automatically regulating the force of traction), running with a vest or belt (up to 8% of body weight), running in cuffs attached to the lower part of the shin, running with the resistance of a partner<sup>13</sup>.

The frequent change of resistance caused by bilateral action contributes to the greatest manifestation of the speed parameters of movements with weights<sup>11,14</sup>.

In order to increase the speed of movement with a weighted resistance, under the influence of working with a light one, a constant alternation of resistances is necessary. After working with weighted resistance, you can perform exercises with increased speed in two or three movements with light resistance<sup>15</sup>.

The use of weighted and lightweight weights makes it possible to selectively influence the increase in the level of use of individual components of special speed

and strength qualities (speed or strength) and allows you to dramatically increase the volume of special exercises<sup>12,16</sup>.

It is known that if we use methods to complicate the conditions for performing a competitive exercise, the number of throwing exercises per year of training will increase 6-10 times for javelin throwers, 2-2.5 times for disco balls, as well as 2-3 times for hammer throwers<sup>14,17</sup>.

A sprinter's strength training program can take the form of weight training, or isometric training, weighted vests, shoes, strength training with a partner, or gymnastic exercises. It is necessary that running with weights be combined with running under normal conditions. This gives a significantly greater increase in running speed than just training in a sprint. The use of resistances makes it possible to increase the frequency and efficiency of leg movements (per second) and increase the strength of the muscles involved in sprinting<sup>13-15</sup>.

The sprinter's speed and strength training began to be carried out not so much by working with a barbell, but by jumping exercises and "power" running, that is, running in difficult conditions. It is important that our training movements be a mirror image of competitive work<sup>2,11,16</sup>.

Running in difficult conditions allows you to maintain the basic structure of a competitive exercise. But there is a problem related to the qualitative and quantitative measurement of variability. This requires deeper and more specific development. There is no consensus on which sequence is best followed when applying complex and ordinary conditions.

There are several studies in the modern scientific literature proving the effectiveness of using one or another technique to develop the physical qualities of athletes in difficult conditions<sup>2,13,16,17</sup>. However, we have presented for the first time a comparative analysis of special learning tools in natural and artificially complicated conditions, which is the novelty of this study.

The hypothesis of the study is that it can be assumed that the use of special means in training, such as weights for various segments, will increase the speed and transience of movement of sprinters aged 17-19 years.

The aim is to explore the possibilities of using special training tools, in particular running in artificially complicated conditions, at the stage of special physical training of sprinters of mass categories.

## Tasks

1. To study the features of using special training equipment for sprinters in accordance with scientific and methodological literature.
2. To develop and test a special set of exercises in artificially complicated conditions for sprinters aged 17-19 years, to propose load components.
3. Compare the results obtained with previous studies on this problem.

## MATERIALS AND METHODS

### *Study participants:*

The study involved young men aged 17-19 years old in the number of 28 people who were engaged in sprint running in the athletics section at the sports school of the Olympic reserve of the city of Kirov (Russia). All athletes underwent an in-depth medical examination and were admitted by a group of doctors to training and competitive activities.

All procedures met the ethical standards of the 1964 Declaration of Helsinki and were approved by the local university ethics committee.

### Organization of the study:

Since November 5, 2022, the general preparatory stage has begun for all athletes of the sports school. At the first lesson, all athletes took control tests. The selection of control tests for the study of the motor training of sprinter athletes was carried out taking into account the information available in the literature on the factorial, correlation significance of tests and their reliability<sup>18-20</sup>:

1. Long jump from a place (assessment of speed and strength qualities);
2. Triple jump from a place (assessment of speed and strength qualities);
3. Running 20 meters per stroke (evaluation of the sprinter's maximum running speed);
4. Running 60 meters from a low start (assessment of speed abilities).

Until February 28, 2023 (before the end of the general preparatory stage), all athletes practiced according to the same methodology and performed the same amount and intensity of load. All athletes used the means to develop the physical qualities necessary for a sprinter<sup>2,13,14,17</sup>:

1. Accelerations of 50-60 meters × 6-8 times.
2. Acceleration of 60-80 meters × 3-4 times.
3. Acceleration of 40 meters × 8-10 times.
4. Acceleration from a low start to 40 meters × 8-10 times.
5. Throwing from various starting positions (weight 3-4 kg).
6. Running on soft ground 2 × 150 meters (freely).
7. Running exercises with a 2 kg belt.
8. Jumping from a place in length 12-15 times.
9. Jumping with getting branches 15-20 times.
10. Variable running -80 meters fast after 20 meters jogging.
11. Running uphill 5 × 100 meters.
12. Running up the stairs.
13. Running down the mountain 3-4 times for 60-80 meters.
14. Jumping exercises of 30-40 meters each exercise, 3-4 repetitions.
15. Repeated running with an intensity of 85-90%.
16. General physical training exercises with weights (25-40 kg): pushes, pulls, jerks, lunges in motion, jumps with a change of leg position – each exercise is 2-3 series 8-10 times.

From March 1 to May 24, 2023, the main pedagogical experiment was conducted at a special preparatory stage.

By random sampling, all children were differentiated into 2 groups of 14 athletes each.

The athletes of the control group practiced according to the generally accepted method, using special training tools in natural conditions of difficulty<sup>11,14,16</sup>: a set of jumping exercises at the stadium, running up stairs, extreme running on sand, running uphill, running in the snow.

The athletes of the experimental group trained according to the proposed method using special means in artificially created difficult conditions. An example of exercises in an experimental group:

1. Running and jumping in ankle cuffs (weight up to 500 grams)
2. Running and jumping in hip cuffs (weight up to

700-900 grams)

3. Running and jumping with a weight belt (weight 2-5 kg)
4. Running with a partner's resistance;
5. Running with a tire;
6. Running with holding on a rubber shock absorber.

Running and running exercises with weights were used in accordance with the basic methodological techniques:

- The length of the segments of running with weights usually does not exceed 50 meters
- The speed of overcoming the segments should be equal to 80-90% of the maximum capabilities of the athlete.
- The number of repetitions in one series – 2-3 times
- Rest between repetitions in a series – 3-4 minutes.
- The number of episodes per workout is 2-4.
- Rest between series of 6-8 minutes or until full recovery (when the heart rate drops to 114-120 beats / min.)

The development of speed in its complex expression is best achieved by using running on segments of 50-80 meters with maximum and near maximum speed.

In order to determine the effectiveness of the use of physical training in both groups, after the end of the special preparatory stage on May 25, 2023, all athletes re-passed the control standards.

*Methods of mathematical statistics*

After the end of the pedagogical experiment, methods of mathematical statistics were used, which are widely used in pedagogical research<sup>21,22</sup>.

The generally accepted characteristics of the static distribution were calculated:

$\bar{X}$  – arithmetic mean;

$\delta$  – standard deviation;

$m$  – standard error of the arithmetic mean;

When comparing the indicators, the reliability of the differences was determined by the Student's T-criterion at  $p \leq 0.05$ .

## ETHICAL CLEARANCE:

This research was conducted in compliance with the needed research ethics. In addition, consent for participation was obtained from the participants before the beginning of their involvement in the study. All data were recorded and analyzed anonymously.

## RESULTS

In order to objectively verify the effectiveness of the proposed method in comparison with the generally accepted method, it is necessary to identify changes in intra-group indicators of physical fitness, as well as their differences in level in groups at the beginning and at the end of the experiment (intergroup differences).

Table 1 shows the baseline values in the control and experimental groups for all tests.

**Table 1.** Comparison of the performance of athletes from both groups before the start of the study

| Control tests | Control group |          |      | Experimental group |          |      | t    | p     |
|---------------|---------------|----------|------|--------------------|----------|------|------|-------|
|               | X             | $\delta$ | m    | X                  | $\delta$ | m    |      |       |
| Jump          | 240,6         | 8,5      | 2,7  | 240,3              | 7,9      | 2,5  | 0,08 | >0,05 |
| Triple jump   | 709           | 31,5     | 9,97 | 707                | 34,7     | 10,9 | 0,14 | >0,05 |
| Run 20m       | 2,87          | 0,19     | 0,06 | 2,89               | 0,16     | 0,05 | 0,26 | >0,05 |
| Run 60m       | 7,76          | 0,32     | 0,1  | 7,84               | 0,32     | 0,1  | 0,57 | >0,05 |

The analysis of the initial data in Table 1 indicates the relative homogeneity of the groups, since before the experiment both groups were almost equal in terms of physical fitness. For all four indicators, the differences are unreliable ( $p > 0.05$ ).

After applying the proposed training tools at the stage of special physical training, repeated testing was carried out for the same indicators and the results were obtained, which are reflected in Tables 2 and 3.

**Table 2.** Comparison of the performance of athletes from the control group at the beginning and at the end of the study (n=14)

| Control tests | Before |          |      | After |          |      | t    | p     |
|---------------|--------|----------|------|-------|----------|------|------|-------|
|               | X      | $\delta$ | m    | X     | $\delta$ | m    |      |       |
| Jump          | 240,6  | 8,5      | 2,7  | 243   | 7,3      | 2,3  | 0,68 | >0,05 |
| Triple jump   | 709    | 31,5     | 9,97 | 730   | 20,5     | 6,5  | 1,76 | >0,05 |
| Run 20m       | 2,87   | 0,19     | 0,06 | 2,8   | 0,13     | 0,04 | 1    | >0,05 |
| Run 60m       | 7,76   | 0,32     | 0,1  | 7,6   | 0,02     | 0,01 | 1,6  | >0,05 |

Table 2 shows that when comparing the indicators for all tests in the control group from the beginning to the end of the study, no significant differences were found, but their positive growth should be noted. At the same time, significant changes occurred in the experimental group for all tests (Table 3).

**Table 3.** Comparison of the performance of athletes from the experimental group at the beginning and at the end of the study (n=14)

| Control tests | Before |          |      | After |          |      | t   | p     |
|---------------|--------|----------|------|-------|----------|------|-----|-------|
|               | X      | $\delta$ | m    | X     | $\delta$ | m    |     |       |
| Jump          | 240,3  | 7,9      | 2,5  | 253   | 6,3      | 2,5  | 3,7 | <0,05 |
| Triple jump   | 707    | 34,7     | 10,9 | 739   | 14,2     | 7,4  | 2,7 | <0,05 |
| Run 20m       | 2,89   | 0,16     | 0,05 | 2,72  | 0,13     | 0,04 | 2,8 | <0,05 |
| Run 60m       | 7,84   | 0,32     | 0,1  | 7,4   | 0,01     | 0,01 | 4,4 | <0,05 |


The analysis of Table 3 shows that from the beginning to the end of the study, the athletes in the experimental group had positive and significant changes in all control tests.

To illustrate the results of the experiment, they can be presented in the form of a diagram, all indicators from the beginning to the end of the study in both groups are presented in percentage terms in Figure 1.

Comparing the data in Figure 1 from the beginning to the end of the study in athletes of the control group and the experimental group and analyzing them, we can consider sufficient grounds to say that the use of special means in training in artificially difficult conditions is more effective than their use in normal and naturally difficult conditions.

## DISCUSSION

Before the start of the pedagogical experiment, the peculiarities of using special training tools for sprinters were studied. An analysis of the scientific and methodological literature on the problem has shown that with a properly organized training process, special training tools at the stage of special physical training can be quite effective. Such features include natural conditions of difficulty in performing the exercise, and artificially created difficult conditions for performing the exercise<sup>2,15,16</sup>.



**Figure 1.** The increase in indicators in both groups from the beginning to the end of the study.

For the first time, a special set of exercises in artificially difficult conditions for sprinters was developed and tested, specific load components for athletes aged 17-19 years were proposed.

According to the results of this pedagogical experiment, it was revealed that athletes from the control group who practiced according to the generally accepted method, using special training tools in natural conditions of difficulty, were able to improve their performance during the study period. In the “Long jump from a place” test, the indicators improved by an average of 2.4cm (1%), and in the “Triple jump from a place” test, the indicators became 21 cm higher (3%). In the test “Running for 20 meters from a stroke” improved by 0.07 seconds (2.4%), in the test “Running for 60 meters from a low start” the indicators improved by 0.16 seconds (2.1%). Despite the positive results in all tests in the control group from the beginning to the end of the study, there was no significant increase in indicators ( $p > 0.05$ ). Such results may indicate a relatively low, but effective application of a standard technique that uses special training tools in natural conditions of difficulty.

As for the athletes from the experimental group, during the study period they significantly and reliably improved their performance on all tests ( $p < 0.05$ ). For example, in the “Long Jump from a place” test, the indicators improved by an average of 12.7 cm (5.3%), and in the “Triple Jump from a place” test, the indicators became 32 cm higher (4.5%). In the test “Running 20 meters from a run”, the indicators became higher by 1.7 seconds (2.4%), and in the test “Running 60 meters from a low start”, the indicators improved by 4.4 seconds (2.1%). Such results certainly indicate the effectiveness of using

a special set of exercises in artificially created difficult conditions and the proposed load components for sprinters aged 17-19 years.

If we compare the results of the study with the methods that exist in the scientific literature, we can note the following.

It has been proven that sprinters include in their training not only exercises with weights, but also jumping exercises, running with a partner’s resistance, which gives a positive training effect and improves the competitive result<sup>13,14</sup>. Such results are confirmed by our research.

Most experts in the field of athletics note the great importance of using running in conditions of varying difficulty to change the speed of running in short stretches. After several jumps in difficult conditions, the speed of running under normal conditions increases from the very beginning more significantly than when training under the most normal conditions<sup>2,16,17</sup>. These results from the literature coincide with ours in the control group, where athletes improved their performance on all tests by training according to the generally accepted methodology, using special training tools in natural conditions of difficulty.

After running in sections of the distance with difficult conditions, in which the runner has to overcome the distance with great effort, at a relatively high pace and speed of movement, running under normal conditions follows. At the same time, special endurance is also improved to some extent, since extremely fast movements in this case are performed in more difficult conditions<sup>11,13,15</sup>. This is confirmed by our results of the pedagogical experiment, which were obtained in both groups, and a positive increase in indicators was recorded for all control standards.

In one study, short-distance runners used weights located at the distal ends of the lower leg. It can be assumed that the impact of loads will disrupt the sensorimotor connections of the multilevel movement system, and the transformed version of the movement system will be connected to the previously established structure of perceptual processes. However, after relieving the load, all athletes increased their running speed compared to the initial data and increased their running pace and reduced the repulsion time<sup>2,6,12</sup>. This is consistent with the results of the study in the experimental group, since athletes were able to significantly and reliably

improve their performance on all tests during the study period using special training tools in artificially created difficult conditions (running with a tire, running with a vest or belt with weights, running handcuffed on the shins or running with a partner's resistance, and others).

Thus, the assumption that the use of special means in training, such as weights for various segments, will increase the speed and transience of movement of sprinters aged 17-19 years, was fully confirmed by the results of the study.

However, despite the fact that the purpose of the study has been achieved and the hypothesis has been solved, the problem has the prospect of further study.

The physical qualities and physical development of school-age children are of great importance<sup>23-26</sup>. The problem of obesity in schoolchildren is relevant<sup>27-30</sup>. It should also be noted the leading role of physical education in solving the problem of obesity in children of different school ages<sup>31-35</sup>. All this indicates the relevance of the research and its further perspective.

## CONCLUSION

If you use a special set of exercises in artificially created difficult conditions when working with sprinters aged 17-19, then the indicators of speed and strength abilities and speed of movement will significantly improve in comparison with athletes who train in natural conditions

of difficulty.

Weights on the shins up to 500 grams of cuffs, on the hips up to 800 grams, a belt of 2-5 kg are recommended. Running with a tire or with a partner's resistance, running with a hold on a rubber shock absorber. Running with artificial traction – 7-8 kg. There are only 2-4 episodes in one training session. The intensity of running with weights should be 80-90% of the maximum capacity. Rest intervals should be sufficient for recovery. Rest for 3-4 minutes between repetitions, and between series until complete recovery. The results of the study are recommended to be used in the training process of sprinters of mass categories.

**Source of fund:** This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**Conflict of interest:** The author declares that he no conflict of interest.

## AUTHOR'S CONTRIBUTION:

Data gathering and idea owner of this study: Polevoy G.G.

Study design: Polevoy G.G.

Data gathering: Polevoy G.G.

Writing and submitting manuscript: Polevoy G.G.

Editing and approval of final draft: Polevoy G.G.

## References

- Rodriguez-Negro, J., Llodio, I., Castillo, D., Romaratezabala, E., & Yanci, J. Changes in selected locomotor skills of young runners after one athletic season: Influence of sex and age. *International Journal of Sports Science & Coaching*, 2021;**16**(5), 1152-1161. <https://doi.org/10.1177/17479541211006676>
- Milloz, M., Hayes, K. & Harrison, A.J. Sprint Start Regulation in Athletics: A Critical Review. *Sports Med* 2021; **51**:21–31. <https://doi.org/10.1007/s40279-020-01350-4>
- Christino MA, Coene RP, O'Neil M, et al. (). Sport specialization, athletic identity, and coping strategies in young athletes. *Orthopaedic Journal of Sports Medicine*, 2021;**7**(13): doi:10.1177/2325967121S00111
- Eric W. MacIntosh, Milena M. Parent & Diane Culver Understanding Young Athletes' Learning at the Youth Olympic Games: A Sport Development Perspective, *Journal of Global Sport Management*, 2022;**7**:1, 1-20, DOI: 10.1080/24704067.2018.1561206
- Pentidis, N., Mersmann, F., Bohm, S. et al. Effects of long-term athletic training on muscle morphology and tendon stiffness in preadolescence: association with jump performance. *Eur J Appl Physiol* 2020;**120**:2715–2727. <https://doi.org/10.1007/s00421-020-04490-7>
- Dijkstra, H. P., Pollock, N., Chakraverty, R., & Alonso, J. M.. Managing the health of the elite athlete: a new integrated performance health management and coaching model. *British journal of sports medicine*, 2014;**48**(7): 523–531. <https://doi.org/10.1136/bjsports-2013-093222>
- Staff, H.C., Solli, G.S., Osborne, J.O. et al. Long-Term Development of Training Characteristics and Performance-Determining Factors in Elite/International and World-Class Endurance Athletes: A Scoping Review. *Sports Med* 2023;**53**: 1595–1607. <https://doi.org/10.1007/s40279-023-01850-z>
- Zouita, A., Darragi, M., Bousselmi, M. et al. The Effects of Resistance Training on Muscular Fitness, Muscle Morphology, and Body Composition in Elite Female Athletes: A Systematic Review. *Sports Med* 2023; **53**: 1709–1735. <https://doi.org/10.1007/s40279-023-01859-4>
- Lundberg Zachrisson, A., Ivarsson, A., Desai, P. et al. Athlete availability and incidence of overuse injuries over an athletics season in a cohort of elite Swedish athletics athletes - a prospective study. *Inj. Epidemiol.* 2020; **7**: 16. <https://doi.org/10.1186/s40621-020-00239-0>
- Thompson, F., Rongen, F., Cowburn, I., & Till, K. The Impacts of Sports Schools on Holistic Athlete Development: A Mixed Methods Systematic Review. *Sports medicine (Auckland, N.Z.)*, 2022;**52**(8): 1879–1917. <https://doi.org/10.1007/s40279-022-01664-5>
- Haugen, T., Seiler, S., Sandbakk, Ø. et al. The Training and Development of Elite Sprint Performance: an Integration of Scientific and Best Practice Literature. *Sports Med - Open* 2019;**5**(44): <https://doi.org/10.1186/s40798-019-0221-0>
- Bezodis, N.E., Willwacher, S. & Salo, A.I.T. The Biomechanics of the Track and Field Sprint Start: A Narrative Review. *Sports Med* 2019;**49**: 1345–1364. <https://doi.org/10.1007/s40279-019-01138-1>
- Rauff, E. L., Herman, A., Berninger, D., Machak, S., & Shultz, S. P. Using sport science data in collegiate athletics: Coaches' perspectives. *International Journal of Sports Science & Coaching*, 2022;**17**(3):500-509. <https://doi.org/10.1177/17479541211065146>
- Jenkins AS, Pollock JR, Moore ML, Makovicka JL, Brinkman JC, Chhabra A. The 100 Most-Cited and Influential Articles in Collegiate Athletics. *Orthopaedic Journal of Sports Medicine*, 2022;**10**(7): doi:10.1177/23259671221108401
- Jones NS, Wieschhaus K, Martin B, Tonino PM. Medical Supervision of High School Athletics in Chicago: A Follow-up Study. *Orthopaedic Journal of Sports Medicine*, 2019;**7**(8):
- Swindell HW, Marcille ML, Trofa DP, et al. An Analysis of Sports Specialization in NCAA Division I Collegiate Athletics. *Orthopaedic Journal of Sports Medicine*, 2019;**7**(1): doi:10.1177/2325967118821179
- McGuigan, M.R., Cormack, S.J., & Gill, N. Strength and Power Profiling of Athletes: Selecting Tests and How to Use the Information for Program Design. *Strength and Conditioning Journal*, 2013;**35**: 7–14. <https://doi.org/10.1519/SSC.0000000000000011>
- Lockie, Robert G., Brett K. Post, and J. Jay Dawes. “hysical Qualities Pertaining to Shorter and Longer Change-of-Direction Speed Test Performance in Men and Women. *Sports* 2019;**7**(2): 45. <https://doi.org/10.3390/sports7020045>
- Chaabene, H., Negra, Y., Bouguezzi, R., Capranica, L., Franchini, E., Prieske, O., Hbacha, H., & Granacher, U. Tests for the Assessment of Sport-Specific Performance in Olympic Combat Sports: A Systematic Review With Practical Recommendations. *Frontiers in physiology*, 2018;**9**: 386. <https://doi.org/10.3389/fphys.2018.00386>
- Mishra, P., Singh, U., Pandey, C. M., Mishra, P., & Pandey, G. Application of student's t-test, analysis of variance, and covariance. *Annals of cardiac anaesthesia*, 2019;**22**(4): 407–411. [https://doi.org/10.4103/aca.ACA\\_94\\_19](https://doi.org/10.4103/aca.ACA_94_19)
- de Winter, J.C.F. () “Using the Student's t-test with extremely small sample sizes,” Practical Assessment, Research, and Evaluation: Vol. 18, Article 10. DOI: <https://doi.org/10.7275/e4r6-dj05>



23. GG, P. The influence of speed and power load on the indicators of the distribution of attention of schoolchildren with different typologies. *Bangladesh Journal of Medical Science*, 2019; **21**(3): 634–638. <https://doi.org/10.3329/bjms.v21i3.59578>
24. Koc, S., Garipağaoğlu, M., Ekinci, Özalp, Kanik, A., & Gültekin, F. Nutritional and Obesity Status of Children and Adolescents with ADHD: a case-control study. *Bangladesh Journal of Medical Science*, 2023;**22**(1):171–179. <https://doi.org/10.3329/bjms.v22i1.61874>
25. A, N. E., Yaacob, L. H., & Azidah, A. Pedometer-based walking intervention with and without group support among sedentary adults in primary care patients in north-east Malaysia: a randomized controlled trial. *Bangladesh Journal of Medical Science*, 2018;**17**(1): 52–57. <https://doi.org/10.3329/bjms.v17i1.35280>
26. Koliadenko, N. V., Zhyvaho, K. S., & Bursa, A. I. Provision of Medical-psychological and Psychiatric Care to Patients with Post-covid Syndrome in Telemedicine Conditions. *Bangladesh Journal of Medical Science*, 2022; **21**(4): 719–730. <https://doi.org/10.3329/bjms.v21i4.60256>
27. Zainol, J., & Salam, A. (). An Audit on Mentor-Mentee Program: Mentees Perceptions on Mentors. *Bangladesh Journal of Medical Science*, 2021;**20**(4): 840–847. <https://doi.org/10.3329/bjms.v20i4.54143>
28. Shantakumar, S. R., Sahabdeen, H. B., Zainal Abidin, F. A. B., Perumal, G., & Kumar, N. (). Association of type and duration of exercise with the mental and physical health and academic performance of Medical undergraduate students- Cross-sectional study. *Bangladesh Journal of Medical Science*, 2022; **21**(1): 135–139. <https://doi.org/10.3329/bjms.v21i1.56339>
29. Babalola, T. K., & Abasi, U. U. *angladesh Journal of Medical Science*, 2018; **17**(2): 270–274. <https://doi.org/10.3329/bjms.v17i2.35883>
30. Rabilal, M., Maharaj, S. S., & Kaka, B. . Efficacy of the Otago exercise programme as a supplementary intervention to exercise classes: On falls, balance, physical function and mobility in a patient with chronic lower back pain: A case report. *Bangladesh Journal of Medical Science*, 2020;**19**(2): 339–342. <https://doi.org/10.3329/bjms.v19i2.45020>
31. Koliadenko, N. V., Zhyvaho, K. S., & Bursa, A. I. Provision of Medical-psychological and Psychiatric Care to Patients with Post-covid Syndrome in Telemedicine Conditions. *Bangladesh Journal of Medical Science*, 2020;**21**(4): 719–730. <https://doi.org/10.3329/bjms.v21i4.60256>
32. Duran, S., & Can ÖZ, Y.. The relationships between Orthorexia nervosa, social appearance anxiety and women's self-esteem: A cross-sectional study. *Bangladesh Journal of Medical Science*, 2020;**21**(3): 675–681. <https://doi.org/10.3329/bjms.v21i3.59584>
33. Wibowo, R. A., Wasityastuti, W., & Sofro, Z. M. Low total physical activity, high total sitting time and high sitting time on a work day are correlated with low fitness in male working adults: a cross sectional study. *Bangladesh Journal of Medical Science*, 2029;**18**(2): 279–287. <https://doi.org/10.3329/bjms.v18i2.40698>
34. Anh, V. T. K., Huong, T. V., Dung, D. V., Khai, N. V., & Huy, T. Q. Working Motivation of Medical Staff and Some Related Factors at a General Hospital in the Mekong Delta, Vietnam in the Context of the COVID-19 Pandemic. *Bangladesh Journal of Medical Science*, 2023;**22**(1): 121–127. <https://doi.org/10.3329/bjms.v22i1.63076>
35. Mosalanejad, L., Razeghi, B., & Ifard, S. A. Educational Game: A Fun and team based learning in psychiatric course and its effects on Learning Indicators. *Bangladesh Journal of Medical Science*, 2018;**17**(4):631–637. <https://doi.org/10.3329/bjms.v17i4.38328>