

Outcome of Pituitary Adenoma Surgery Through Transphenoidal Endoscopic Endonasal Approach -A Study of 96 Cases

Nath HD¹, Razib KO², Halder R³, Rahman MA⁴, Hasan MM⁵, Uddin ANW⁶, Karim MR⁷, Barua S⁸, Munir SKF⁹, Barua KK¹⁰

Conflict of interest: There is no Conflict of interest relevant to this paper to disclose.

Funding Agency: Was not funded by any institute or any group.

Contribution to authors: Pro. Dr Haradhan Deb Nath, Dr Md Motasimul Hasan

Manuscript Preparation: Dr Rathin Halder, Dr Md Rabiul Karim

Data Collection: Dr KH. Olinur Razib, Dr Sudip Barua, SK Farhad Munir

Editorial formatting: Pro. Dr Haradhan Deb Nath, Dr Md Motasimul Hasan

Copyright: ©2022bang. BJNS published by BSNS. This article is published under the creative commons CC-BY-NC license. This license permits use distribution (<https://creativecommons.org/licenses/by-nc/4-0/>) reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Received: 17 January, 2023

Accepted: 8 February 2023

Abstract:

Background: Transphenoidal endoscopic approach is minimal invasive surgery in case of pituitary adenoma.

Objective : To see the outcome of transphenoidal endoscopic approach in case of pituitary tumour.

Results : This study showed among the 96 patients, 64 (66.7%) patients were male and 32(33.33%) were female. It was documented that in 96 patients, 88(91.66.3%) were macroadenoma and 8(8.33%) were microadenoma. Among 96 patients, 88 (91.66%) had preoperative visual disturbance and 8(8.33%) had normal vision. Tumor was totally removed in 88(91.66%) patients in endoscopic approach. Clinically 92(95.8%) patients were improved in endoscopically. Two (2.1%) patient developed pneumocephalus and meningitis, between them one died due to meningitis and other cured by conservative treatment. In 2 (2.1%) patient, carotid artery was partially torn, which was managed by nasal pack.

Conclusion: It was concluded that endoscopic endonasal approach is safe and effective procedure.

Key word: Pituitary adenoma, transphenoidal approach, endoscopic endonasal approach, acromegally, cushing's syndrome.

Bang. J Neurosurgery 2023; 12(2): 29-33

Introduction:

The role of the pituitary gland has intrigued scientists and doctors for many centuries. It was only in 1887 that its clinical role was discovered by Minkowski, one year after acromegaly was described by Pierre Marie.¹ The first pituitary tumor operations were performed at the end of the 19th century. Transsphenoidal surgical techniques have been

improving since 1907 when Schioffer performed the first successful transnasal operation in Vienna. In the USA,² Hirsch (1910) and cushing (1912)³ contributed to the evolution of transsphenoidal surgery and had surgical mortality rates as low as 9.5 and 4.8% without the use of cortisone or antibiotics. Cushing initially used a head light and otologic mirrors but eventually discontinued trans-sphenoidal surgery because of

1. Pro. Dr. Haradhan Deb Nath, Professor, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka.
 2. Dr. KH Olinur Razib, Resident, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka
 3. Dr. Rathin Halder, Medical Officer, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka,
 4. Dr. Md. Aatur Rahman, Registrar, Department of Neurosurgery, SSMC Mitford Hospital, Dhaka.
 5. Dr. Md. Motasimul Hasan, Associate Professor, Department of Neurosurgery, Dhaka Medical College & Hospital, Dhaka.
 6. Dr. Abu Naim Wakil Uddin, Medical Officer, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka.
 7. Dr Md. Rabiul Karim, Assistant Professor, Department of Neurosurgery, Chittagong Medical College Hospital, Chittagong.
 8. Dr Sudip Barua, Medical Officer, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka.
 9. Dr. SK Farhad Munir, Research Assistant, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka.
 10. Prof. Kanak Kanti Barua, Professor (Rtd.), Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka.
- Address of Correspondence:** Dr. KH. Olinur Razib, Resident, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka. Mobile: 01727374275, E-mail: olinurrazib1084@gmail.com

insufficient vision on the surgical field. Later, Guiot and Derome (1967), in Paris, and Hardy (1969), in Montreal, again popularized the transsphenoidal approach by introducing the surgical microscope, intraoperative radiographs and image intensification. With further technical improvements in neurosurgery and the introduction of antibiotics and cortisone, surgical mortality dropped under 1%. The recent introduction of endoscopy has reduced nasal complications and improved patient comfort. The endoscope had already been used by G. Guiot during transsphenoidal surgery more than 40 years ago but the pure endoscopic approach to pituitary lesions was introduced by Jankowski and Jho in the mid 1990s.¹

Tumors are classified according to size and invasiveness, which is predictive of surgical results. There is no consensus on the use of a universal classification but one which has stood the test of time was proposed by Hardy in 1976. This classification is based on modifications of the sella turcica on plain x-ray studies and is now adapted to MRI findings. The tumors are divided in four groups as follows.⁵

Grade I tumors, or microadenomas, are lesions up to 10mm in diameter.

Grade II tumors are more than 1 cm in diameter and cause diffuse enlargement of the sella turcica with or without suprasellar extension. These non-invasive macroadenomas remain enclosed in the anatomical structures containing the pituitary gland (diaphragma sellar, medial wall of the cavernous sinus, dura of sellar floor and dorsum sellar) that can be displaced but not invaded.

Grade III tumors are adenomas which show signs of local invasion. Some of them can be smaller than 10mm (invasive microadenomas). Grade IV tumors show extensive and diffuse invasion of the skull base and/or the intracranial space.¹

Radiologic assessment of parasellar invasion presents the problem of distinguishing whether the inner wall of the cavernous sinus has been infiltrated or simply displaced.⁶

Methods:

It was a prospective cross sectional study. Sampling technique was purposive consecutive. Total 48 cases of pituitary adenoma by transsphenoidal endoscopic approach from January 2010 to December 2019 at

the Department of Neurosurgery, Bangababdu Sheikh Mujib Medical University, Dhaka. Patients with neuro-images suggesting pituitary adenoma. Patient with pituitary adenoma who had visual impairment, hormonal disturbance and requiring surgery for it. Patient who had post-operative histological confirmation of pituitary adenoma were included in this study. Exclusion criteria are patients who were done reoperation, patients who had concomitant intra-ocular disease making visual assessment difficult, systemic disorders other than pituitary adenoma that affected visual function, presence of any other intracranial pathology. Data were collected pre-designed data collection sheet. Data were analysis using computer based programme statistical package for social science (SPSS) for windows version 16.

Operation procedure

Under general anaesthesia with supine position head is raised slightly from body. Nose was parallel to the floor, head was 20° flexed and rotated to left shoulder, surgeon will stand towards the right shoulder. Disinfectant was used providone iodine soaked gauze. Killian type nasal speculum and Hardy nasal speculum were used. The middle turbinate was identified and passage was made between the narrow space of middle turbinate and nasal septum. Ostium of sphenoidal air sinus and choana were identified.

Coagulation was done at bony part of nasal septum. Bilaterally nasal mucosa was dissected. Keel of the vomer was identified and was removed. Anterior wall of sphenoidal air sinus, sinus cavity, sinus mucosa and posterior wall of sphenoidal air sinus was removed. Dura was opened after coagulation and tumour was excised by suction ring curette, and micro ronger. Sellar floor was reconstructed and nasal pack was given with ribbon gauze and merocele.

Results:

We did transsphenoidal microscopic operation of 96 cases from January 2010 to May 2022.

Table-I
Distribution of the patients by sex (n=96)

Sex	Number	Percentage (%)
Male	64	66.66
Female	32	33.33
Total	96	100.00

It was evident that male 64(66.66%) predominate the female 32(33.33%).

Table-II
Distribution of the patients by age (n= 96)

Age	Number	Percentage (%)
1-20	2	2.08
21-40	50	52.08
41-60	30	31.25
>60	14	14.58

It was documented that 21-40 years belonged to the highest age group 50(52.08%)

Table-III
Distribution of the patients by size of the tumor (n=96)

Pituitary adenoma	Number	Percentage (%)
Microadenoma	8	8.33
Macroadenoma	88	91.66

It was evident that the majority of the tumour were macroadenoma 88(91.66%).

Table-IV
Distribution of the patients by visual field defect, pituitary apoplexy and seller size (n=96)

Visual disturbance	Number	Percentage (%)
Present	88	91.66
Absent	8	8.33
Seller size		
Normal	20	20.83
Enlarged	76	79.16
Pituitary apoplexy		
Present	24	25
Absent	72	75

It was evident that majority of the patients had visual disturbance 88(91.66%) and 76(79.16%)enlarged sella.

Table-V
Distribution of the patients by functional types of tumor (n=96)

Functional Type	Number	Percentage (%)
Prolactinoma	4	4.16
Acromegally	20	20.83
Cortisol secreting tumor	2	2.08
Non functional tumor	70	72.91

It was documented that majority of the tumor were non functional 70(72.91%).

Table-VI
Distribution of the patient by extent of removal of tumor (n=96)

Extent of removal	Number	Percentage (%)
Total removal	88	91.16
Subtotal removal	8	8.33

It was evident that majority of the tumour 88(91.16%) were totally removed.

Table-VII
Distribution of the patients by postoperative complication (n=42)

Complication	Number	Percentage (%)
Diabetes insipidus	30	71.42
Meningitis	2	4.76
Cerebrospinal fluid leakage (CSF)	6	14.28
Pneumocephalus	2	4.76
ICA partially torn	2	4.76

Table-VIII
Distribution of the patients by postoperative visual outcome and hormonal status (n=96)

Visual outcome	Number	Percentage (%)
Improved	92	95.83
Same as before	4	4.16
Hormonal status (n=13)		
Improved	14	53.84
Same as before	10	38.46
Detoriated (Permanent DI)	2	7.69

It was documented that majority 92 (95.83%) the patients improve after surgery

Discussion:

A endoscopic transsphenoidal approach can be used for about 95% of all pituitary adenomas. Tumors with significant anterior extension into the anterior cranial fossa or lateral extension into the middle fossa, either a transcranial approach consisting of a pteroinal or a subfrontal craniotomy or a combination of both these approaches is required. In dumbbell-shaped tumors with both a significant intrasellar component and a suprasellar-intracranial extension separated by a narrow neck, endoscopic transsphenoidal surgery can first be tried to decompress the visual pathways. Postoperative imaging will indicate if a sub-sequent transcranial approach will be required.⁷

Since the mortality rate is very low when the endoscopic transsphenoidal approach is used by experienced surgeons (less than 1%), priority must be given to this technique in difficult cases.⁸ In our study 20(66.67%) had diabetes insipidus. The situation must be re-evaluated after this first surgical step. Repeated transsphenoidal resections can be proposed to decompress any residual mass. This is because intracranial pressure will cause the residual suprasellar tumor to progressively descend into the seller in the weeks following surgery.

Combined approaches and staged procedures have to be considered in difficult cases as do alternative treatment modalities such as tumor-suppressive drugs and radiation therapy. In the case of very fibrous adenomas that cannot be removed by the transsphenoidal approach, care should be taken to avoid worsening of visual function by manipulation of the optic nerves and optic chiasm during subsequent cranial surgery.⁵ In our study one (6.66%) case was fibrous variety and abandoned due to excessive bleeding later operations was done by transcranial approach.

Mortality and morbidity related to hypothalamic ischemia, chiasmatic manipulations and uncontrolled bleeding during cranial surgery for large pituitary adenomas is mentioned in the literature.⁴

As a corridor can be safely drilled into the sphenoid bone under navigation or fluoroscopic control during the endoscopic procedure, transsphenoidal surgery can be performed on young children (> 5 years) or when the sphenoid sinus is poorly pneumatized.⁴

A small-normal sized sella turcica in the presence of a macrotumor with suprasellar extension is not suited for transsphenoidal surgery and implies reconsidering the diagnosis of pituitary adenoma. Using fully endoscopic transnasal approach, endocrinological remission rates in non-invasive functioning adenomas are about 75 to more than 80%. Results achieved in non-invasive grade 2 macroadenomas are similar to those obtained in microadenomas.⁵

The invasive character of the tumor is a factor influencing the results negatively: correction of hormonal hypersecretion in parasellar invasive macroadenomas is only exceptionally observed.⁶

Visual improvement is nearly always obtained after good decompression of the visual pathways from below, using a transsphenoidal approach, unless a

long preoperative evolution has resulted in optic atrophy.⁸ In our study among 95.83% with visual disturbance improved.

Careful tumor resection should avoid postoperative hypopituitarism; however, literature mentions new anterior pituitary insufficiency in up to 14% of the cases. Postoperative transient diabetes insipidus is the most frequent complication and generally recovers in 48 hours. Using endoscope, postoperative CSF leakage rates vary from 1.2 to 6%.¹⁰ In our study 30(71.42%) patients had postoperative diabetes insipidus and improved.

Complications can be due to a wrong track causing damage to the optic nerve perforating the cribriform plate or penetrating the cavernous sinus and damaging the with nerve or the carotid artery. Among the 42 cases 6(14.28%) patients had CSF. Frequent fluoroscopic or navigation controls to check the anatomical landmarks during the surgical approach are recommended.⁸

Hypothalamic damage, CSF leak and diaphragmatic breach or hypopituitarism can result from too aggressive tumor resection. Gentle handling of smooth ring curettes and avoiding any traction on the tumor or the surrounding structures should prevent these complications.⁶

Nasal and sinuses infectious complications will be avoided by repositioning the nasal septum and the middle turbinate at the end of surgery to ensure good ventilation. The use of prophylactic antibiotics for 48 hours is recommended.⁵

Pituitary adenomas represent more than 10% of intracranial tumors and are a frequent cause for neurosurgical consultation. Pituitary diseases show multiple and different pathological and clinical aspects, requiring multidisciplinary diagnostic approaches and multimodality treatment options including endocrinology, surgery and radiotherapy. Despite recent improvements in medical therapy, surgery still has an important place in the treatment of these lesions and requires well-trained pituitary surgeons. In experienced hands, pituitary surgery and especially endoscopic transsphenoidal surgery is safe with very low morbidity.^{7,10}

Vieira have treated 270 pituitary tumors, all macroadenomas and locally invasive; there were malignant tumors. All had been previously treated one or more times by some other modalities. Microsurgery alone was used in 90.3%, radiation therapy and micro-surgery in 8.2%, and radio therapy alone in

1.5%. Tumor volume ranged from 0.9 to 32 mL, with an average volume of 11 mL. In our study 88(91.16%) case tumor was completely removed and 8(8.33%) patients had subtotal tumour removal.¹¹

Tumors were treated with a maximum dose of to 60 Gy (average, 37.5 Gy). Periphery dose ranged from 3 to 28 Gy (average, 15 Gy). This discrepancy is caused by variations in patient profile. Patients that had received previous radiation therapy or that had a tumor close to the optic apparatus in general received lower doses. Currently, Gamma Plan, the planning software for the gamma knife, allows for very specific shielding of the optic nerves and chiasm, and this has allowed for higher dose administered when these structures are nearby.¹¹

Previous authors have treated 92 nonsecretory pituitary tumors, 82 of which have radiographic and endocrinologic follow-up of a minimum of 6 months and an average of 34 months. Of these, 55(67%) had a decrease in the volume of their tumors; 21 (26%) had no change in the size; and 6 (7%) increased in size. New hypopituitarism occurred in 12 patients (15%). The only indication we have to date for treating these tumors is for postoperative residual tumors, in order to lower the incidence of tumor progression or pro-gression in spite of previous radiation therapy.¹¹

Conclusion:

Transphenoidal endoscopic removal of tumour is one of the safe procedures.

Reference:

1. Cappabianca P, Cavallo LM, de Divitiis E. Endoscopic endonasal transphenoidal surgery. *Neurosurgery* 2004;55: 933-940; discussion 940-941.
2. Cavallo LM, Dal Fabbro M, Jalalod'din H, Mcssina A, Esposito I, Esposito F, de Divitiis E, Cappabianca P. Endoscopic endonasal transphenoidal surgery. Before scrubbing in: tips and tricks. *Surg Neurol* 2007;67: 342-347.
3. Fahlbusch R, Buchfelder M (1988) Transsphenoidal surgery of parasellar pituitary adenomas. *Acta Neurochir (Wien)* 1988;92: 93-99.
4. Frank G, Pasquini E, Farneti G, Mazzatenta D, Sciarretta V, Grasso V, Faustini Fustini M. The endoscopic versus the traditional approach in pituitary surgery. *Neuroendocrinology* 2006; 83: 240-248.
5. Hardy J, Vezina JL. Transsphenoidal neurosurgery of intracranial neoplasm. *Adv Neurol* 1976;15: 261-273.
6. Kassam A, Thomas AJ, Snyderman C, Carrau R, Gardner P, Mintz A, Kanaan H, Horowitz M, Pollack IF. Fully endoscopic expanded endonasal approach treating skull base lesions in pediatric patients. *J Neurosurg* 2007; 106: 75-86.
7. Laws ER. Surgery for acromegaly: evolution of the techniques and out-comes. *Rev Endocr Metab Disord* 2008;9: 67-70.
8. Ludecke DK, Abe T. Transsphenoidal microsurgery for newly diagnosed acromegaly: a personal view after more than 1000 operations. *Neuroendocrinology* 2006;83: 230-239.
9. Nieman LK, Biller BM, Findling JW, Newell-Price J, Savage MO, Stewart PM, Montori VM. The diagnosis of Gushing's syndrome: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 2008; 93: 1526-1540.
10. Shahinian HK. Endoscopic skull base surgery. Humana Press, Totowa 2008.
11. Vieira JO Jr, Cukiert A, Liberman B. Evaluation of magnetic resonance imaging criteria for cavernous sinus invasion in patients with pituitary adenomas: logistic regression analysis and correlation with surgical findings. *Surg Neurol* 2006; 65:130-135.