



**BJP**

**Bangladesh Journal of Pharmacology**

**Research Article**

*In vitro* anthelmintic activity of *Heliotropium indicum*, *Senna fistula* and *Spigelia anthelmia* used as worm expeller in South West Nigeria

## ***In vitro* anthelmintic activity of *Heliotropium indicum*, *Senna fistula* and *Spigelia anthelmia* used as worm expeller in South West Nigeria**

**Oluwakemi K. Sobiyi and A.O. Tom Ashafa**

*Phytomedicine and Phytopharmacology Research Group, Department of Plant Sciences, University of the Free State, Qwaqwa Campus, Phuthaditjhaba 9866, South Africa.*

### Article Info

Received: 15 March 2015

Accepted: 26 April 2015

Available Online: 13 May 2015

DOI: 10.3329/bjp.v10i2.22589

### Cite this article:

Sobiyi OK, Ashafa AOT. *In vitro* anthelmintic activity of *Heliotropium indicum*, *Senna fistula* and *Spigelia anthelmia* used as worm expeller in South West Nigeria. Bangladesh J Pharmacol. 2015; 10: 417-22.

### Abstract

The anthelmintic potential of *Heliotropium indicum*, *Senna fistula* and *Spigelia anthelmia* were investigated using nematodes larvae from sheep and adult earthworms in order to justify the folkloric claim of the plants as worm expeller in south west of Nigeria. The acetone, ethanol, hydro-alcohol and distilled water extracts showed dose-dependent anthelmintic activities at the different concentrations (0.3, 0.5, 1.0 mg/mL) when tested against nematodes larvae. The order of anthelmintic effect for the plants was *H. indicum*>*S. fistula*>*S. anthelmia*. Acetone and ethanol extracts of the three plants showed the most effective activity (100% mortality) against adult earth worm (*Pheretima posthuma*) after 30 min of exposure. The reference standard drug (Vermox®) showed less effectiveness compared to the medicinal plants used in the study. Overall, the study indicates that *H. indicum*, *S. fistula* and *S. anthelmia* are potential anthelmintic herbal drugs, which in turn validates the use of the three species as worm expellers by the Yorubas tribe of South West Nigeria.

## Introduction

*Trichiuris trichiuria*, *Ascaris lumbricoides* and *Ancylostoma duodenale* have been reported to be most prevalent in developing world where there is poor sanitation facilities and inadequate water supply (Savioli, 2004). In Africa, intestinal cestode infection is of great concern and it is closely linked with poor environmental hygiene and poverty (Carabinet al., 2006; Abunnaet al., 2007). Intestinal parasites have been long recognised among Nigerian children between the ages 0-18 years as a major health problem (Ajayi and Akinyinka, 1999; Oguoma et al., 2008), in which majority of the children were found to be stunted with retarded growth, anaemic and underweight due to malnutrition.

*Heliotropium indicum* is highly valued in the folklore medicine in the treatment of different diseases (Muthuet al., 2006; Togolaet al., 2005). *Senna fistula* is used as purgative having active ingredients anthraqui-

none derivatives and their glycosides. It is also used as a laxative acting on the lower bowel especially in alleviating constipation (Spiller et al., 2003). *Spigelia anthelmia* L. is known as 'Aparan' or 'ewe aran' (Dalziel, 1937), among the Yoruba tribe of south western Nigeria. According to Ezikeet al. (2013), the native inhabitants of south western Nigeria typically use *S. anthelmia* for expelling worms.

Chemotherapy is used to control helminthic infection in Nigeria; however the drugs are not readily available to the dwellers of the aforementioned areas (Mbaria et al., 1998). This study presents an alternative approach for treating intestinal worms in order to validate the folkloric claims about these plants as worm expellers.

## Materials and Methods

### Plant sampling








Figure 1: Comparative percentage mortality of 0.25 mg/mL (A), 0.5 mg/mL (B) and 1.0 mg/mL (C) of the extracts against adult earth worm (*Pheretima posthum*)

activity than Vermox®. There was 100% mortality rate at higher concentrations (1 mg/mL) of each plant used. At 30 min, the 0.3 mg/mL extracts of the three plants exhibited 55–100% mortality on the larvae while at 0.5 mg/mL, 70–100% mortality was recorded (Table I). Acetone, ethanol and hydro-alcohol extracts of the three plants showed 100% effectiveness as potential anthelmintic at 0.5 mg/mL concentration at 1 hour of observation. At 1 hour, the water extract of *H. indicum* and *S. fistula* showed better anthelmintic property compared to *S. anthelmia* having mortality rate of (80, 90, 100%), (90, 95, 100%) and (70, 85, 90%) respectively.

The earthworms selected for the anthelmintic activity were most sensitive to acetone and ethanol root extracts of all the three plants (Table II). There was 100%

mortality at all the three concentrations at a minimum of 30 min of exposure. *H. indicum* exhibited better anthelmintic activity with the hydro-alcohol extract than those of *S. fistula* and *S. anthelmia*. There was no death observed at lower concentrations (0.3 and 0.5 mg/mL) in all the three plant water extracts for all the time(s) the test was carried out. At 1.0 mg/mL, *H. indicum* had 50% mortality on the earthworms while it was 75% for both *S. fistula* and *S. anthelmia* respectively at this concentration which indicated that significant anthelmintic activity is concentration dependent. 25% mortality was only observed at 1.0 mg/mL after 3 hours in the positive control (Vermox®) while all the earthworms in distilled water (negative control) were living and active after 3 hours.

There was a significant difference ( $F_{2,3}= 11$ ,  $p<0.05$ ) between the plants at the lowest concentration of 0.3 mg/mL used in the study (Figure 1A). There was no anthelmintic activity observed at the least concentration of *S. anthelmia* hydro-alcohol plant extract and from distilled water extract of the three plants against the earthworms. At the 0.5 mg/mL, there was also a significant difference ( $F_{2,3}= 17.4$ ,  $p<0.05$ ) observed for the three plants (Figure 1B) and this significant difference ( $F_{2,3}= 6.1$ ,  $p<0.05$ ) was equally observed at the highest concentration (1.0 mg/mL) used in the study (Figure 1C). *S. anthelmia* showed the least effect with hydro-alcohol extract while *H. indicum* was observed to have the least anthelmintic activity with the distilled water extract.

## Discussion

The result from the anthelmintic activity suggests that all the three plants used for the study are potent worm expeller and this agrees with the work of Sofowora, (2006) that ethno-medicinal prescription of plant for worm eradication had achieved great success with almost 63% of tropical plants showing anthelmintic properties.

The anthelmintic activity of the three plants carried out using gastrointestinal nematode larvae of sheep showed a 100% death rate in ethanol and acetone after 30 min of observation at 0.5 mg/mL concentration while for the hydro-alcohol and distilled water extracts this mortality was observed at the concentration 1.0 mg/mL also after 30 min. This confirms the folkloric claim that the plants are more effective when soaked into the local gin (more or less alcohol) to kill/expel intestinal worms, this is also in agreement with Eloff, (1998) that the isolation of botanical compounds from plant materials largely depends on the medium or solvent and method of extraction.

Earthworms have been widely used for *in vitro* initial evaluation of anthelmintic compounds due to its easy

availability (Sollmann, 1918; Sundeep Kumar et al., 2010). The three plants showed 100% mortality in both acetone and ethanol extract for the times (30 min - 3 hours) of observation. It can, therefore, be inferred that ethanol and acetone were able to extract more of the constituents that is required to kill the worms, this result exhibited the same pattern with the work of Chavan et al. (2010), where the *in vitro* anthelmintic activity of the fruit extract of *Barleeia prionitis* was tested against adult earthworm and a better result was observed with the ethanol extract over the water extract at time 7.1 min of observation. In hydro-alcohol extracts, *H. indicum* was observed to be most effective while in distilled water extracts, *S. anthelmia* showed the highest effect against the Indian earthworms. The effect of the anthelmintic activity of *S. anthelmia* root is consistent with the report of Ezike et al. (2013), in which the whole plant methanol extract used against earthworm showed best potential anthelmintic effect of the three plants used for the study, he further claimed that the effect could be accounted for due to its alkaloids constituent as such compounds have been associated with anthelmintic activity (Makkaret al., 2007). *H. indicum* has been proven by Fu et al. (2002), to contain tumorigenic pyrrolizidine alkaloids (normal cell that are transformed into cancer cells) and should not be consumed in larger quantity. The vermifugal activity of the extracts against earthworms suggests that these plants would be effective against human intestinal parasitic infections.

---

## Conclusion

The use of acetone, ethanol and hydro-alcohol extracts are recommended for better effect of anthelmintic activity of *H. indicum*, *S. fistula* and *S. anthelmia* as there was less or almost no effect was observed with the use of distilled water extract. *H. indicum* and *S. fistula* are recommended to be used as human intestinal worm expeller instead of *S. anthelmia* which is widely known in Nigeria.

---

## Acknowledgements

This research was supported by the Postgraduate School Bursary of the University of the Free State. The authors are also grateful to the research Committee UFS-Qwaqwa Campus for funding (211427604). We thank Ms. Getrude Mahanke (Research Assistant) for her support during all experiments.

---

## References

Abunna F, Tilahun G, Megersa B, Regassa A. Taeniasis and its socio-economic implication in Awassa town and its surroundings, Southern Ethiopia. *East Afr J Public Health*. 2007; 4: 73-79.

- Ajaiyeoba EO, Onocha P, Olarenwaje OT. *In vitro* anthelmintic properties of *Buchholzia coriacea* and *Gynandropis gynanadra* extract. *Pharm Biol*. 2001; 39: 217-20.
- Ajayi EOS, Akinyinka OO. Evaluation of the nutritional status of first year school students in Ibadan South. *West Afr J Med*. 1999; 28: 59-63.
- Ashafa AOT, Afolayan AJ. Screening the root extracts from *Biden pilosa* L. var. *radiata* (Asteraceae) for anti-microbial potentials. *J Med Plants Res*. 2009; 3: 568-72.
- Ashok Kumar BS, Lakshman K, Jayaveera KN, Nandeesh R, Manoj B, Ranganayakulu D. Comparative *in vitro* anthelmintic activity of three plants from the *Amaranthaceae* family. *Arch Biol Sci Belgrade*. 2010; 62: 185-89.
- Carabin H, Krecek RC, Cowan LD, Michael L, Foyaca-Sibat H, Nash T, Willingham AL. Estimation of the cost of *Taeniasolium* cysticercosis in Eastern Cape Province, South Africa. *Trop Med Int Health*. 2006; 11: 906-16.
- Chatterjee KD. Parasitology, protozoology and helminthology. Calcutta, Guha Ray Sree Saraswaty Press Ltd, 1967, pp 168-69.
- Chavan CB, Hogade MG, Bhinge SD, Kumbhar M, Tamboli A. *In vitro* anthelmintic activity of fruit extract of *Barleeia prionitis* L against *Pheretima posthuma*. *Int J Pharm Pharmacol Sci*. 2010; 2: 49.
- Dalziel JM. The useful plants of West Tropical Africa. The crown agents for the colonies. London, 1937.
- Eloff JN. Which extractant should be used for the screening and isolation of antimicrobial components from plants? *J Ethnopharmacol*. 1998; 60: 1-8.
- Ezike AC, Akah PA, Okore V, Okoli CO, Okoye TC, Okoye AC. *In vitro* evaluation of the anthelmintic and antibacterial activities of three Nigerian medicinal plants. *Bio Med Res*. 2013; 1: 254-57.
- Fu PP, Yang YC, Xia Q, Chou MC, Cui YY, Lin G. Pyrrolizidine alkaloids-tumorigenic components in Chinese herbal medicines and dietary supplements. *J Food Drug Anal*. 2002; 10: 198-11.
- Makkar HPS, Sidhuraju P, Becker K. Plant secondary metabolites. New Jersey, USA., Humana Press Inc, 2007.
- Margono SS, Wandura T, Swasono MF, Murni S, Craig PS, Ito A. Taeniasis/cysticercosis in Papua (Irian Jaya), Indonesia. *Parasitol Int*. 2006; 55: S143-48.
- Martinez M, Ramo JD, Torreblanca A, Diazmayans J. Effect of cadmium exposure on zinc levels in the brine shrimp *Artemia parthenogenetica*. *Aquaculture* 1999; 172: 315-25.
- Mbaria JM, Maithe TE, Mitema ES, Muchiri DJ. Comparative efficacy of pyrethrum marc with albendazole against sheep gastrointestinal nematodes. *Trop Anim Health Prod*. 1998; 30: 17-22.
- Muthu C, Ayyana M, Raja N, Ignacimuthu S. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. *J Ethnobiol Ethnomed*. 2006; 2: 43.
- Oguoma VM, Anyasodor AE, Mbata TI. Prevalence of intestinal protozoan parasites among children aged 5-12 years in Owerri Metropolis South Eastern Nigerian. *Trop J*

- Health Sci. 2008; 15: 43-48.
- Savioli L, Albonico M. Soil-transmitted helminthiasis. *Nat Rev Microbiol.* 2004; 2: 618-19.
- de Silva NR, Brooker PJ, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminth infections: Updating the global picture. *Trends Parasitol.* 2003; 19: 547-51.
- Sofowora AB. Medicinal plants and traditional medicine in Africa. 3rd ed. Nigeria, Spectrum Book Ltd, 1993, pp 195-238.
- Sofowora AB. Medicinal plants of West Tropical Africa. Lagos, Sanders Publishers Ltd, 2006, pp 45-57.
- Sollmann T. Anthelmintics: Their efficiency as tested on earthworms. *J Pharmacol Exp Ther.* 1918; 12: 129-70.
- Spiller HA, Winter MI, Weber JA, Krenzelok EP, Anderson DL, Ryan ML. Skin breakdown and blisters from senna-containing laxatives in young children. *Annals Pharmacother.* 2003; 37: 636-39.
- Sundeeep Kumar HK, Bose A, Raut A, Sahu SK, Raju MBV. Evaluation of anthelmintic activity of *Pistia stratiotes* Linn. *J Basic Clin Pharm.* 2010; 1: 103-05.
- Togola A, Diallo D, Demb'el'e S, Barsett H, Paulsen BS. Ethnopharmacological survey of different uses of seven medicinal plants from Mali, (West Africa) in the regions Doila, Kolokani and Siby. *J Ethnobiol Ethnomed.* 2005; 1: 7.
- Vidyarthi RD. A textbook of Zoology. New Delhi, 1967, S Chand and Co, pp 329-70.
- 

**Author Info**

A. O. Tom Ashafa (Principal contact)  
e-mail: ashafaot@qwa.ufs.ac.za

Table II

Effect of Vermox®, *Heliotropium indicum*, *Senna fistula* and *Spigelia anthelmia* root extracts on earthworms (*Pheretima posthuma*)

Test sample	Conc. (mg/mL)	Con. tol 30 min	% mortality																										
			Vermox						Acetone extract						Ethanol extract						Hydro-alcohol extract						Distilled water		
			1 hour	2 hours	3 hours	30 min	1 hour	2 hours	3 hours	30 min	1 hour	2 hours	3 hours	30 min	1 hour	2 hours	3 hours	30 min	1 hour	2 hours	3 hours	30 min	1 hour	2 hours	3 hours				
<i>Heliotropium indicum</i>	0.25	0	0	0	100	a	a	a	100	a	a	a	100	a	a	a	75	100	b	b	b	0	0	0	0				
	0.5	0	0	0	100	a	a	a	100	a	a	a	100	a	a	a	100	a	a	a	a	0	0	0	0				
	1.0	0	0	25	100	a	a	a	100	a	a	a	100	a	a	a	100	a	a	a	a	25	50	50	50				
<i>Senna fistula</i>	0.25	0	0	0	100	a	a	a	100	a	a	a	100	a	a	a	0	0	25	50	50	0	0	0	0				
	0.5	0	0	0	100	a	a	a	100	a	a	a	100	a	a	a	50	50	75	100	100	0	0	0	0				
	1.0	0	0	25	100	a	a	a	100	a	a	a	100	a	a	a	100	a	a	a	a	50	50	75	75				
<i>Spigelia anthelmia</i>	0.25	0	0	0	100	a	a	a	100	a	a	a	100	a	a	a	0	0	0	0	0	0	0	0	0				
	0.5	0	0	0	100	a	a	a	100	a	a	a	100	a	a	a	0	0	25	50	50	0	0	0	25				
	1.0	0	0	25	100	a	a	a	100	a	a	a	100	a	a	a	25	25	50	75	75	0	25	50	75				

<sup>a</sup>All earthworm died within 30 min. <sup>b</sup>All earthworm died within 1 hour