



BJP

Bangladesh Journal of Pharmacology

Research Article

GC-MS analysis, evaluation of phytochemicals, anti-oxidant, thrombolytic and anti-inflammatory activities of *Exacum bicolor*

GC-MS analysis, evaluation of phytochemicals, anti-oxidant, thrombolytic and anti-inflammatory activities of *Exacum bicolor*

Appaji Mahesh Ashwini¹, Latha Puttarudrappa², Belagumba Vijaykumar Ravi² and Mala Majumdar¹

¹Department of Biotechnology, Centre for Postgraduate Studies, Jain University, Bangalore, Karnataka 560 011, India;

²Departments of Biochemistry, Kempegowda Institute of Medical Sciences, Bangalore, Karnataka 560 070, India.

Article Info

Received: 9 June 2015

Accepted: 26 June 2015

Available Online: 22 September 2015

DOI: 10.3329/bjp.v10i4.23610

Cite this article:

Ashwini AM, Puttarudrappa L, Ravi BV, Majumdar M. GC-MS analysis, evaluation of phytochemicals, anti-oxidant, thrombolytic and anti-inflammatory activities of *Exacum bicolor*. Bangladesh J Pharmacol. 2015; 10: 745-52.

Abstract

The aim of the present study was to investigate the GC-MS analysis, phytochemical screening, anti-oxidant, thrombolytic and anti-inflammatory activities of methanol extract of leaves of *Exacum bicolor*. FTIR analysis confirmed the presence of alcohol, phenols, alkanes, aromatic compounds, aldehyde and ethers. GC-MS analysis revealed the presence of eight phyto-constituents. The total phenol, flavonoid and alkaloid contents were 18.0 ± 0.2 mg/GAE/g, 13.1 ± 0.4 mg QE/g and 108.0 ± 1.2 mg AE/g respectively. The DPPH assay exhibited potent anti-oxidant abilities with IC_{50} $8.8 \mu\text{g/mL}$. Significant thrombolytic activity was demonstrated by clot lysis method ($45.1 \pm 0.8\%$). The methanol extract showed significant membrane stabilization on human red blood cell with IC_{50} value of $37.4 \mu\text{g/mL}$. There was a significant correlation ($R^2 > 0.98$) with total phenolic content versus anti-oxidant and anti-inflammatory activity. The above results confirmed that *E. bicolor* could be a promising anti-oxidant, thrombolytic and anti-inflammatory agent.

Introduction

Oxidative stress plays a pivotal role in the development of human diseases (Rajendran et al., 2014). Anti-oxidants that can scavenge or neutralise the reactive oxygen species are beneficial in reducing the oxidative stress (Bandyopadhyay et al., 1999). Plants rich in anti-oxidants have ability to protect against oxidative cell damage that can lead in the treatment of many human diseases including diabetes, cancer, Alzheimer's, cardiovascular diseases, chronic inflammation, thrombus formation and several degenerative diseases in humans (Danino et al., 2008; Deore et al., 2008; Dinstel et al., 2013). The free radical scavenging molecules such as flavonoids, phenols, tannins, alkaloids, amines, vitamins and other metabolites possess anti-inflammatory, thrombolytic, anti-carcinogenic, antibacterial and antiviral activities (Filomena et al., 2008). Inflammation is a key factor in all aspects of coronary disease including

the initiation and progression of atherosclerotic plaque, plaque rupture, and thrombosis (atherothrombosis) where the oxidative stress is known to play a significant role (Freedman, 2008). Oxidative stress and inflammation are intimately linked with both the evolution of cardiovascular disease and acute coronary syndromes (Pashkow, 2011). Due to short comings present in the synthetic drugs, research has been directed towards the development of herbal medicine which are considered safer due to their natural activity.

Exacum bicolor Roxb., a member of family *Gentianaceae*, is a herbaceous plant possessing anti-oxidant and anthelmintic activities (Ashwini and Majumdar, 2014; Ashwini and Majumdar, 2015). Ethanopharmacologically *E. bicolor* is used for curing human ailments like diabetes, malaria, skin disorders, fungal diseases and inflammation (Marles and Farnsworth, 1995; Reddi et al., 2005; Pullaiah, 2006; Khare, 2007). No scientific report is



available till date to validate these folkloric uses. *E. bicolor* is endemic to peninsular India and presently considered as an endangered species (Sreelatha et al., 2007; Brilliant et al., 2012). Chemically *E. bicolor* consists of procatechuic, apigenin, luteolin, vanillic, ρ -coumaric acids, secoiridoids and ρ -hydroxybenzoic (Das et al., 1985; Khare, 2007). Hence, the present study is focussed to evaluate the anti-inflammatory and thrombolytic activities for the first time.

Materials and Methods

Plant collection and extraction

Plant material was collected from Kumar Parvatha, Western Ghats, Karnataka and authenticated by Regional Research Institute Bangalore, India (Accession No.: 557). The leaves were dried at room temperature in the shade. Twenty grams of the dried powdered leaves sample was soaked in 150 mL of methanol and was shaken intermediately. After 7 days the solution was filtered and was evaporated to dryness.

Infrared spectral analysis

The methanol leaf extract of *E. bicolor* was subjected to IR spectrum which was determined using Fourier transform infrared spectrophotometer (FTIR-Perkin-Elmer). The extract was ground with KBr powder and then pressed into pellets for FTIR measurement in the frequency range of 4,000–400 cm^{-1} (Bunghez et al., 2011).

Gas chromatography-mass spectrometry (GC-MS) analysis

The methanol leaf extract was subjected for GC-MS analysis equipped with Thermo GC-Trace ultra Ver: 5.0, Thermo MS DSQ II and equipped with column DB - 5MS capillary standard non-polar (length 30 m x inner diameter 0.25 mm film thickness 0.25 μm) was used for analysis. Helium gas was used as the carrier gas at constant flow rate 1 mL/min and an injection volume of 1 μL . The oven injector temperature 70°C and raised to 260°C at 6°C/min. Overall runtime was 40.5 min.

Identification of components

Interpretation on mass spectrum GC-MS was conducted using the database of National Institute Standard and Technology (NIST) having more than 62,000 patterns. The spectrum of the unknown component was compared with relative retention time and mass spectra of the known components stored in the NIST library.

Preliminary phytochemical screening

The preliminary qualitative phytochemical study of the methanol leaves extract were screened for alkaloids (Meyer's test), flavonoids (Shinoda test), saponin, ste-

roids, terpenoids, glycosides (Salkowski's test), phenols, tannins, amino acids, proteins (ninhydrin test), carbohydrates (Fehling's test), coumarins, quinones, oxalates and phlobotannins and acids (Harborne, 1973).

Total phenols

The total phenolics content in methanol extract was determined by FC method with minor modifications (Singleton and Rossi 1965). To 1 mL of extract (25, 50, 75, 100 $\mu\text{g}/\text{mL}$) 0.5 mL FCR (diluted 1:10 v/v) was added and allowed to stand for 5 min. To the above solution 20% sodium carbonate (1 mL) was added and allowed to stand for half hour in dark. The sample was read against the blank at 765 nm using UV-spectrophotometer. Results were expressed as in mg/g of dry weight of gallic acid equivalent (GAE) which was used as standard. All experiments were carried out in triplicates and represented as mean \pm SE.

Estimation of flavonoids

Total flavonoid content was determined by aluminium chloride method using quercetin as a standard (Chang et al., 2002). 1 mL of plant extract (25, 50, 75, 100 $\mu\text{g}/\text{mL}$) and 0.5 mL of 5% sodium nitrite was added to the above solution 0.5 mL of 10% aluminium chloride was added. Incubated at room temperature for 6 min and 2 mL of 1 M sodium hydroxide was added to the reaction mixture. Volume was made up with distilled water and the absorbance was measured at 420 nm spectrophotometrically. Results were expressed as quercetin equivalents (mg QE/g dry weight). All the tests were performed in triplicates and represented as mean \pm SE.

Estimation of alkaloids

Alkaloids were estimated according to Shamsa et al. (2008). *E. bicolor* methanolic extract (100 mg) was dissolved in 2 N HCL. 1 mL of the filtrate was washed with chloroform. The pH of the solution was adjusted to 7 with 0.1 N NaOH. Atropine was used as standard to which BCG solution and phosphate buffer was added. The mixture was shaken well and using chloroform and made up to 10 mL in a volumetric flask. The absorbance was measured at 470 nm against the blank. The values were expressed as atropine equivalent (AE) mg/g dry weight. All experiments were carried out in triplicates and represented as mean \pm SE.

Anti-oxidant assay

DPPH(1,1-Diphenyl-2-picrylhydrazyl) radical scavenging activity

Free radical scavenging activity of *E. bicolor* methanol extract was determined according to Braca et al. (2003). The reaction mixture consist of extract (25, 50, 75, 100 $\mu\text{g}/\text{mL}$), 2 mL of 0.002% methanol solution of DPPH. This solution was incubated for 30 min in dark. The absorbance was read at 517 nm using spectrophotometer. Control was prepared by omitting the extracts,

ascorbic acid was used as standard and the percentage inhibition activity was calculated using the equation:

$$\% \text{Inhibition} = [(A_{\text{Control}} - A_{\text{Extract}}) / A_{\text{Control}}] \times 100$$

Where A_{Control} is the absorbance of the control and A_{Extract} the absorbance of the extract. All the tests were performed in triplicates represented as mean \pm SE.

ABTS⁺ [2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid)⁺] radical cation scavenging assay

ABTS radical cation decolorization assay was determined with minor modifications (Katalinic et al., 2006). ABTS⁺ was produced by reacting 2 mM ABTS with 17 mM potassium persulfate and allowed to stand in dark at room temperature for 12-16 hours before use. To 1 mL of extract (25, 50, 75, 100 $\mu\text{g}/\text{mL}$) 2 mL of ABTS reagent was added. The absorbance was read at 734 nm and percentage inhibition was calculated using formula. Ascorbic acid was used as standard.

$$\% \text{ scavenging activity} = [(A_0 - A_t) / A_0] \times 100.$$

Where A_0 is the absorbance of the control and A_t is the absorbance of the extract. All the tests were performed in triplicates and represented as mean \pm SE.

In vitro thrombolytic activity

The thrombolytic activity was evaluated by the previously developed method (Prasad et al., 2007). Streptokinase (SK) vial was procured commercially of the brand name myokinase manufactured by Biocon (India) Ltd., Bangalore of 15,00,000 I.U. 5 mL of distilled water was used to dissolve. 100 μL (30,000 I.U) from the stock was used for further studies. Blood sample was drawn from human volunteers ($n=50$) who are healthy and do not show any history of acquiring oral contraceptive or undergoing anticoagulant therapy.

100 mg of powdered extract was dissolved in 10 mL distilled water. Blood was collected and distributed in pre weighed sterile eppendorf tube (0.5 mL/tube) and incubated at 37°C for 45 min. After clot formation, the serum was completely removed without disturbing the clot and each tube with the clot was again weighed to determine the clot weight (clot weight = weight of clot with tube - weight of empty tube). To each pre-weighed tube containing the clot, 100 μL of *E. bicolor* methanol leaf extract (test extract), SK (positive control) and distilled water (negative control) were separately added to the tubes. Tubes were incubated at 37°C for 90 min. The fluid released after the incubation period was removed and tubes were weighed again to observe the difference in weight after clot disruption. Percentage of clot lysis was calculated by calculating the weight that is taken before and after clot.

In vitro anti-inflammatory activity

Anti-inflammatory activity was assessed by human red blood cell (HRBC) membrane stabilization method

(Gandhisan et al., 1991). The blood sample was collected from healthy human volunteer. Equal volume of alsever solution (2% dextrose, 0.8% sodium citrate, 0.5% citric acid and 0.42% NaCl) was mixed and centrifuged at 3,000 rpm. The packed cells were washed with isosaline for 3 times and 10% suspension was made with isosaline. Various concentration of extracts (25, 50, 75, 100 $\mu\text{g}/\text{mL}$), 1 mL of phosphate buffer, 2 mL hyposaline and 0.5 mL of HRBC suspension were added. It was incubated at 37°C for 30 min and later centrifuged at 3,000 rpm for 20 min. The haemoglobin content of the supernatant solution was estimated at 560 nm spectrophotometrically. Diclofenac (10, 25, 50, 75, 100 $\mu\text{g}/\text{mL}$) was used as reference standard and control was prepared by omitting the extracts. The percentage of hemolysis and protection of HRBC membrane was calculated as follows:

$$\% \text{Hemolysis} = (\text{OD of test sample} / \text{OD of control}) \times 100$$

$$\% \text{Protection} = 100 - \% \text{Hemolysis}$$

Statistical analysis

All the results are expressed as mean \pm SEM. IC₅₀, percentage of clot lysis analysis was done using Graph pad prism 6 using ANOVA with post hoc analysis by DMRT. p values ≤ 0.05 were considered to be significant. The Correlation coefficient (R^2) values of all the different concentration of extracts between total phenols with DPPH assay and anti-inflammatory activities were evaluated.

Results

The FTIR spectrum (Figure 1) analysis revealed the identity of the functional groups of the active components present in the plant based on the peaks and the values in the IR region. *E. bicolor* exhibited characteristic absorption bands at 3436, 2919, 1689, 1604, 1402, 1370, 1239, 1101 and 1080 cm^{-1} . These results of FTIR analysis confirmed the presence of alcohols, phenols, alkanes, amines, aromatic compound, aldehyde and ethers.

The GC-MS analysis of phytoconstituents in methanol extract of leaves of *E. bicolor* revealed the presence of eight major phytoconstituents (Figure 2; Table I). The identification of the compounds was confirmed based on the retention time, peak area and the molecular formula. The major phytoconstituents reported are 1-methyl 2-(3-oxocyclohexenyl)imidazole (5.8%), erythrocentaurin (1.0%), neophytadiene (4.0%), hexadecanoic acid (5.4%), 6-octadecenoic acid (12.0%), (+)-inophylum D (2.7%), 4,6,8(14)-cholestatriene (6.0%) and methyl 3,4-diphenylpyrrolo[2,1,5-cd]indolizine-1-carboxylate (5.5%).

The preliminary qualitative phytochemical screening of the methanol extract revealed the presence of alkaloids, flavonoids, saponin, steroids, terpenoids, glycosides,




Figure 1: FTIR spectrum of *E. bicolor* (leaf) methanol extract

phenols, tannins, amino acids, proteins, carbohydrates and acids. The quantitative analysis of *E. bicolor* leaf methanol extract was based on the total phenols, flavonoids and alkaloids. The total phenols, flavonoid and alkaloid content in methanol extract were found to be 18 ± 0.2 mg/ GAE/g, 13.1 ± 0.4 mg QE/g and 108 ± 1.2 mg AE/g respectively.

The results of DPPH assay showed that methanol extract of *E. bicolor* had a good inhibitory activity in dose-dependent manner. The percentage inhibition of methanol extract was 91.8% at 100 $\mu\text{g}/\text{mL}$ concentration with an IC_{50} value of 8.8 $\mu\text{g}/\text{mL}$. In terms of ABTS cation radical scavenging assay, the percentage inhibition of *E. bicolor* leaf extract was 76.9% at 100 $\mu\text{g}/\text{mL}$ concentration with an IC_{50} value of 43.7 $\mu\text{g}/\text{mL}$. The extract also exhibited dose dependent inhibitory activity. Ascorbic acid was used as standard for both DPPH and ABTS assay with the IC_{50} value of 6.0 $\mu\text{g}/\text{mL}$ and 11.3 $\mu\text{g}/\text{mL}$ respectively.

In the present study, thrombolytic activity was evaluated by clot disruption method. Streptokinase, positive control, showed $54.8 \pm 0.7\%$ clot lysis. Clots when treated with sterile distilled water (negative control) showed only $8.3 \pm 0.5\%$ of clot lysis. The mean difference in clot lysis percentage between positive and negative control was significant ($p < 0.0001$). The tested methanolic extract of *E. bicolor* exhibited $45.1 \pm 0.8\%$ of clot lysis.

In vitro anti-inflammatory activity of *E. bicolor* extracts at different concentrations showed significant stabiliza-




Figure 2: GC-MS analysis of *Exacum bicolor* methanol leaf extract

Table I

Chemical profile identified by GC-MS analysis of methanol leaf extract of *E. bicolor*

SL. No.	RT	Compounds	Molecular formula	MW	%Area
1	16.51	1-Methyl 2-(3-oxocyclohexenyl)imidazole	$\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}$	176	5.8
2	17.72	Erythrocentaurin	$\text{C}_{10}\text{H}_8\text{O}_3$	176	1.0
3	20.26	Neophytadiene	$\text{C}_{20}\text{H}_{38}$	278	4.0
4	22.01	Hexadecanoic acid	$\text{C}_{17}\text{H}_{34}\text{O}_2$	270	5.4
5	25.33	6-Octadecenoic acid	$\text{C}_{19}\text{H}_{36}\text{O}_2$	296	12.0
6	28.78	(+)-Inophylum D	$\text{C}_{25}\text{H}_{24}\text{O}_5$	404	2.7
7	33.13	4,6,8(14)-Cholestatriene	$\text{C}_{27}\text{H}_{42}$	366	6.0
8	34.35	Methyl 3,4-Diphenylpyrrolo[2,1,5-cd] indolizine-1-carboxylate	$\text{C}_{24}\text{H}_{17}\text{NO}_2$	351	5.5




Figure 3: *In vitro* anti-inflammatory activity by HRBC membrane stabilization method of standard diclofenac and *E. bicolor*

tion towards HRBC membranes. The percentage protection (Figure 3) at concentration 100 µg/mL was more when compared to the other concentrations in *E. bicolor* ($84.6 \pm 2.6\%$) and diclofenac ($92.3 \pm 4.4\%$) as standard. The IC_{50} values of *E. bicolor* and diclofenac were 37.4 and 28.0 µg/mL respectively.

A positive, correlation of $R^2=0.9867$ was found between the total phenols and DPPH assay which was highly significant ($p=0.007$). Whereas the total phenols were also significantly ($p=0.006$) correlated with anti-inflammatory activity ($R^2= 0.9916$).

Discussion

FT-IR showed typical bands arising as a result of strong O-H stretching ($3550-3200\text{ cm}^{-1}$) intramolecular bonds. Other bands, occurred at $3000-2840\text{ cm}^{-1}$ (C-H stretching), $1710-1685\text{ cm}^{-1}$, showed strong (C=O stretching) conjugated aldehyde peaks at $1650-1600\text{ cm}^{-1}$ resulted in medium (C=C stretching) conjugated alkene, signals at $1150-1085\text{ cm}^{-1}$ gave strong (C-O stretching) aliphatic ether, $1450-1375\text{ cm}^{-1}$ resulted medium C-H bending alkane with methyl group, at $1085-1050\text{ cm}^{-1}$ strong (C-O stretching) primary alcohol. Based on the FT-IR spectrum, *E. bicolor* leaf extract contains various phyto-compounds with functional groups such as phenols, amines, aldehyde, alkanes, carboxylic acids and alcohols.

The GC-MS analysis of the plant extracts are becoming a valuable tool for detection of phytochemicals which can be aimed before the process of large scale purification (Vinay et al., 2014). In the present study, the GC-MS analysis of methanol extract of *E. bicolor* majorly contained phenols, alkanes, terpenoids, alkaloids which might be responsible for various medicinal activities (Sellamuthu et al., 2009). According to

the previously reported literature erythrocentaurin, a monoterpene alkaloid has also been identified from *Enicostemma hypsipifolium* and *Swertia lawii* (Ghosal et al., 1974). Neophytadiene (major component) and hexadecanoic acid were also present in *Centaurium erythraea* (Jovanovic et al., 2009) which belongs to family *Gentianaceae*. The present study was in accordance with the above literature.

Phytochemicals or secondary metabolites are chemical compounds which are formed during the plants normal metabolic processes and plants use them to protect themselves during stress related conditions (Ning et al., 2009). In the current study, the leaf extract of *E. bicolor* possesses $18 \pm 0.2\text{ mg/GAE/g}$ of phenols, $13.1 \pm 0.4\text{ mg QE/g}$ of flavonoids, $108 \pm 0.0\text{ mg AE/g}$ of alkaloids when determined spectrophotometrically. According to Baba and Malik (2014), *Gentiana kurroo* methanol leaf extract showed $34 \pm 1.8\text{ mg GAE/g}$ of phenols and $20 \pm 1.5\text{ mg RE/g}$ flavonoids respectively. The total phenols in methanol extract of *S. chirata* were about $38.4 \pm 0.4\text{ mg GAE/g}$ (Tupe et al., 2013).

Phenolic compounds are widely distributed in various plant species which have received considerable attention (Li et al., 2006). Phenolic anti-oxidants provided tremendous potential benefits because of their ability to scavenge reactive oxygen species (Bakirel et al., 2008). The total anti-oxidant power evaluates health beneficial effects because of the co-operative action of anti-oxidants. Therefore, it is desirable to measure the radical scavenging capacity level by more than one method (Fu et al., 2014).

DPPH anti-oxidant scavenging assay is based on the ability of DPPH, to decolorize in the presence of anti-oxidants. The visible deep purple color is produced due to the DPPH free radical which contains an odd electron. When DPPH accepts an electron donated by an anti-oxidant compound, the DPPH is decolorized which can be quantitatively measured by absorbance (Nuengchamnong et al., 2009). The methanolic extract of *E. bicolor* in the present study exhibited potent anti-oxidant activity at 100 µg/mL ($91.2 \pm 0.7\%$) with lower IC_{50} value (8.8 µg/mL). According to Vajjanathappa et al. (2008) the IC_{50} value of methanol extract of *Enicostemma axillare* was $325.5 \pm 5.9\text{ µg/mL}$. In case of *S. chirata*, IC_{50} value was 87.6 ± 0.4 (Tupe et al., 2013). According to Baba and Malik (2014), *Gentiana kurroo* showed 91% of inhibition at 600 µg/mL. In comparison to earlier findings, *E. bicolor* exhibited relatively strong radical scavenging activities and might serve as effective radical scavenger.

ABTS method has chromophores which are soluble in both aqueous and organic solvents, and may therefore serve the need to simultaneously measure hydrophilic and lipophilic anti-oxidants (Cekic et al., 2009). In this assay, ABTS radical cation was generated directly in

stable form using potassium persulfate and the anti-oxidant activity is measured in terms of decolorization (Sanchez-Moreno 2002). The free radical scavenging ability of *E. bicolor* was determined using ABTS radical cation which exhibited $76.9 \pm 0.9\%$ inhibition and the IC_{50} value was $43.7 \mu\text{g/mL}$. Tupe et al., (2013) reported that in *S. chirata* the IC_{50} value was $71.7 \pm 1.7 \mu\text{g/mL}$. But the present study with *E. bicolor* showed better anti-oxidant capacities when compared to *S. chirata*.

Atherothrombotic diseases occur as serious impacts of the thrombus formed in the blood vessels. Thrombolytics dissolve blood clots and are used in treating myocardial infarction, cerebral stroke, high cholesterol hypertension, hypotension, arrhythmias, congestive heart failure, coronary artery disease, unwanted blood clots and arteriosclerosis (American Pharmacist Association, 2010). The synthetic drugs have limitations and can lead to serious and sometimes fatal consequences. Studies have been conducted to find plants having anti-thrombotic compound which might lead to lower or no adverse effects (Prasad et al., 2007). In case of *E. bicolor*, the methanol leaf extract was assessed by clot disruption method which revealed $45.1 \pm 0.8\%$ of clot lysis while the streptokinase (positive control) and water (negative control) demonstrated $56.2 \pm 0.6\%$ and $8.3 \pm 0.5\%$ lysis of clot respectively. The results were also statistically significant ($p < 0.0001$). According to the previous study, *S. chirata* showed clot lysis of 46.1% and 31.9% in ethanolic and chloroform extract respectively (Hossain et al., 2012). According to Dhande et al., (2014) the crude extracts of *S. chirata* were found to have significant ($p < 0.01$) thrombolytic activity at 1 mg/mL . Whereas, the *E. bicolor* methanolic leaf extract showed better thrombolytic activity with lower concentration ($100 \mu\text{g/mL}$) when compared to the above study.

Plants with anti-oxidants properties are used for minimizing the inflammation related diseases (Wong et al., 2006). Erythrocyte membrane is an analogue of lysosomal membrane. The plant extract plays an important role in stabilizing lysosomal membrane which limits the inflammatory response by preventing the release of activated neutrophils, which might further cause tissue damage and inflammation (Murugasan et al., 1981). The extract may inhibit this process and acts as an anti-inflammatory compound. Investigation on *in vitro* anti-inflammatory activity by human red blood cell membrane stabilization method in *E. bicolor* exhibited $87.2 \pm 2.6\%$ of protection at $100 \mu\text{g/mL}$ with IC_{50} value of $37.4 \mu\text{g/mL}$, whereas Leelaprakash and Dass (2011) reported $30 \pm 0.0\%$ inhibition of hemolysis in *Encostemma axillare* methanol extract of which the present study exhibited potent anti-inflammatory activity.

In order to understand the relationship between the anti-oxidant activity with total phenolic content, the correlation between the DPPH scavenging activity and

the total phenolics were measured. Our results revealed that there is a strong and significant correlation ($R^2=0.9867$) between total phenolic content and DPPH free radical scavenging activity. There was also a positive correlation between total phenol contents and anti-inflammatory activity ($R^2 = 0.9916$).

Conclusion

Methanolic leaf extract of *E. bicolor* possesses phytochemicals which might be playing a major role in anti-oxidant, anti-inflammatory and thrombolytic activities. The positive correlation between total phenolic content with DPPH scavenging capacities and anti-inflammatory activity showed that total phenols might be the major contributor for these biological activities. *E. bicolor* extracts may be exploited as a source of beneficial compounds for oxidative stress related diseases in humans.

Ethical Issue

The collection of blood sample from the human volunteers was done with the approval by Institutional Ethics Committee of Kempegowda Institute of Medical Sciences (KIMS), Bangalore, Karnataka with registration No. ECR/216/Inst/Kar/2013. The volunteer donors were supplied with a consent form which informed detail of investigations.

Acknowledgement

The authors are thankful to management of Jain University for providing all the infrastructural facilities and financial support to carry out the research work.

References

- American Pharmacist Association. The pharmacy technician, 4th ed. Englewood, Morton Publishing Company, 2010, p 434.
- Ashwini AM, Majumdar M. Qualitative phytochemical screening and *in vitro* anthelmintic activity of *Exacum bicolor* Roxb., an endemic medicinal plant from Western Ghats in India. Acta Biol Ind. 2014; 3: 510-14.
- Ashwini AM, Majumdar M. Quantification of phytochemical contents and *in vitro* anti-oxidant activity of *Exacum bicolor* (Roxb.), an endemic medicinal plant. Int J Pharm Pharma Sci. 2015; 7: 225-30.
- Baba SA, Malik SA. Evaluation of anti-oxidant and antibacterial activity of methanolic extracts of *Gentiana kurroo* Royle. Saudi J Biol Sci. 2014; 21: 493-98.
- Bakirel T, Bakirel U, Keleş OU, Ülgen SG, Yardibi H. *In vivo* assessment of antidiabetic and anti-oxidant activities of rosemary (*Rosmarinus officinalis*) in alloxan-diabetic rabbits. J

- Ethnopharmacol. 2008; 116: 64-73.
- Bandyopadhyay U, Das D, Banerjee RK. Reactive oxygen specie: Oxidative damage and pathogenesis. *Curr Sci*. 1999; 77: 658-66.
- Braca A, Politi M, Sanogo R, Sanou H, Morelli I, Pizza C, Tommasi ND. Chemical composition and anti-oxidant activity of phenolic compounds from wild and cultivated *Sclero caryabirrea* (*Anacardiaceae*) leaves. *J Agric Food Chem*. 2003; 51: 6689-95.
- Brilliant R, Vincy MV, Joby P, Pradeepkumar. AP Vegetation analysis of Montane forest of Western Ghats with special emphasis on RET species. *IJBC* 2012; 4: 652-64.
- Bunghaz IR, Raduly M, Doncea S, Aksahin I, Ion RM. Lycopene determination in tomatoes by different spectral techniques (UV-VIS, FTIR, and HPLC). *Dig J Nanomater Bioest*. 2011; 6: 1349-56.
- Cekic SDC, Baskan KN, Tüttem E, Apak R. Modified cupric reducing anti-oxidant capacity (CUPRAC) assay for measuring the anti-oxidant capacities of thiol-containing proteins in admixture with polyphenols. *Talanta* 2009; 79: 344-51.
- Chang C, Yang M, Wen H, Chern J. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal*. 2002; 10: 178-82.
- Danino O, Gottlieb HE, Grossman S, Bergman M. Anti-oxidant activity of 1,3-dicaffeoylquinic acid isolated from *Inula viscosa*. *Food Res Int*. 2008; 42: 1273-80.
- Das S, Barua, RN, Sharma RP, Baruah, JN, Kulanthaivela P, Herza W. Secoiridoids from *Exacum tetragonum*. *Phytochemistry* 1985; 23: 908-09.
- Deore SL, Khadabadi SS, Daulatkar VD, Deokate UA, Farooqui IU. Evaluation of hypoglycemic and anti-diabetic activity of bark of *Butea monosperma*. *Pharmacog Mag*. 2008; 4: 134-38.
- Dhande SR, Aakruti AK, Kalpana AP, Vilasrao K. Antihyperlipidemic activity of *Bambusa bambos* (Druce.) and *Suertia chirata* (Buch-ham) in cholesterol suspension induced hypercholesterolemia in rats. *Int J Pharm Pharm Sci*. 2014; 6: 607-10.
- Dinstel RR, Cascio J, Koukel S. The anti-oxidant level of Alaska's wild berries: High, higher and highest. *Int J Circumpolar Health*. 2013; 72: 796-802.
- Filomena C, Sosa S, Mariangela M, Statti F, Uzunov G, Tubaro D, Menichini A, Loggia F, Della R. *In vivo* anti-inflammatory and *in vitro* anti-oxidant activities of Mediterranean dietary plants. *J Ethnopharma*. 2008; 116: 144-51.
- Freedman JE. Oxidative stress and platelets. *Arterioscl Throm Vas*. 2008; 28: 11-16.
- Fu R, Zhang YT, Guo YR, Liu FX, Chen F. Determination of phenolic contents and anti-oxidant activities of extracts of *Jatropha curcas* L. seed shell, a by-product, a new source of natural anti-oxidant. *Indian Crop Prod*. 2014; 58: 265-70.
- Gandhisan R, Thamarachelvan A, Baburaj. Anti-inflammatory action of *Lannea coromandelica* HRBC membrane stabilization. *Fitotherapy* 1991; 62: 82-83.
- Ghosal S, Singh AK, Sharma PV, Chaudhuri RK. Chemical constituents of *Gentianaceae* IX: Natural occurrence of erythrocentaurin in *Enicostemma hissiopifolium* and *Suertia lawii*. *J Pharm Sci*. 1974; 63: 944-45.
- Harborne JB. *Phytochemical methods*. London, Chapman and Hall Ltd, 1973.
- Hossain SM, Sharmin IF, Sheikh T, Hasan H, Rashedul AM. *In vitro* anti-oxidant, membrane stabilizing and thrombolytic activities of *Glycosmis arborea*. *J Pharm Biol Sci*. 2012; 15: 141-43.
- Jovanović O, Radulović N, Stojanović G, Palić R, Zlatković B, Gudžić B. Chemical composition of the essential oil of *Centaurium erythraea* Rafn (*Gentianaceae*) from Serbia. *J Essent Oil Res*. 2009; 21: 317-22.
- Katalinic V, Milos M, Kulisic T, JukicM. Screening of 70 medicinal plant extracts for anti-oxidant capacity and total phenols. *Food Chem*. 2006; 94: 550-57.
- Khare CC. *Indian medicinal plants: An illustrated Dictionary*. Berlin, Springer Verlag, 2007, pp 699-700.
- Leelaprakash G, Dass SM. *In vitro* anti-inflammatory activity of methanol extract of *Enicostemma axillare*. *Int J Drug Dev Res*. 2011; 3: 189-96.
- Li BB, Smith AB, HossainMM. Extraction of phenolics from citrus peels: II. Enzyme-assisted extraction method. *Sep Purif Technol*. 2006; 48: 189-96.
- Marles RJ, Farnsworth NR. Antidiabetic plants and their active constituents. *Phytomedicine* 1995; 2: 137-89.
- Murugasan N, Vember S, Damodharan C. Studies on erythrocyte membrane IV. *In vitro* haemolytic activity of oleander extract. *Toxicol Lett*. 1981; 8: 33-38.
- Ning G, Tianhua L, Xin Y, He P. Constituents in *Desmodium blandum* and their antitumor activity. *Trad Her Drug*. 2009; 40: 852-56.
- Nuengchamnon N, Krittasilp K, Ingkaninan K. Rapid screening and identification of anti-oxidants in aqueous extracts of *Houttuynia cordata* using LC-ESI-MS coupled with DPPH assay. *Food Chem*. 2009; 117: 750-56.
- Pashkow FJ. Oxidative stress and inflammation in heart disease: Do anti-oxidants have a role in treatment and/or prevention? *Int J Inflam*. 2011; 514623.
- Prasad S, Kashyap RS, Deopujari JY, Purohit HJ, Taori GM, Dagainawala HF. Effect of *Fagonia arabica* (Dhamasa) on *in vitro* thrombolysis. *BMC Complement Altern Med*. 2007; 7: 36.
- Pullaiah T. *Encyclopaedia of World medicinal plant*. Vol. II. New Delhi, Regency Pub, 2006.
- Rajendran P, Nandakumar N, Rengarajan T, Palaniswami R, Gnanadhas N, Lakshminarsaiah U, Gopas J, Nishigaki I. Anti-oxidants and human diseases. *Clinica Chimica Acta*. 2014; 436: 332-47.
- Reddi STV, Naidu BVAR, Prasanthi S. In: *Herbal remedies for diseases*. Alikhan I, Khanum A (eds). Hyderabad, Ukay Publications, 2005, pp 13-67.
- Sanchez-Moreno C. Methods used to evaluate the free radical scavenging activity in foods and biological systems. *Food Sci Tech Int*. 2002; 8: 121-37.

- Sellamuthu PS, Muniappan BP, Perumal SM, Kandasamy M. Antihyperglycemic effect of mangiferin in streptozotocin induced diabetic rats. *J Health Sci.* 2009; 55: 206-14.
- Shamsa F, Monsef H, Ghamooshi R, Verdianrizi M. Spectrophotometric determination of total alkaloids in some Iranian medicinal plants. *Thai J Pharm Sci.* 2008; 32: 17-20.
- Singleton VL, Rossi JA. Colorimetry of phenols with phosphomolybdic-phosphotungstic and reagents. *Am J Enol Viticult.* 1965; 37: 144-48.
- Sreelatha U, Baburaj TS, Kutty NC, Nazeem P, Bhaskar J. Cultivation prospects of *Exacum bicolor* Royle: An endangered, ornamental & anti-diabetic herb. *Nat Prod Rad.* 2007; 6: 402-04.
- Tupe RS, Kemse NG, Khaire AA. Evaluation of anti-oxidant potentials and total phenolic contents of selected Indian herbs powder extracts. *Int Food Res J.* 2013; 20: 1053-63.
- Vaijanathappa J, Baami S, Bhojraj S. *In vitro* anti-oxidant activity of *Enicostemma axillare*. *J Health Sci.* 2008; 54: 524-28.
- Vinay V, Karimulla S, Saravanan D. GC-MS analysis, preliminary phytochemical screening, physicochemical analysis and anti-diabetic activity of ethanol extract of *Jasminum cuspidatum* leaves. *Bangladesh J Pharmacol.* 2014; 9: 610-16.
- Wong CC, Li HB, Cheng KW, Chen F. A systematic survey of anti-oxidant activity of 30 Chinese medicinal plants using the ferric reducing anti-oxidant power assay. *Food Chem.* 2006; 97: 705-11.
-

Author Info

Mala Majumdar (Principal contact)
e-mail: malamajumdar51@gmail.com