



**BJP**

**Bangladesh Journal of Pharmacology**

**Research Article**

**Phytochemical analysis, cytotoxic, anti-oxidant and anti-diabetic activities of the aerial parts of *Sorghum halepense***

## Phytochemical analysis, cytotoxic, antioxidant and anti-diabetic activities of the aerial parts of *Sorghum halepense*

Muhammad Abdur Rehman Shah, Rahmat Ali Khan and Mushtaq Ahmed

Department of Biotechnology, University of Science and Technology, Bannu 28100, KPK, Pakistan.

### Article Info

Received: 18 February 2019  
Accepted: 2 August 2019  
Available Online: 21 August 2019  
DOI: 10.3329/bjp.v14i3.40292

#### Cite this article:

Shah MAR, Khan RA, Ahmed M. Phytochemical analysis, cytotoxic, antioxidant and anti-diabetic activities of aerial parts of *Sorghum halepense*. Bangladesh J Pharmacol. 2019; 14: 144-51.

### Abstract

Phytochemical screening, cytotoxic activity, total phenolic content, antioxidant and anti-diabetic activities of the *Sorghum halepense* methanolic extract and its different fractions were evaluated. Methanolic extract and its various fractions revealed the presence of reducing sugars, tannins, steroids, glycosides, flavonoids in the methanolic extract where they were absent in n-hexane fraction except flavonoids. Gums and saponins were absent in all the samples. Methanolic extract indicated the highest cytotoxic ( $80.7 \pm 1.3\%$ ) and anti-diabetic ( $62.5\%$ ) activities. The maximum total phenolic contents ( $28.7 \pm 1.4$  mg/mL) were found in the chloroform fraction. An aqueous fraction expressed the highest antioxidant activity  $74.1$  and  $97.1\%$  free radical scavenging properties in DPPH and ABTS assays respectively whereas, in the case of  $H_2O_2$ , methanolic extract indicated maximum ( $36.9\%$ ) activity. In conclusion, the extract of aerial parts of *S. halepense* is a source of compounds against cancer, diabetic and free radical associated disorders.

## Introduction

The damage caused by free radicals is the crucial ethological factor concerned with several chronic diseases such as cancer, atherosclerosis, diabetes mellitus, arthritis, and neurodegenerative diseases and also in the aging process. The epidemiological studies have revealed that the use of diets rich in fruits and vegetables and those containing selected natural antioxidants such as plant polyphenols, vitamin C and flavonoids are linked to reduced prevalence of cardiovascular and some cancerous chronic diseases (Gheldof and Engeseth, 2002; Liu et al., 2000; Siddhuraju and Becker, 2003; Zuo et al., 2002).

The anti-diabetic activities of *Achillea santolina* (Grover et al., 2002), *Adiantum capillus-veneris* (Al-Snafi, 2013), *Agrimony eupatoria* (Al-Snafi, 2015), *Aristolochia longa* (El Omari et al., 2019), *Atractylis gummifera* (Bouabid et al., 2019), *Eruca sativa* (Kishore et al., 2017), *Gaultheria trichophylla* (Alam and Saqib, 2017), *Leucas aspera* (Annapandian and Sundaram, 2017), *Myrica gale*,

*Rhodiola rosea*, *Rumex acetosa*, *Taraxacum officinale* (Sekhon-Loodu and Rupasinghe, 2019), *Trixis angustifolia* (Salazar-Gómez et al., 2019) and *Urtica dioica* (Ahangarpour et al., 2012) have been reported.

The antidiabetic effect of *Sorghum bicolor* (Chung et al., 2011) has been reported. However, the antidiabetic effect of *S. halepense* is not known.

The aim of this study was to investigate the phytochemicals, cytotoxic properties, total phenolic content, antioxidant and anti-diabetic properties of crude methanolic extract of aerial parts of *S. halepense* and its n-hexane, chloroform and aqueous fractions through brine shrimp lethality, DPPH, ABTS and  $H_2O_2$  assays and quantification of total phenolic content.

## Materials and Methods

### Chemicals for biological activities

Chemical reagents sodium phosphate monobasic, sodi-



um phosphate dibasic, DPPH (2, 2-diphenyl-1-picrylhydrazyl), ABTS•+ (2,2-azinobis (3-ethyl-benzothiazoline)-6-sulfonic acid disodium salt), hydrogen peroxide, amylase, DNS (dinitrosalicylic) acid, sea salts, gluco-phage, potassium persulfate were of analytical grade. Methanolic extract and its *n*-hexane, chloroform and aqueous fractions of the aerial part of *S. halepense*. Biotechnology Medical Services (BMS, UV-1602) spectrophotometer was used for measuring absorbance.

#### Plant material

The aerial parts of *S. halepense* were collected in March 2017 from the District Bannu, Pakistan. It was identified by Prof. Abdur Rehman, Govt. Postgraduate College, Bannu, Khyber Pakhtoon Khwa, Pakistan.

#### Preparation of crude extract

The aerial parts of young *S. halepense* were collected, washed with tap water, dried under shade and ground into fine powder with the help of pestle and mortar. The powder (600 g) was extracted in 1.5 L 70% methanol by keeping the powder in mentioned solvents at room temperature with frequent agitation for 72 hours and the resultant liquid was filtered (Whatman No. 3 filter paper, Whatman Ltd., England). The filtrate was sited at room temperature to vanish the solvent content. The resultant gummy methanolic extract (26.8 g) was collected and stored for future use.

#### Preparation of fractions

Methanolic crude extract (20 g) of *S. halepense* was sequentially extracted each with 300 mL *n*-hexane, chloroform and water using separating funnel to avoid any sort of damages to the filtrate. The respective solvents were evaporated completely at room temperature. The resulting extracts of *n*-hexane (2 g), chloroform (3.2 g) and water (7.9 g) were stored for further designed assays.

#### Phytochemicals screening

The methanolic extract of *S. halepense* and its different fractions were subjected to phytochemicals screening by using standard methods to investigate the presence of cardiac glycosides, saponins, flavonoids, tannins, steroids, reducing sugar and gums (Rice-Evans et al., 1996; Trease and Evans, 1989).

#### Cytotoxic assay

Methanolic extract and its various fractions of *S. halepense* were subjected to brine shrimp lethality bioassay to find out their cytotoxic activities (Meyer et al., 1982). A small amount of brine shrimp eggs (1 mg) were put in 4% artificial seawater contained in a cabin covered by aluminum foil and placed for 24 hours. After hatching, the shrimps crossed the porous central wall of the cabins and came into an illuminated one. The working solution of methanolic extract and its

different fractions were prepared in methanol with different concentrations i.e. 100, 250, 500 and 1000 µg/mL. Test tubes containing test samples were placed at room temperature for complete evaporation of methanol followed by inoculation with 10 shrimps to each test tube and incubation for 24 hours. Then counted live shrimps in each test tube, calculated %lethality with the help of Abbot's formula and compared the results (lethality) of control with experimental.

$$\% \text{Death} = [(\text{Sample} - \text{control}) / \text{control}] \times 100$$

#### Total phenolic content

The total phenolic content present in the methanolic extract and its fractions of *S. halepense* was determined by the method described elsewhere (Singleton and Rossi, 1965) with slight modification. The sample solution (250 µL; 1-5 mg/mL) was added to 2.5 mL Folin-Ciocalteu reagent (10 x diluted in distilled water) and incubated at room temperature for 5 min. A saturated solution of Na<sub>2</sub>CO<sub>3</sub> (2.5 mL, 60 mg/mL) was mixed to the reaction mixture and placed again for 2 hours at the mentioned temperature. Gallic acid was used as the standard. Measured the absorbance at 725 nm spectrophotometrically and articulated the results as gallic acid equivalent. Gallic acid solutions (1-5 mg/mL) were used to prepare the calibration curve and expressed the results as gallic acid equivalents (GAE).

#### Antioxidant assays

##### ABTS radical cation assay

The capability of methanolic extract and its various fractions of *S. halepense* to scavenge ABTS•+ (2, 2-azinobis (3-ethyl-benzothiazoline)-6-sulfonic acid disodium salt) free radicals was determined by making slight changes in (Re et al., 1999) protocol. Prepared ABTS (7 mM) and potassium persulfate solutions (2.45 mM), mixed together and placed for eight hours in dark. The mentioned stock solution was diluted with the relevant solvent (50%) and adjusted its absorbance of about 0.900 (± 0.02) at 745 nm at 30°C. 300 µL (125-2000 µg/mL in a respective solvent) of each sample was mixed with ABTS working solution and measured the absorbance. The percentage scavenging property of samples applied and the ascorbic acid was calculated using the following formula:

$$\text{Scavenging effect (\%)} = [(\text{control absorbance (ABTS)} - \text{sample absorbance}) / (\text{control absorbance})] \times 100$$

##### Hydrogen peroxide scavenging (H<sub>2</sub>O<sub>2</sub>) assay

The hydrogen peroxide scavenging activities of *S. halepense* methanolic extract and its different fractions were estimated according to the protocol mentioned elsewhere (Wettasinghe and Shahidi, 2000) protocol. The prepared sample solutions (125-2000 µg/mL), phosphate buffer (100 mM, pH 7.4) and hydrogen peroxide solution (43 mM) in the phosphate buffer. The



coefficient was determined in further analyses of results between total phenolic contents and different antioxidant and anti-diabetic activities. The  $p < 0.05$  was considered to be statistically significant.

## Results

### Phytochemical screening

The phytochemical screening of *S. halepense* methanolic extract and its various fractions revealed the presence of reducing sugars, tannins, steroids, glycosides, flavonoids in the methanolic extract whereas they were absent in the *n*-hexane fraction except flavonoids (Table I). Gums and saponins were absent in all the samples.

### Cytotoxic assay

The cytotoxic activities using brine shrimps lethality assay showed  $80.7 \pm 1.3$ ,  $60.3 \pm 1.1$ ,  $50.4 \pm 1.6$  and  $40.6 \pm 1.5$  dead brine shrimp in crude methanolic extract and its chloroform, aqueous and *n*-hexane fractions respectively at the concentration of 1,000  $\mu\text{g/mL}$  (Table II).

### Total phenolic contents

The maximum amount of phenolic contents was observed in chloroform fraction ( $28.7 \pm 1.4$  mg GAE/g) followed by methanol ( $22.5 \pm 1.3$  mg GAE/g), aqueous ( $15.1 \pm 1.3$  mg GAE/g) and *n*-hexane fraction ( $5.6 \pm 1.4$  mg GAE/g).

### Antioxidant assays

#### DPPH (1,1-diphenyl-2-picrylhydrazyl) method

The aqueous fraction indicated highest antioxidant activity (74.1%) followed by crude methanolic extract (73.5%) and hexane fraction (32.9%) whereas chloroform revealed lowest activity (24.6%) at the concentrations of 2 mg/mL. Ascorbic acid expressed 86.4% antioxidant activity at the same concentration. The antioxidant characteristics of *S. halepense* methanolic extract and its different fractions were concentration-dependent.

#### ABTS-radical cation assay

The highest antioxidant activity was revealed by an aqueous fraction (97.1%) followed by crude methanolic extract (96.7%) and chloroform fraction (79.0%) while the said activity was not observed in *n*-hexane fraction. The mentioned activity of aqueous fraction (97.1%) was comparable with ascorbic acid (96.2%) (Figure 1). An indirect correlation was observed between phenolic contents and antioxidant activities of the applied samples, the chloroform fraction exhibiting maximum phenolic contents ( $28.7 \pm 1.4$  mg/mL) expressed minimum antioxidant activity (79.0%), the methanolic extract had moderate phenolic contents ( $22.5 \pm 1.3$  mg/mL) had shown moderate antioxidant activity (96.7%).

#### Hydrogen peroxide scavenging ( $\text{H}_2\text{O}_2$ ) capacity

Table I					
Phytochemical screening of <i>S. halepense</i> methanolic extract and its chloroform and aqueous fractions					
Phytochemical	Tests	Crude methanolic extract and its fractions			
		Aqueous	Methanol	Chloroform	<i>n</i> -Hexane
Saponins	Alkaline reagent test	-	-	-	-
Flavonoids	Ferric chlorides test	+	+	-	+
Glycosides	Fehling test	+	+	+	-
steroids	Millon's test	+	+	+	-
Reducing sugar	Benedict's test	+	+	+	-
Tannins	Ferric chloride test	+	+	+	-
Gums	Molish's reagent test	-	-	-	-

Table II					
%Lethality of brine shrimps caused by crude methanolic extract and its various fractions of <i>S. halepense</i>					
Concentration ( $\mu\text{g/mL}$ )	%Lethality of crude methanolic extract and its fractions				
	Methanolic extract	Chloroform	Aqueous	<i>n</i> -Hexane	Control
100	$40.7 \pm 1.3$	$30.6 \pm 1.2$	$20.3 \pm 1.5$	$10.2 \pm 1.7$	$00 \pm 00$
250	$50.8 \pm 1.5$	$30.8 \pm 1.1$	$30.9 \pm 1.3$	$10.3 \pm 1.6$	$00 \pm 00$
500	$70.6 \pm 1.7$	$50.4 \pm 1.8$	$30.1 \pm 1.5$	$30.7 \pm 1.9$	$10.2 \pm 1.1$
1000	$80.7 \pm 1.3$	$60.3 \pm 1.1$	$50.4 \pm 1.6$	$40.6 \pm 1.5$	$00 \pm 00$




Figure 1: DPPH free radical scavenging capability (A), ABTS free radical scavenging capability (B), H<sub>2</sub>O<sub>2</sub> free radical scavenging capability (C) and anti-diabetic activity (D) of the *S. halepense* methanolic extract and its fractions;

Hf: n-Hexane fraction, Cf: chloroform fraction, Me: methanolic extract, Aqf: aqueous fraction and Asa: ascorbic acid; n-hexane fraction did not show anti-oxidant activity in B; hexane fraction and Cf: chloroform fraction did not show anti-oxidant activities in C; hexane fraction and Aqf: aqueous fraction did not express anti-diabetic activities in D

Table III		
Correlation between phenolics and <i>S. halepense</i> extract soluble fractions		
Assays	Correlation R <sup>2</sup> phenolics	Significance
%DPPH radical scavenging ability	0.5	ns
%ABTS scavenging ability	0.6	ns
%H <sub>2</sub> O <sub>2</sub> scavenging	0.9	ns
%Alpha-amylase inhibition	0.8	ns

The applied extract sample of *S. halepense* expressed scavenging of H<sub>2</sub>O<sub>2</sub> in a concentration-dependent manner. The crude methanolic extract and its water fraction indicated 36.9 and 36.5% scavenging activity respectively on hydrogen peroxide while ascorbic acid showed 51.0% at the concentration of 2000 µg/mL. The chloroform and *n*-hexane fraction did not express H<sub>2</sub>O<sub>2</sub> scavenging activities (Figure 1).

#### Anti-diabetic assay

Crude methanolic extract and its different fractions of *S. halepense* were subjected to an anti-diabetic assay to assess their anti-diabetic properties (Figure 1). The said characteristic of glucophage (commercially available medicine used as a standard) and crude methanolic extract were observed (48.9%) and (62.5%) respectively.

#### Correlation among total phenolic contents, anti-

#### oxidant and anti-diabetic activities

Chloroform fraction exhibited maximum phenolic content (28.7 ± 1.4 mg GAE), with minimum DPPH and ABTS free radical scavenging properties while did not show anti-diabetic and hydrogen peroxide radicals scavenging activities (Table III).

#### Discussion

The phytochemical assessment of aerial parts of young *S. halepense* extract and its diverse fractions showed the existence of reducing sugars, tannins, steroids, glycosides, flavonoids in the methanolic extract where they were absent in *n*-hexane fraction except flavonoids. Gums and saponins were absent in all samples. Aerial parts of young *S. halepense* methanolic extract and its chloroform, aqueous and *n*-hexane fractions showed brine shrimp lethality up to 80.7 ± 1.3%, 60.3 ± 1.1%, 50.4 ± 1.6% and 40.6 ± 1.5% respectively at a concentration of 1000 µg/mL. Different results were observed in different antioxidant assays i.e. aqueous fraction expressed highest antioxidant activity 74.1% and 97.1% free radical scavenging properties in DPPH and ABTS respectively whereas in case of H<sub>2</sub>O<sub>2</sub> methanolic extract indicated maximum (36.9%) activity. It is found that plant extracts with higher total phenolic contents have lower antioxidant properties. Aerial parts of young *S. halepense* methanolic extract and its different fractions were subjected to amylase inhibition assay to find out

their anti-diabetic properties. The anti-diabetic activities of glucophage (used as a standard) and methanolic extract of aerial parts of young *S. halepense* were recorded (48.9%) and (62.5%) respectively. The higher anti-diabetic behavior of methanolic extract than glucophage indicates that the mentioned extract is an effective source of anti-diabetic compounds. Dose-dependent percentage inhibition of alpha amylase was observed.

The major chemical constituents of *S. halepense* are sorigoleone and dihydrosorigoleone (Baerson et al., 2008).

Comparable phytochemicals constituents were observed in various extracts of leaves of *T. chebula* (Kathirvel and Sujatha, 2012). The reported cytotoxic activities of *Calligonum polygonoides* (Khan et al., 2015) and *Solanum surrantance burm* (Shah et al., 2013). *Arcangelisia flava*, *Coscinium blumeianum* and *Fibraurea tinctoria* (Keawpradub et al., 2005) are harmonious with our results. Previously, unlike antioxidant results have reported in different antioxidant assays (Hagerman et al., 1998). Our results are congruent with earlier studies which reported indirect relations between phenolic content and antioxidant activities exhibited by plant extracts (Kahkonen et al., 1999; Ismail et al., 2004). Our results resemble with curry tree and petai that contained high total phenolic content but lower anti-oxidant activities (Wong et al., 2006). Many previous studies have reported anti-diabetic properties and antioxidant activities of phenolic compounds in different parts of medicinal plants (Krings and Berger, 2001; Samal, 2013). Resembling dose-dependent and higher anti-diabetic results were revealed by the extract of *P. guajava* leaves *in vitro* (Manikandan et al., 2016).

Phytochemical assessment of plant extract and its different fractions revealed the presence of various phyto constituents that demonstrated specific therapeutic effects. Further study of plant extract is dependent on the presence of these compounds. Phenolic compounds, like flavonoids and terpenoids are the major antioxidants that exercise scavenging of free radicals and hence neutralization of reactive oxygen species (Hossain et al., 2011). Flavonoids use several mechanisms such as scavenging of free radicals and the inhibitions of enzymes involved in the production of free radicals and thus prevent the development of oxidative stress (Benavente-García et al., 1997). Usually the phytochemicals are extracted in different solvents where each one has specific influence on the extraction of said constituents depending on the nature of solvent and its interface with desired compounds. Therefore, the extracted compounds and their dry weight will be variable extracted in various solvents (Li et al. 2006; Pereira et al., 2008; Zhang et al., 2009). The cytotoxic findings imply that the assessed extracts are probable potential sources of antimicrobial and anticancer compounds. The differences observed in scavenging

behavior of test sample and its fractions against various free radicals system may depict that different oxidation reduction mechanisms are involved in the different assays. The higher free radical scavenging activity of aerial parts of young *S. halepense* aqueous fraction and methanolic extract to forage more free radicals (ABTS+) might be owing to higher molecular weight phenolics (tannins) and their convenience depends on molecular weight, the number of aromatic rings and nature of hydroxyl group's exchange than the explicit functional groups. Various antioxidant compounds such as  $\beta$ -carotene, tocopherol, selenium, vitamin C or phenolic compounds perform in different ways with divers mechanisms (Ismail et al., 2004). Serious health problems like obesity and diabetic results from metabolic disorder of carbohydrates. Problems in insulin secretion or deterioration in its function decreases the breakdown of disaccharides and polysaccharides and eventually cause the onset of diabetes mellitus (Funke and Melzig, 2006). Inhibition of alpha amylase, an enzyme responsible for digestion of carbohydrate slows down the process of carbohydrate digestion and hence reduce the uptake of glucose by intestines (Cheng and Fantus, 2005). Searching of natural inhibitor of amylase can be conducted by *in vitro* amylase inhibition assay and hence ultimate estimation of anti-diabetic characteristics of plant extracts.

---

## Conclusion

The aerial parts of young *S. halepense* contain considerable cytotoxic, antioxidant and anti-diabetic properties. The chloroform fraction exhibited maximum phenolic contents but minimum antioxidant activities.

---

## Conflict of Interest

The author declares that no conflict of interests.

---

## Acknowledgement

We are thankful to all my friends and colleague who guided us.

---

## References

- Ahangarpour A, Mohammadian M, Dianat M. Antidiabetic effect of hydroalcoholic *Urtica dioica* leaf extract in male rats with fructose-induced insulin resistance. *Iranian J Med Sci.* 2012; 37: 181.
- Alam F, Saqib QN. Antidiabetic potential of *Gaultheria trichophylla* in mice. *Bangladesh J Pharmacol.* 2017; 12: 292-98.
- Al-Snafi A E. Chemical constituents and pharmacological activities of milfoil (*Achillea santolina*). A Review. *Int J Pharm*

- Tech Res. 2013; 5: 1373-77.
- Al-Snafi A E. The pharmacological and therapeutic importance of *Agrimonia eupatoria*: A review. *Asian J Pharm Sci Technol*. 2015; 5: 112-17.
- Annapandian V, Sundaram RS. *In vitro* antidiabetic activity of polar and nonpolar solvent extracts from *Leucas Aspera* (Willd.) link leaves. *Pharmacogn Res*. 2017; 9: 261-65.
- Baerson SR, Dayan FE, Rimando AM, Nanayakkara NP, Liu CJ, Schröder J, Fishbein M, Pan Z, Kagan IA, Pratt LH, Cordonnier-Pratt MM, Duke SO. A functional genomics investigation of allelochemical biosynthesis in *Sorghum bicolor* root hairs. *J Biol Chem*. 2008; 283: 3231-47.
- Benavente-García O, Castillo J, Marin FR, Ortuño A, Del Río JA. Uses and properties of citrus flavonoids. *J Agric Food Chem*. 1997; 45: 4505-15.
- Bouabid K, Lamchouri F, Toufik H, Boulfia M, Senhaji S, Faouzi ME. *In vivo* anti-diabetic effect of aqueous and methanolic macerated extracts of *Atractylis gummifera*. *Bangladesh J Pharmacol*. 2019; 14: 67-73.
- Cheng AY, Fantus IG. Oral antihyperglycemic therapy for type 2 diabetes mellitus. *Can Med Assoc J*. 2005; 172: 213-26.
- Chung IM, Kim EH, Yeo MA, Kim SJ, Seo MC, Moon HI. Antidiabetic effects of three Korean sorghum phenolic extracts in normal and streptozotocin-induced diabetic rats. *Food Res Int*. 2011; 44: 127-132.
- El Omari N, Sayah K, Fettach S, El Bliidi O, Bouyahya A, Faouzi MEA, Barkiyou M. Evaluation of *in vitro* antioxidant and antidiabetic activities of *Aristolochia longa* extracts. *Evid Based Complement Altern Med*. 2019; 2019.
- Funke I, Melzig MF. Traditionally used plants in diabetes therapy: Phytotherapeutics as inhibitors of alpha-amylase activity. *Rev Bras Farmacogn*. 2006; 16: 1-5.
- Gheldof N, Engeseth NJ. Anti-oxidant capacity of honeys from various floral sources based on the determination of oxygen radical absorbance capacity and inhibition of *in vitro* lipoprotein oxidation in human serum samples. *J Agric Food Chem*. 2002; 50: 3050-55.
- Grover J, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. *J Ethnopharmacol*. 2002; 81: 81-100.
- Gyamfi MA, Yonamine M, Aniya Y. Free-radical scavenging action of medicinal herbs from Ghana: *Thonningia sanguinea* on experimentally-induced liver injuries. *Gen Pharmacol*. 1999; 32: 661-67.
- Hagerman AE, Riedl KM, Jones GA, Sovik KN, Ritchard NT, Hartzfeld PW, Riechel TL. High molecular weight plant polyphenolics (tannins) as biological antioxidants. *J Agric Food Chem*. 1998; 46: 1887-92.
- Hossain MA, Shah MD, Gnanaraj C, Iqbal M. *In vitro* total phenolics, flavonoids contents and antioxidant activity of essential oil, various organic extracts from the leaves of tropical medicinal plant *Tetragium* from Sabah. *Asian Pac J Trop Biomed*. 2011; 4: 717-21.
- Ismail A, Marjan ZM, Foong CW. Total antioxidant activity and phenolic content in selected vegetables. *Food Chem*. 2004; 87: 581-86.
- Kahkonen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, Heinonen M. Antioxidant activity of plant extracts containing phenolic compounds. *J Agric Food Chem*. 1999; 47: 3954-62.
- Kathirvel A, Sujatha V. *In vitro* assessment of antioxidant and antibacterial properties of *Terminalia chebula* Retz. leaves. *Asian Pac J Trop Biomed*. 2012; 2: 788-95.
- Keawpradub N, Dej-adisai S, Yuenyongsawad S. Antioxidant and cytotoxic activities of Thai medicinal plants named Khaminkhruea: *Arcangelisia flava*, *Coscinium blumeianum* and *Fibraurea tinctoria* Songklanakarin. *J Sci Technol*. 2005; 27: 455-67.
- Khan A, Khan RA, Ahmed M, Mushtaq N. *In vitro* antioxidant, antifungal and cytotoxic activity of methanolic extract of *Calligonum polygonoides*. *Bangladesh J Pharmacol*. 2015; 10: 316-20.
- Kishore L, Kaur N, Kajal A, Singh R. Extraction, characterization and evaluation of *Eruca sativa* against streptozotocin-induced diabetic nephropathy in rat. *Bangladesh J Pharmacol*. 2017; 12: 216-27.
- Krings U, Berger R. Antioxidant activity of some roasted foods. *Food Chem*. 2001; 72: 223-29.
- Kwon YI, Apostolidis E, Shetty K. Evaluation of pepper (*Capsicum annuum*) for management of diabetes and hypertension. *J Food Biochem*. 2007; 31: 370-85.
- Li BB, Smith B, Hossain MM. Extraction of phenolics from citrus peels: II. Enzyme-assisted extraction method. *Sep Purif Technol*. 2006; 48: 189-96.
- Liu S, Manson JE, Lee IM, Cole SR, Hennekens CH, Willett WC, Buring JE. Fruit and vegetable intake and risk of cardiovascular disease: The Women's health study. *Am J Clin Nutr*. 2000; 72: 922-28.
- MacDonald-Wicks LK, Wood LG, Garg ML. Methodology for the determination of biological antioxidant capacity *in vitro*: A review. *J Sci Food Agric*. 2006; 86: 2046-56.
- Manikandan R, Anand AV, Kumar S. Phytochemical and *in vitro* anti-diabetic activity of *Psidium guajava* leaves. *Pharmacogn J*. 2016; 8: 392-94.
- Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL. Brine shrimp: A convenient general bioassay for active plant constituents. *Planta Med*. 1982; 45: 31-34.
- Moon JK, Shibamoto T. Antioxidant assays for plant and food components. *J Agric Food Chem*. 2009; 57: 1655-66.
- Pereira JA, Oliveira I, Sousa A, Ferreira ICFR, Bento A, Estevinho L. Bioactive properties and chemical composition of six walnut (*Juglans regia* L.) cultivars. *Food Chem Toxicol*. 2008; 46: 2103-11.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med*. 1999; 26: 1231-37.
- Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med*. 1996; 20: 933-56.



- Salazar-Gómez A, Vargas-Díaz ME, Garduño-Siciliano L. Hypoglycemic potential of *Trixis angustifolia* aqueous extract in alloxan-induced diabetic mice. Bangladesh J Pharmacol. 2019; 14: 74-79.
- Samal PK. Antioxidant activity of *Strobilanthes asperimus* in albino rats. Asian J Pharm Res. 2013; 3: 71-74.
- Sekhonor-Loodu S, Rupasinghe H. Evaluation of antioxidant, antidiabetic and antiobesity potential of selected traditional medicinal plants. Front Nutr. 2019; 6: 53.
- Shah MAR, Khan H, Khan S, Muhammad N, Ullah Khan F, Shahnaz, Muhammad A, Khan YM. Cytotoxic, anti-oxidant and phytotoxic effect of *Solanum surattense* Burm F fruit extracts. Int J Pharmacogn Phytochem. 2013; 28: 1154-58.
- Siddhuraju P, Becker K. Anti-oxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. J Agric Food Chem. 2003; 51: 2144-55.
- Singh HP, Batish DR, Kohli R. Allelopathy in agroeco-systems: An overview. J Crop Prod. 2001; 4: 1-41.
- Singleton V, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. Am J Enol Vitic. 1965; 16: 144-58.
- Trease GE, Evans WC. Phytochemical screening. Pharmacognsy. 11th ed. London, Macmillian Publishers, 1989, pp 45-50.
- Wettasinghe M, Shahidi F. Scavenging of reactive-oxygen species and DPPH free radicals by extracts of borage and evening primrose meals. Food Chem. 2000; 70: 17-26.
- Wong SP, Leong LP, Koh JHW. Anti-oxidant activities of aqueous extracts of selected plants. Food Chem. 2006; 99: 775-83.
- Zhang Z, Liao L, Moore J, Wu T, Wang Z. Anti-oxidant phenolic compounds from walnut kernels (*Juglans regia* L.). Food Chem. 2009; 113: 160-65.
- 

**Author Info**

Rahmat Ali Khan (Principal contact)

e-mail: rahmatgul\_81@yahoo.com