

## DETERMINATION OF AN EFFECTIVE MEDIA AND ITS HORMONE AND PROTEIN SUPPLEMENTATION FOR *IN VITRO* MATURATION OF OOCYTES OF INDIGENOUS ZEBU COWS

M. M. Rahman, S. M. N. Morshed, N. S. Juyean and M. M. U. Bhuiyan\*

Department of Surgery and Obstetrics, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

### ABSTRACT

*In vitro* maturation (IVM) of oocytes is the first important step for successful *in vitro* embryo production of any mammalian species. The objectives of the present study were to determine an effective basic medium and its hormone and protein supplementation for IVM of oocytes of indigenous zebu cows. The oocytes were derived from ovaries of locally slaughtered cows after aspiration of follicle. The oocytes were cultured in medium for 24 hrs at 38.5°C with 5% CO<sub>2</sub> in humidified air for maturation. The maturation of oocytes was evaluated by examining the presence of first polar body extrusion in denuded oocytes under inverted microscope. To determine an effective basic medium, the oocytes were cultured in fetal bovine serum (FBS) supplemented tissue culture medium (TCM), modified synthetic oviduct fluid (mSOF) and Tyrodes albumin lactate pyruvate (TALP) medium. The maturation rate was significantly higher (74±4.2) in TCM medium than that of TALP medium (58.2±6.2). To determine an effective hormone supplementation for maturation medium, the oocytes were cultured in either in follicle stimulating hormone (FSH) or gonadotrophin supplemented TCM. The maturation rate of oocytes was significantly ( $p>0.05$ ) higher (73.3±4.0) in FSH supplemented medium than that of gonadotrophin supplemented counterpart (60.2±6.6). To determine an effective protein supplementation, the oocytes were cultured in FBS, oestrus cow serum (OCS) and bovine serum albumin (BSA) supplemented TCM 199. The maturation rate of oocytes were 73.0±5.9, 71.1±2.8, and 62.5±9.4 in medium supplemented with FBS, OCS and BSA respectively ( $p>0.05$ ). In conclusions, TCM supplemented with either FBS, OCS or BSA as protein and FSH as hormone may be used as medium for IVM of oocytes of indigenous zebu cows.

**Keywords:** Hormone, IVM, Media, Oocytes, Protein, Zebu cows

### INTRODUCTION

Livestock plays a major role in the life of farmers in developing countries. It provides food, income, employment and many other contributions including production of hide, draft power, fertilizer and fuel. Bangladesh is highly density populated country in the world. There are mark deficit of milk and meat production in Bangladesh. The indigenous zebu (*Bos indicus*) cattle of Bangladesh are lower yielding than the exotic (*Bos taurus*) cattle although the indigenous cattle are highly adapted to tropical environment and resistant to maximum diseases. Therefore, it is essential to upgrade the locally adapted indigenous zebu cattle for increasing milk and meat production. The up-gradation of indigenous zebu cows by artificial insemination using semen of exotic breed has been in practice for more than 4 decades in Bangladesh (Ahmed and Islam, 1987). However, the progress of genetic up-gradation is slower than expectation. So it is essential for application of other assisted reproductive technologies (ARTs) for rapid production of crossbred cattle to meet up the requirement of milk and meat.

Rapid generation of F1 offspring can be done by applying ARTs such as multiple ovulation and embryo transfer (MOET) and *in vitro* embryo production (IVEP) followed by embryo transfer (ET) in zebu cows. The IVEP and ET is more suitable than MOET as IVEP can use slaughterhouse derived oocytes for generation of F1 crossbred embryos. By this time, the IVEP and ET techniques have already been established with satisfactory results in many animals and humans in most of the country in the world. However, this IVEP-ET technique has not yet been established in this country. *In vitro* maturation (IVM) of oocytes is the first and most important step for any IVEP-ET programme in cattle. However, although several preliminary studies have been conducted on IVM of oocytes of zebu cows in Bangladesh (Das *et al.*, 2006; Islam *et al.*, 2007; Talukder *et al.*, 2008; Morshed *et al.*, 2014; Singha *et al.*, 2015; Choudhury *et al.*, 2017) still the IVM rate needs to be improved. The maturation medium and supplementation of protein and hormones in it may play an important role for IVM rate and subsequent development after *in vitro* fertilization (IVF) (Bavister *et al.*, 1992). Therefore, the present study was conducted to select an efficient basic medium including supplementation of suitable hormone and protein in it for optimal rate IVM of oocytes of indigenous Zebu cows in Bangladesh.

---

\*Corresponding e-mail address: mmubhuiyan@gmail.com

## **MATERIALS AND METHODS**

The study was carried out at the Community-based Dairy Veterinary Foundation (CDVF) Laboratory, Department of Surgery and Obstetrics, Bangladesh Agricultural University, Mymensingh.

### **Chemicals and media**

All the chemicals, reagents and media constituents were purchased from Sigma-Aldrich Inc., St Louis, USA. Media and reagents were prepared using standard protocol and under aseptic condition. All media were filtered using 0.22 $\mu$ m pore size filter (Durapore<sup>®</sup> membrane filter, Carrigtwohill, Ireland) and culture medium was routinely equilibrated in incubator (VS-9000C, Vision Scientific Co. Ltd. South Korea) at 38.5°C with 5% CO<sub>2</sub> in humidified air for at least 2 hrs before use.

### **Collection of ovaries**

The ovaries of indigenous zebu cows were collected from local slaughter house after slaughtering within 2 hrs and carried to the laboratory in a thermo flask containing warm normal saline (37°C, 0.9% sodium chloride solution, w/v).

### **Collection of oocytes**

In the laboratory, the ovaries were rinsed three times with warm (37°C) normal saline. The follicular fluid of 2 to 8 mm diameter follicles was aspirated using an 18 gauge needle (TERUMO<sup>®</sup>, Beijing, China) fitted with a 10 ml disposable plastic syringe (JMI Syringes and Medical Devices Ltd <sup>®</sup>, Chauddagam, Comilla, Bangladesh)

### **Selection of oocytes for culture**

The aspirated follicular fluid was transferred in a 60 mm petridish (Greiner bio-one, Frickenhausen, Germany) and left for 5 minutes for sedimentation. The retrieved follicular aspirate was diluted with HEPES-buffered tissue culture medium (TCM) 199 supplemented with bovine serum albumin (BSA). Oocytes were selected under a stereo-microscope (LABOMED, USA). The cumulus-oocyte-complexes (COCs) with more than 3 compact cumulus cell layers and homogenous ooplasm were selected for maturation. The COCS were washed three times in washing medium followed by once washing in maturation medium.

### **Culture of oocytes for maturation**

Four 50 $\mu$ l drops of maturation medium were prepared in 35mm petridish (FALCON, Becton Dickinson Labware, USA) and covered with embryo tested mineral oil. For *IVM*, 8-10 COCs were cultured in each drop of medium in incubator at 38.5°C with 5% CO<sub>2</sub> in humidified air for 24 hrs.

### **Evaluation of oocytes for maturation**

The culture drops were examined under the stereo microscope for expansion of cumulus cell after 24 hrs of culture in the maturation media. Presumptive maturation was confirmed by the degree of cumulus expansion. To examine the presence of first polar body extrusion, the COCs were denuded by using denuding agent (3% sodium citrate, w/v in HEPES buffered TCM 199 medium) and pipetting. After pipetting, the denuded oocytes were kept in 10 $\mu$ l drops of HEPES buffered TCM 199 and examined for presence of polar body under inverted microscope (Leica DM IRE2, Germany) with the help of a mouth controlled pipette.

## **Experimental design**

### **Experiment 1**

To determine an effective basal maturation medium, the COCs were cultured in TCM 199 (TCM, Earle's salts with sodium bicarbonate), modified synthetic oviductal fluid (mSOF) medium and Tyrodes albumin lactate pyrovate (TALP) medium supplemented with 10% fetal bovine serum (FBS), 5 $\mu$ g/ml follicle stimulating hormone (FSH) and 1 $\mu$ g/ml oestradiol (OE<sub>2</sub>). Each experiment was repeated for 10 times across the days.

### **Experiment 2**

To determine an effective hormone supplementation in maturation medium, the oocytes were cultured in TCM 199 supplemented with 5 $\mu$ g/ml FSH or 10 IU/ml Gonadotrophin, 10% FBS and 1 $\mu$ g/ml OE<sub>2</sub>. Each experiment was repeated for 8 times across the days.

### Experiment 3

To determine an effective protein supplementation in maturation medium, the COCs were cultured in TCM 199 supplemented with 10% FBS, 10% OCS (Oestrous Cow Serum) or 3% BSA (Bovine serum albumin) and 5µg/ml FSH and 1µg/ml OE<sub>2</sub>. Each experiment was repeated for 8 times across the days.

### Statistical analysis

The data were recorded in Microsoft Excel spread sheet and descriptive statistics was performed. The maturation rates were expressed as mean ± SD and Z test was done to determine the significant effect of different treatments on the maturation rate using SPSS Version 20. The difference between groups was considered significant when P-value was < 0.05.

## RESULTS AND DISCUSSION

The objectives of the present study were to select an effective basic medium and its effective hormone and protein supplementation for IVM of oocytes of indigenous zebu cows. A total of 2340 oocytes were collected from 585 ovaries and the mean number of oocytes collection from each ovary was 4.00. The present mean number of retrieved oocytes per ovary is higher than that of previous study (Morshed *et al.*, 2014; Singha *et al.*, 2015; Choudhury *et al.*, 2017) and lower than that of an earlier study (Talukder *et al.*, 2008). The reason for variations in oocyte retrieval rate among studies may be due to variation in skillness of follicle aspirators. Moreover, seasons of oocytes retrieval and cyclic status of cows may influence the oocyte retrieval rate from ovaries (Dode and Adona, 2001). Additionally, an increase in FSH level in blood may influence the number of oocytes retrieved (Fortune, 1994). Further, nutrition and temperature may influence the gonadotrophin concentrations and affect the population of follicles and number of oocytes retrieved (Zeitoun *et al.*, 1996).

In the present study, the rate of overall oocyte maturation was 67.6% (785/1161). The present maturation rate is lower than previous study reported by Singha *et al.* (2015) in indigenous zebu cows. However, contrasting to the present findings, lower maturation rate was reported by Morshed *et al.* (2014) (53.8%) and Das *et al.* (2006) (65.4%) in indigenous zebu cows. The reasons for variation in maturation rate among studies might be due to variation in basic media and percentage of serum supplementation in it used for oocyte maturation. Moreover, grades of oocytes may influence the *in vitro* maturation rates of oocytes as variation in rate of maturation *in vitro* was demonstrated between good and poor grade oocytes (Goswami 2002). However, all retrieved oocytes were cultured for maturation irrespective of grading which may contribute for obtaining lower maturation rate by Morshed *et al.*, (2014) than that of present study. In the present study, oocytes with at least 3 compact cumulus cell layers were used for maturation which might contribute to obtaining satisfactory rate of oocyte maturation *in vitro*.

### Experiment I. Determination of an effective basic medium for oocytes maturation

The maturation rate of oocytes in three basic media is presented in Table 1. The maturation rates of oocytes were 74.5 ± 4.2%, 67.7 ± 5.8% and 58.2±6.2 in TCM 199, mSOF and TALP medium, respectively. The deference in maturation rate was significant between TCM and TALP media (P<0.05).

Table1. Effects of basic maturation media on IVM rate of Zebu oocytes

Basic medium	Number of oocytes cultured	Number of oocytes matured	Maturation rate (%)
TCM	145	108	74.5 ± 4.2 <sup>a</sup>
mSOF	133	90	67.7 ± 5.8 <sup>ab</sup>
TALP	134	78	58.2 ± 6.2 <sup>b</sup>

Number of replicates is 10. Proportion values are mean ± SD.

<sup>a,b</sup> The values with superscripts within same column were significantly different from each other (P<0.05).

For maturation of oocytes *in vitro*, ingredients of a culture media play an important role. Different culture media such as TCM-199 (Kharche *et al.*, 2006; Amer *et al.*, 2008), SOF (Totey *et al.*, 1992), minimum essential medium (MEM) (Ravindranatha *et al.*, 2001) and Ham's F-10 (Totey *et al.*, 1993; Tamilman *et al.*, 2005) have been used for IVM of mammalian oocytes elsewhere. Among them, TCM 199 is the most widely used culture medium for such purposes (Arunakumari *et al.*, 2007). The beneficial effect of TCM-199 medium on IVM of animal oocytes may be attributed due to presence of some factors in its composition such as essential amino acids and glutamine that may stimulate DNA and RNA synthesis and enhance cell division (Pawshe *et al.*, 1996; Gordon, 2003). Moreover, there is a report that TCM 199 improved the rate of IVM of oocytes better than MEM in buffaloes (Roushandeh *et al.*,

2006). It has been known for many years that glucose and glutamine are poor energy substrates for the cumulus cell-free rodent oocytes (Downs and Verhoeven, 2003). The lower IVM rate in MEM may be explained by the fact that it contains higher glucose and glutamine than that of TCM 199. In the present investigation, the maturation rate was significantly higher in TCM than TALP media but did not vary with SOF medium ( $P > 0.05$ ). Similar to the present study, Prasad *et al.* (2013) did not obtain any difference in IVM rates when maturation rate of buffalo oocytes were compared between TCM 199 and SOF. Contrasting to the present study, Totey *et al.* (1992) reported higher percentage of oocytes maturation in TCM than that of SOF in buffaloes. However, when compared the rate of cleavage and blastocyst formation after IVF, there were no difference in cleavage and blastocyst formation in bovine oocytes matured either in defined TCM 199 or SOF (Lonergran *et al.*, 1994). The reasons for variations in IVM rate among studies may be due to variation in supplementations in basic media.

### Experiment 2. Determination of an effective hormone supplementation for oocytes maturation

The maturation rate of oocytes in two hormone supplemented media is presented in Table 2. The maturation rate of oocytes was higher ( $73.3 \pm 4.0\%$ ) in FSH supplemented TCM 199 than that of gonadotrophin supplemented counterpart ( $60.2 \pm 6.6\%$ ). The difference in maturation rate was statistically significant between two hormone supplementations ( $P < 0.05$ ).

Table 2. Effect of hormone supplementation on IVM rate of Zebu oocytes

Hormone supplementation	Number of oocytes examined	Number of oocytes matured	Maturation rate (%)
FSH	161	118	$73.3 \pm 4.0^a$
Gonadotrophin	161	97	$60.2 \pm 6.6^b$

Number of replicates is 8. Proportion values are mean  $\pm$  SD.

<sup>a,b</sup> The values with superscripts within same column were significantly different from each other ( $P < 0.05$ ).

In IVM, gonadotropins are the main stimulator of the oocyte development and FSH being deemed vital for the oocytes becomes qualified to be *in vitro* fertilized (Lu *et al.*, 2014; Khan *et al.*, 2015). In the present study, the maturation rate of oocytes was significantly ( $P < 0.05$ ) higher in FSH supplemented TCM 199 than that of gonadotrophin supplemented counterpart. Contrasting to the present study, although IVM rate was not compared, the embryo development rate after IVF did not vary between bovine oocytes cultured either in FSH or gonadotrophin supplemented medium (Groza *et al.*, 2008). Moreover, the IVM rate did not vary in buffalo oocytes matured in either FSH or PMSG supplemented medium (Hegab *et al.*, 2009). Further, no difference in IVM rate of bovine oocytes was observed when compared between FSH and LH supplementation (Younis *et al.*, 1989). Supplementation of reproductive hormones in maturation media is essential because it improves IVM rate of mammalian oocytes. Addition of FSH (Chauhan *et al.*, 1998), PMSG (Roy *et al.*, 1968), LH and oestradiol (Nandi *et al.*, 2002) to maturation media has been made to improve the developmental competence of *in vitro* matured oocytes. In many mammalian species, gonadotrophin has been found to stimulate cumulus cells to synthesize molecules able to drive germinal vesicle breakdown (GVBD) as meiosis activating sterols (Tsafiriri *et al.*, 2005). Oestradiol has been found to improve the completion of maturational changes and also to support the synthesis of presumed male pronuclear growth factor (Fukui and Ono, 1989). Moreover, FSH is essential for cumulus cell expansion and maturation of oocytes *in vitro* as it enhanced the expansion of cumulus cells in buffaloes (Alok *et al.*, 2010). Nevertheless, in addition to either FSH or gonadotrophin, oestradiol was always supplemented in maturation medium in the present study.

### Experiment 3. Determination of an effective protein supplementation for oocytes maturation

The maturation rate of oocytes in three protein supplemented media is presented in Table 3. The maturation rates of oocytes were  $73.0 \pm 5.9\%$ ,  $71.1 \pm 2.8$  and  $62.5 \pm 9.4$  in medium supplemented with FBS, OCS and BSA, respectively. However, the maturation rate did not vary among three protein supplementations ( $P > 0.05$ ).

Table 3. Effect of protein supplementation on IVM rate of Zebu oocytes

Protein supplementation	Number of oocytes examined	Number of oocytes matured	Maturation rate (%)
FBS	141	103	73.0 ± 5.9
OCS	142	101	71.1 ± 2.8
BSA	144	90	62.5 ± 9.4

Number of replicates is 8. Proportion values are mean ± SD.

The maturation rate was not significantly different from each other ( $P > 0.05$ ).

There are several known functions of protein in culture medium. Serum may provide energy substrates, amino acids, growth factors and vitamins to the culture medium. The positive effect of serum on maturation rate of oocytes may be due to presence of growth factors that is manifested by improved embryo development following IVF (Eppig *et al.*, 1992). Moreover, it is important to include serum in the IVM medium to prevent hardening of zona pellucida which could adversely affect fertilization (Downs *et al.*, 1986). Albumin chelates heavy metals and provides some  $p^H$  buffering, which can also be achieved by inclusion of amino acids in culture medium (Mehta and Kiessling, 1990). The surfactant property of albumin prevents adhesion of cells to glass and plastic surface (Pinyopummintr and Bavister, 1991) and also can act as a scavenger of reactive oxygen species (Natsuyama *et al.*, 1993). Protein supplements during IVM medium can have profound effects both on the rate of development and on the overall efficiency of development shown by the morula and blastocyst yields. (Atef ali and Marc- Andre Sirard, 2002). Additionally, there is a report that fetuin, a major glycoprotein constituent of fetal calf serum, can prevent hardening of zona pellucida during IVM (Schroeder *et al.*, 1990). Kan and Yamane (1983) reported another beneficial action of serum due to its antioxidant properties as evidenced by reducing superoxide formation. In addition, serum added to the maturation medium provides a source of albumin that balances the osmolarity (Thompson, 2000).

In the present study, the maturation rate of oocytes did not vary in OCS supplemented TCM 199 than that of FBS or BSA supplemented counterpart ( $P>0.05$ ). Contrasting to the present finding, Lu and Gordon (1987) reported that OCS had a significant and marked effect on oocytes maturation compared to FBS supplementation. Further, similar result to earlier report has been demonstrated elsewhere (Schellander *et al.*, 1990). There are different sources for supplemented sera such as FBS (Kobayeshi *et al.*, 1994; Nandi, 1998; Mahmoud and Nawito, 2005; Das *et al.*, 2006; Talukder *et al.*, 2008), OCS (Schellander *et al.*, 1990; Singha *et al.*, 2015), steer serum (Roy *et al.*, 1968; Nandi *et al.*, 2001) and superovulated cow serum (Boediono *et al.*, 1994). Although, there are reports to use different sera sources for supplementation in maturation medium, comparison was made within FBS, OCS and BSA only in the present study. Because, FBS, OCS and BSA are the most widely used protein supplementation for maturation culture of bovine oocytes in the world.

In conclusions, selection of a suitable oocytes culture medium is crucial for optimal results in bovine IVM/IVF. For IVM, both TCM and mSOF may be used as a basic medium supplemented with FBS, OCS or BSA as protein source and FSH may be used as a hormone supplement in basic medium for optimum IVM rate of indigenous Zebu cow's oocytes in Bangladesh.

#### ACKNOWLEDGEMENTS

The research work has been supported by the research project entitled “Optimization of *in vitro* maturation and fertilization of oocytes in the bovine in Bangladesh”, Funded by Ministry of Education, People's Republic of Bangladesh.

#### REFERENCES

1. Ahmed ZU and Islam TS (1987). Cattle breeding program through Artificial Insemination in Bangladesh. A report from Central Cattle Breeding Station, Savar, Dhaka. Department of Livestock Services, Bangladesh.
2. Ali A and Sirard MA. (2002) Effect of the absence or presence of various protein supplements on further development of Bovine oocytes during *in vitro* maturation. *Biology of Reproductin* 66: 901–905.
3. Alok P, Gupta SC and Neelam G (2010). Effect of FSH and LH hormones on oocyte maturation of buffalo and gene expression analysis of their receptors and Cx43 in maturing oocytes. *Zygote* 18: 231-234.

4. Amer HA, Hegab AO and Zaabal SM (2008). Effects of ovarian morphology on oocyte quantity and quality, granulose cells, *in vitro* maturation and steroid hormone production in buffaloes. *Animal Reproduction* 5: 55-62.
5. Arunakumari G, Vagdevi R, Rao BS, Naik BR, Naidu KS, Suresh Kumar RV and Rao VH (2007). Effect of hormones and growth factors on *in vitro* development of sheep preantral follicles. *Small Ruminant Research* 70: 93-100.
6. Bavister BD, Rose-Hellekant TA and Pinyopummintr T (1992). Development of *in vitro* matured/*in vitro* fertilized bovine embryos into morulae and blastocysts in defined culture media. *Theriogenology* 37: 127-146.
7. Boediono A, Takaji M, Saha S and Suzuki T (1994). Influence of day-0 and day-7 super ovulated cow serum during development of bovine oocytes *in vitro*. *Reproduction Fertility and Development* 6: 261-284.
8. Chauhan MS, Singla SK, Palta P, Manik RS and Madan ML (1998). *In vitro* maturation and fertilization and subsequent development of buffalo (*Bubalus bubalis*) embryos: Effects of oocyte quality and type of serum. *Reproduction Fertility and Development* 10: 173 - 177.
9. Choudhury SM, Bhuiyan MMU †, Rahman MM, Rahman MM, Sharif MN, Bhattacharjee J, Bari FY and Juyena NS (2017). Comparison between Two Cryo-devices for Vitrification of Immature Oocytes of Indigenous Zebu Cows in Bangladesh. *Journal of Embryo Transfer* 32:311-317.
10. Das ZC, Siddiqui MAR, Bhattacharjee, Bhuiyan MMU and Shamsuddin M (2006). *In vitro* fertilization using zebu and crossbred bull semen. Bangladesh Society for Veterinary Education and Research (BSVER) Publication 31: 31.
11. Dode MAN and Adona PR (2001). Developmental capacity of *Bos Indicus* oocytes after inhibition of meiotic resumption by 6-dipethylaminopurine. *Animal Reproduction Science* 65: 171-180.
12. Downs SM and Verhoeven A (2003). Glutamine and the maintenance of meiotic arrest in mouse oocytes: Influence of culture medium, glucose and cumulus cells. *Molecular Reproduction and Development* 66: 90-97.
13. Downs SM, Schroeder AC and Epping JJ (1986). Serum maintains the fertilizability of mouse oocyte *in vitro* by preventing hardening of zona. *Gamete Research* 15: 115-122.
14. Eppig J, Wigglesworth JK and Brien MJO (1992). Comparison of embryonic development competence of mouse oocytes grown with and without serum. *Molecular Reproduction and Development* 32: 33-40.
15. Fortune JE (1994). Ovarian follicular growth and development in mammals. *Biology of Reproduction* 50: 225-232.
16. Fukui Y and Ono H (1989). Effects of sera, hormones and granulosa cells added to culture medium for *in vitro* maturation, fertilization, cleavage and development of bovine oocytes. *Journal of Reproduction and Fertility* 86: 501-506.
17. Gordon I (2003). Laboratory production of cattle embryos: Maturing the oocyte, 2th edition, Cambridge, MA 02138, CABI publishing.
18. Goswami PC (2002). Collection and grading of bovine cumulus oocyte-complexes (COCs) from slaughterhouse ovaries *in vitro* maturation, fertilization and culture. *MS Thesis*, Department of Animal Breeding and Genetics, Bangladesh Agricultural University, Mymensingh.
19. Groza i, Ciupe S, Stegeran B, Cenariu M, Petrean A and Ercean E (2008) Researches regarding the optimization of the *in vitro* maturation protocol of bovine oocytes used for *in vitro* fertilization, Bulletin of the University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, *Veterinary Medicine* 65:119-123.
20. Hegab AO, Montasser AE, Hammam AM, Abu EI-Naga EMA and Zaabel SM (2009). Improving *in vitro* maturation and cleavage rates of buffalo oocytes. *Animal Reproduction* 6: 416-421.
21. Islam MF, Bhuiyan MMU, Banu TA, Bhattacharjee J and Shamsuddin M (2007). *In vitro* maturation and fertilization of oocytes in zebu cattle. *Bangladesh Veterinary Journal* 41: 58-76.
22. Kan M and Yamane I (1983). Oxygen-dependent growth declining and effect of vitamin E for human diploid fibroblasts in serum-free, BSA containing culture. *The Tohoku Journal of Experimental Medicine* 139: 389-398.
23. Khan DR, Guillemette C, Sirard MA and Richar FJ (2015) Characterization of FSH signaling networks in bovine cumulus cells: a perspective on oocyte competence acquisition. *Molecular Human Reproduction* 21:688-701.
24. Kharche SD, Goel AK, Jindal SK and Sinha NK (2006). Technical note: *in vitro* maturation of caprine oocytes in different concentration of estrous goat serum. *Small Ruminant Research* 64: 186-189.
25. Kobayeshi K, Yamashita S and Hoshi H (1994). Influence epidermal growth factor and transforming growth factor-  $\alpha$  on *in vitro* maturation of cumulus cell-enclosed bovine oocytes in a defined medium. *Journal of Reproduction Fertility* 100: 439-446.
26. Lonergan P, Carolan C and Mermillod P (1994). Development of bovine embryos *in vitro* following oocyte maturation under defined conditions. *Reproduction Nutrition Development* 34: 329-339.
27. Lu CL, Wang TR, Yan LY, Xia X, Zhu Xu, Li R, Zhao HC, Yan J, Yin TL Jin HY, Zhang Y, Yhang, WX, Feng HL and Qiao J (2014).Gonadotropin-mediated dynamic alterations during bovine oocyte maturation *in vitro*. *Biology of Reproduction* 91:1-9.

*Effect of hormone and protein for in vitro maturation of oocytes of cows*

28. Lu KH and Gordon I (1987). Effect of serum, hormones and cumulus cells on the in vitro maturation of bovine oocytes. Proceeding 7<sup>th</sup> meeting of the European embryo transfer association (Cambridge) 63.
29. Mahmoud KGM and Nawito MF (2005). Influence of serum source on buffalo *in vitro* embryonic development with reference to heat inactivation. *Livestock Production Science* 96: 285-289.
30. Mehta TS and Kiessling AA (1990). Development potential of mouse embryos conceived in vitro and cultured in ethylenediaminetetraacetic acid with or without amino acids or serum. *Biology and Reproduction* 43:600-606.
31. Morshed NSM, Bhuiyan MMU, Rahman MM, Singh JK and Juyena NS (2014). Culture condition for *in vitro* maturation of abattoir derived oocytes of native zebu cows in Bangladesh. *Journal of Embryo Transfer* 29: 201-206.
32. Nandi S, Raghu HM, Ravindranatha BM and Chauhan MS (2002). Production of buffalo (*Bubalus bubalis*) embryos *in vitro*. Premises and promises. *Reproduction in Domestic Animal* 37: 65-74.
33. Nandi S, Chauhan MS and Palta P (1998). Influence of cumulus cells and sperm concentration on cleavage rate and subsequent embryonic development of buffalo (*Bubalus bubalis*) oocytes matured and fertilized *in vitro*. *Theriogenology* 50: 1251- 1262.
34. Natsuyama S, Noda Y, Narimoto K, and Tori T. (1993). Role of protein supplements in the culture of mouse embryos. *Theriogenology* 40:149-157.
35. Pawshe CH, Palanisamy A, Taneja M, Jain SK and Totey SM (1996). Comparison of various maturation treatments on *in vitro* maturation of goat oocytes and their early embryonic development and cell numbers. *Theriogenology* 46: 971-982.
36. Pinyopummintr T and Bavister BD (1991). In vitro maturation/ in vitro fertilization bovine oocytes can develop in to morulae/ blasyocysts in chemically defined protein- free culture medium. *Biology Reproduction* 45:736-742.
37. Prasad CS, Palanisamy A, Gomathy VS, kumar SS, Thangavel A and Dhinakar RG (2013). Effect of TCM-199 and synthetic oviductal fluid (SOF) Medium and cysteamine supplementation to in vitro maturation media on maturation, cleavage rate and subsequent embryonic development of buffalo oocytes. *Buffalo Bulletin* 32:182-188.
38. Ravindranatha BM, Nandi S, Gupta PSP and Sharma PV (2001). Comparison of three different media on maturation of buffalo oocytes *in vitro*. *Indian Journal Animal Science* 71: 841-843.
39. Roushandeh AM, Mooghahi MHN, Pasbakhsh P, Abddvahabi A, Akbari M, Shokrgozar A and Sobhani A (2006). Effect of cysteamine on the rate of *in vitro* maturation of oocytes in two media. *Acta Medica Iranica* 44: 167-171.
40. Roy A, Razada BC, Tiwari RBI, Andey MDP, Yadavand PO and Sengupta BP (1968). Effect of management on fertility of buffalo cows bred during summer. *Indian journal of Veterinary Science* 38: 554-560.
41. Schellander KF, Fuhrer BG, Brackett B, Korb H and Schleger W (1990). In vitro fertilization and cleavage of bovine oocytes mature in medium supplemented with oestrus cow serum. *Theriogenology* 33: 477-485.
42. Schroeder A, Schultz RM, Kopf GS, Taylor ER, Becker RB and Eppig JJ (1990). Fetuin inhibit zona pellucida hardening and conversion of ZP2=ZP2f during spontaneous mouse oocyte maturation in the absence of serum. *Biology of Reproduction* 43: 891.
43. Singha JK, Bhuiyan MMU, Rahman MM and Bari FY (2015). Comparison of culture media for in vitro maturation of oocytes of indigenous Zebu cows in Bangladesh. *Journal of Embryo Transfer* 30: 327-333.
44. Talukder AK, Shamsuddin M, Rahman MB, Bari FY and Parish JJ (2008). Normaland abnormal in vitro fertilization of zebu cattle oocytes. *Journal of Embryo Transfer* 24: 89-95.
45. Tamilmann G, Rao BS, Vegdevi R, Amarnath D, Naik BR, Mutharao M and Rao VH (2005). Nuclear maturation of bovine oocytes in cultured preantral follicles. *Small Ruminant Research* 60: 295-305.
46. Thompson JG (2000). *In vitro* culture and embryo metabolism of cattle and sheep embryos. *Animal Reproduction Science* 30: 273-280.
47. Totey SM, Pawshe CH and Singh GP (1993). Effects of bull and heparin and sperm concentrations on *in vitro* fertilization of buffalo (bubalus bubalis) oocytes matured *in vitro*. *Theriogenology* 39: 887-898.
48. Totey SM, Singh G, Taneja M, Pawshe CH and Talwer GP (1992). *In vitro* maturation, fertilization and development of follicular oocytes from buffalo (*Bubalus bubalis*). *Journal of Reproduction Fertility* 95: 597-607.
49. Tsafiriri A, Cao X, Ashkenazi H, Mottola S, Popliker M and Pomerantz SH (2005). Resumption of oocyte meiosis in mammals: On models, meiosis activating sterols, steroids and EGF like factors. *Molecular Cellular Endocrinology* 234: 37- 45.
50. Younis AI, Brackett BG and Fayer-Hosken RA (1989). Influence of serum and hormones on bovine oocyte maturation and fertilization in vitro. *Gamete Research* 23: 189-201.
51. Zeitoun MM, Rodriguez-Martinez H and Randel RD (1996). Effect of season on ovarian follicular dynamics in Brahman cows. *Theriogenology* 45: 1577-1581.