

## **Causal factors and treatment outcome of genitourinary fistula in a teaching hospital**

MK Fatema<sup>1</sup>, E Saha<sup>2</sup>, F Begum<sup>3</sup>, SN Lucky<sup>4</sup>, F Rahman<sup>5</sup>

### **Abstract**

**Background:** Urogenital fistula, majority of which is vesicovaginal fistula is a public health problem especially of developing countries. It is commonly caused by prolonged and obstructed labor. With the advancement of health care delivery system etiology of urogenital fistula is changing in our country.

**Objective:** This study was done to find out the causal factors and to determine the success rate of operative procedure for treatment of urogenital fistula.

**Method:** This was an observational type of cross sectional study, carried out in Department of Obs and Gynae in Khulna Medical College Hospital from January 2013 to June 2014 among 30 women at different ages suffering from urogenital fistula due to gynaecological or obstetrical causes or malignancy.

**Result:** Majority of the patients were young primipara, short stature and malnourished, coming from lower socioeconomic condition of rural areas. Prolonged labor 12 (40%) was the most common cause of urogenital fistula, followed by gynaecological surgeries mainly hysterectomies 9 (30%) and malignancy 2 (6.6%) rare cause. The success rate of repair following first and second attempt was 92% and 8% respectively. Overall success rate was 24 (96%) and functionally failed with a failure in 1 (4%) cases. This study showed main cause for developing fistula showing prolonged labor and majority of patient were treated pervaginally. Most of the cases (96%) had successful repair which is very encouraging and correlates well with recently published series.

**Conclusion:** Fistula can be totally preventable by proper antenatal care, identification of high risk cases, timely referral, proper intranatal, postnatal care, and proper training. So improvement of health care services and dedication will prevent this type of morbidity. The best results are obtained when repair of urogenital. fistulae is carried out under optimal conditions.

**Keywords:** Genito urinary fistula, Causal factor, Outcome.

### **Introduction**

An abnormal communication between urinary and genital tract is termed urogenital fistula. The commonest type of genitourinary fistula is vesicovaginal fistula. Vesicovaginal fistula is an abnormal fistulous communication tract between vagina and urinary bladder, which leads to continuous involuntary passage of urine into vagina. Genitourinary fistula is a major problem in many developing countries especially the vesicovaginal fistula commonly caused by prolonged and obstructed labor is one of the worst complication of child birth.<sup>1</sup> Genitourinary fistula is a devastating condition affecting the physical and psychological health of women. With advance

obstetric care these fistula are rare in industrialized world.

Globally about 3.5 million women are living with genitourinary fistula, a miserable condition.<sup>2</sup> An incidence of 12 per 1000 deliveries has estimated worldwide, with an annual incidence of up to 50,000 to 100,000. In Bangladesh 1.9 percent women are suffering from Vesicovaginal fistula (BIRRERTH).<sup>3</sup> According to UNFPA & Engender Health, the number of women living with fistula is estimated to be 1.69 per 1000 ever married women.<sup>4</sup> Vesicovaginal fistulas result from mainly obstetrical and gynaecological cause. In the developing countries, 80-90 percent vesicovaginal fistulas are of obstetrical origin.<sup>5</sup> Lacks of privacy

1. Muckti Kaniz Fatema FCPS, Junior consultant (Obs & Gynae), Khulna Medical College Hospital.

2. Eti Saha FCPS, Associate Professor (Obs & Gynae), Khulna Medical College.

3. Fouzia Begum MCPS, Consultant (Obs & Gynae), Khulna Medical College Hospital.

4. Shamsun Nahar Lucky MS, Professor (Obs & Gynae), Khulna Medical College.

5. Farzana Rahman FCPS, consultant (Obs & Gynae), Abedin Hospital, Dhaka.

n hospital, objection from families in different attitude of husband leads to prolong labor, encourage obstructed labor. Recently the incidence of genitourinary fistula following gynecological surgery, especially hysterectomy has increased.<sup>6</sup> Repair of the vesicovaginal fistula is a challenging task for the fistula surgeons worldwide. The outcome of vesicovaginal fistula repair depends on many factors like site, size and number of fistula, urethral length, bladder capacity and amount of scarring etc.<sup>7</sup>

This study was designed to find out the causal factors and to assess the outcome of vesicovaginal fistula repair in Khulna Medical College Hospital, Khulna.

## Methods

The study was observational cross sectional study & was done in the department of Obstetrics and Gynecology, KMCH, Khulna from January 2013 to June 2014 among admitted women (30) suffering from urogenital fistula. After admission all eligible patients were informed and written consent was obtained. Ethical clearance was taken from the ethical review committee of Khulna Medical College Hospital. Inclusion criteria: Patient admitted with signs of dribbling of urine due to history of previous operative procedure and obstructed or prolonged labor or malignancy. Exclusion criteria: patients also having rectovaginal fistula or complete perineal tear. A detailed history has taken to pin point the casual factor. This included respondent's personal information, socio economic condition, past obstetrical and medical history, pattern of medical care and problem faced by the women suffering from urogenital fistula. Each woman was evaluated with a detailed history as regards age, parity, antecedent event leading to fistula e.g., obstetric or gynecological. Obstetrical events including duration of labor, place, type of delivery, pregnancy outcome, and duration of fistula were detailed. The assessment included the women's general physical condition, size, site and number of fistula, amount of scarring of fistulous, margins or stenosis of the vagina. Detailed examination were done including examination under anaesthesia. After the assessment of the patient appropriate operative measure were taken.

All the result were noted in predesigned history sheet. Surgery was considered to be successful if patient can hold urine and there is no leakage of urine in between the act of voiding after removal of catheter and before discharge from the hospital. After collection of data, analysis was done by computer aided statistical software SPSS. Data

were presented in the form of tables and graphs. Data were analyzed with descriptive statistics. The level of significance of 0.05 were used for this study.

## Results

During a period of one and half year, 30 women that satisfied the inclusion criteria were prospectively evaluated & the results were analyzed. Among 7388 patients admitted in gynecological and obstetrics department of KMCH Khulna there were 30 urogenital fistula, yielding a frequency to be 0.4%, and among total Gynae patients of 1916 yielding a frequency to be 1.5%. Age distribution of patients (30)- Majority [11(36%)] patients belongs to age group 21-30 years, second commonest age was 31-40 years about 8(27%). Most of the patients [13 (43.3%)] in this study were young primipara, next [9 (30.0%)] were para (2-9) and 3(10%) of them were grand multipara. Distribution of cases according to aetiology of fistula (Table I): (A) Obstetrical causes 20 (66.66%) patients, among them prolonged labor (vaginal delivery) 13 (43.33%), obstructed labor (caesarean. section) 7 (23.33%). (B) Gynaecological causes 10 (33.33%) patients, among them total abdominal hysterectomy 9 (30.00%), malignancy 1 (33.33%).

Causative factors	Number of patients	%
(A) Obstetrical causes		
Prolonged labor (Vag deli)	13	43.33
Obstructed labor (Caes sec)	7	23.33
Total	20	66.66
(B) Gynae causes		
Total abdo Hysterectomy	9	30.00
Malignancy	1	3.33
Total	10	33.33

Table II shows distribution of cases according to types of fistula (n=30). Vesicovaginal fistula was the commonest and constituted 60%.

Types of fistula	Percentage	Number of fistula
Vesico vaginal	18	60
Vesico cervical	3	10
Uretero vaginal	9	30

**Table III**

Distribution of cases according to size (n=30)

Size of fistula	Number of patients	Percentage
Small (up to 2 cm)	16	53.3
Medium (2.1-3 cm)	9	30
Large (>3 cm.)	5	16

In this study 53.3% of patients had small size fistula (commonest) which is shown in Table III. Out of 30 patients with urogenital fistulae, 2 patients with small fistulae healed conservatively with continuous catheter drainage where one patient could not be operated upon because they suffered from carcinoma cervix in terminal stages. 2 patients were advised to come after 3 months because of presence of unhealthy granulation tissues around the fistulae. Therefore, 25 patients underwent surgery. The commonest approach used was transvaginal 17 (68%) and 8 patients (32%) were repaired through abdominal procedure with layer closure.

**Table IV**

Patient distribution as per previous attempt of repair (n=25)

Number of previous attempt	Number of patients	Percentage
No attempts	21	84
one attempt	13	12
Two attempt	1	4

Majority (84%) patients in this series had no history of previous attempts of repair. 12% patients had single attempt. Table IV shows previous attempts of repair.

**Table V**

Outcome of the study (n=30)

Outcome	Number of patients	Percentage	P value
Conservative management	5	16.67	
Surgical management			
Cured	24	96	<0.001
Failure	1	4	
Total	25	83.33	

Among 25 patients cure rate was 96% (n=25). In this study failure rate was 4% (Table-V). P value was <0.001 the test is significant.

Repair of fistula was difficult due to associated additional cofactor - vaginal stenosis [2 (8.8%)], attachment to pubic bone [2 (8.8%)], associated hydronephrosis [4 (16%)]. Post operative problems among patients were retention of urine due to catheter blockage [2 (8%)], urethral leakage [1 (4%)], UTI [4 (16%)]. In 8% patients catheter needed to be changed due to catheter block.

## Discussion

Obstetric fistula has gained international attention in the last 10-15 years. The condition has been researched mostly in developing countries like Nigeria, Ethiopia, Niger and Tanzania. Most studies of obstetric fistula uses observational, analytical study designs, mainly cross sectional studies. The research conducted are typically hospital based retrospective analysis of case records/patient records. Case control studies are not commonly used. The true incidence of urogenital fistula is unknown as many women do not reach hospital and continue to be neglected by their husbands and ostracized from society. Overall prevalence has been estimated at 0.2 to 2% in different societies. Its occurrence reflects the level of maternity care in a community and most are a consequence of mismanaged labor, a sequelae to obstructed labor. In my study period the frequency of urogenital fistula among all admitted patients of Obstetric and Gynae Department was 0.40% and among Gynae patients the frequency was 1.5%. The highest prevalence is in poor communities of Africa and Asia and constitute 0.5-1.7% of gynaecological admission in teaching hospital.<sup>6</sup>

Only three studies were identified using a case control design and that was one study in Nigeria, and a recent study in Zambia and in north eastern Nigeria.<sup>8-10</sup> There is a shared view that the main cause of obstetric fistula is prolonged labor and that the major outcome is stillbirths.<sup>11</sup> Research uncovers that many women are in labor for several days, often in the presence of a traditional birth attendant and little or no access to emergency obstetric care.<sup>12</sup> The majority of research highlights the following predispositions: low reproductive age, biological factors such as short stature and incomplete pelvis growth.<sup>10-13</sup> Cultural aspects such as female genital mutilation and socioeconomic aspects such as low education and poverty.<sup>8,12</sup> Many studies found malnutrition to be a risk factor, however some claimed that this needs to be further researched. Many studies found that VVFs occurred mostly in first pregnancies.<sup>13</sup>

However some studies did not agree with this.<sup>8</sup> The success rate of obstetric fistula repair is in general in between 80 to 90 percentile.<sup>13</sup> Success rate is generally lower for completely cured patients, sometimes as low as 60%.<sup>13</sup> One main predictor of surgery outcome is identified in some studies to be the amount of vaginal scarring.<sup>13,14</sup> The majority of iatrogenic fistulas develop subsequent to caesarean section. In our study the incidence of urological fistula arising out of obstetrical complication was 66.66% (20/30). In one study obstetrical complications were responsible for 88% cases and gynecological surgeries for 9% cases.<sup>6</sup>

One study conducted in Pakistan revealed a total number of 287 patients with genitourinary fistula.<sup>6</sup> The mean age of patients with urinary fistula was 31.5 + 7.5 years, parity was 4.2 ±2.8, and duration of labor was 38.4 ±6.5 hours. The most common type of urinary fistula was vesico vaginal fistula [250 (89.9%)]. A total of 268 patients underwent surgery. The success rate following first, second and third attempt was 85%, 91% and 96% respectively. In our study 25 patients underwent surgery among 30 patients. In our study prolonged labor with spontaneous delivery was responsible for 43.3% cases and obstructed labor with surgical interference was responsible for another 23.3%. So the study demonstrates the high rate of successful closure of the fistula in a specialized fistula unit, but highlights the problems of persistent urinary incontinence following closure.

There were some limitations in this study: period of study was insufficient to conduct a quality study, sample size was small that was not correlated with sample size calculation, it was a hospital based study, not representing the community population, this study was conducted in a single hospital which may not be the representative for the whole country.

### Conclusion

It was found that fistulas occurred mainly due to prolonged and obstructed labour, which can be totally preventable by proper antenatal care, identification of high risk cases, timely referral, proper intra natal and postnatal care. Iatrogenic fistula can be prevented by proper training. Repair of genito urinary fistula in appropriate time and in appropriate route by an expert surgeon can minimize failure rate.

### References

1. S Nahar, MK Sarker, KP Das, N Akter, Iatrogenic Genitourinary Fistulae: A survey in Khulna. Medical College. Journal of Bangladesh College of Physicians and Surgeons. 2005;23: 68-71.
2. Uprety D K. Subedi S. Budhatholki B. Regmi M. Vesicovaginal fistula at tertiary center at Eastern Nepal. J Nepal Med Ass .2008; 47: 120-22.
3. Sayeba A. Draft Training Manual on Female Genital Tract Fistula. 2004; 3-10.
4. Directorate General Health Service. Participants Hand Book for Doctors on Fistula Surgery and management Dhaka. 2008; 3-21.
5. Hiralal Konar. Genital Fistulae Textbook of DC Dutta's gynecology including Contraception, 6th ed. Daryaganj New Delhi 110002, India, Jaypee Brothers Medical Publishers (P) Ltd;2013: 417-429.
6. P.S. Sachdev, N. Hassan, RM. Abbasi, C.M. Das, Genito urinary fistula: a major morbidity in developing countries, J Ayub Medical College Abbottabad. 2009 ;21: 8-11.
7. Begum B, Khandakar S. Rahaman F. A study on outcome of VVF repair at Kumudini Women's Medical College Hospital. Bangladesh Medical Journal. 2011; 40:18-24.
8. Ezegwui HU. Nwogu IkqJo EE. Vesico vaginal fistula in Eastern Nigeria. Obstet Gynaecol. 2005; 25:589-591.
9. Porcaro AB, Zicari M, Zecchini AS, Pianon R, Monaco C. Migliorini F. Vesicouterine fistulas following cesarean section: report on a case, review and update of the literature. Int Urol Nephrol. 2002; 34: 335-344.
10. Melah GS, Massa AA, Yahaya UR, Bukar M, Kizaya DD, El Nafaty AU. Risk factors for obstetric fistulae in north eastern Nigeria. I Obstet Gynaecol 2007; 27: 819-823.
11. Muleta M. Obstetric fistula in developing countries: a review article. J Obstet Gynaecol Can. 2006; 28 : 962-966.
12. Wall LL. Obstetric vesicovaginal fistula as an international public health problem. Lancet 2006; 568: 1201-9.
13. Wall LL, Karshima JA, Kirschner C, Arrowsmith SD. The obstetric vesicovaginal fistula: characteristics of 899 patients from Jos, Nigeria. Am J Obstet Gynecol.2004; 190 : 1011-1019.
14. Holme A, Breen M, MacArthur C, Obstetric fistulae: a study of women managed at the Monze Mission Hospital, Zambia. BJOG. 2.007; 114: 1010-1017.