

RESEARCH PAPER

Short-term Evaluation of Bowel Dysfunction after Anterior Resection for Carcinoma Rectum

Md. Abdul Munim Sarkar^{1*}, Md. Setabur Rahman², AKM Minhaj Uddin Bhuiyan², Laila Shirin²,
Sonia Rahman³, Md. Washif Shakir⁴

¹Department of Surgery (Surgical Oncology), Rajshahi Medical College, Rajshahi, ²Department of Surgical Oncology, NICRH, Dhaka, Bangladesh, ³Department of Surgery, Rupganj UHC, Narayanganj, Bangladesh,

⁴Dept. of Surgery (Surgical Oncology), Rangpur Medical College, Rangpur, Bangladesh

Abstract

Background: Colorectal cancer (CRC) is the third commonest diagnosed cancer and the second leading cause of cancer-related mortality. Though anterior resection with total mesorectal excision (TME) is the gold standard for the treatment of non-disseminated rectal or distal sigmoid cancer, but almost all patients have experienced some degree of bowel dysfunctions.

Objective: To evaluate short-term bowel dysfunction by LARS score after anterior resection of carcinoma rectum.

Methods: This prospective observational study was carried out in the Department of Surgical Oncology, National Institute of Cancer Research & Hospital (NICRH), Mohakhali, Dhaka. The study was undertaken with the patients who already underwent AR for carcinoma rectum. Data collection was started from January 2020 to March 2021 and the 35 case was collected from outpatient department (OPD) and indoor of NICRH maintaining inclusion and exclusion criteria at 3rd month and 6th month. Statistical analysis was done according to the objective of the study by using Statistical Package for Social Science software version 27.0 for windows.

Results: Out of 35 eligible patients, the majority of 21 (60.0%) patients were male. The mean age was found 45.83±11.92 years. Regarding tumor distance from the anal verge, 8(22.86%) patients' tumors were found at the high rectum, 22(62.86%) at mid rectum (8-11 cm) and 5(14.29%) at low rectum. Bowel dysfunction was observed in 94% of cases; out of them, 37% had major dysfunction at 6th month. The mean LARS score was found 28.6±7.2 at 3rd month and 25.7±6.8 at 6th month. The difference was statistically significant ($p \leq 0.05$) between the two groups. Bowel dysfunction, i.e., Incontinence, clustering, urgency and frequency were documented at 6th month 74.29%, 77.14%, 71.43% and 82.86% respectively and were not statistically significant ($p > 0.05$) between at 3rd and 6th months. At 6th month, bowel dysfunctions were higher in ultra-low anterior resection than high anterior resection.

Conclusion: High prevalence of bowel dysfunction, i.e., Incontinence, clustering, urgency and frequency were documented in patients with carcinoma rectum after anterior resection.

Key words: Carcinoma Rectum, Bowel Dysfunction, Anterior Resection (AR), Anterior Resection Syndrome

Introduction:

Colorectal cancer (CRC) is the third most common diagnosed cancer and the second highest cause of cancer-related mortality. The World Health Organization (WHO) GLOBOCAN 2020 database estimates 1,931,590 (10%) new CRC cases and

935,173 (9.4%) CRC-related deaths in 2020. The number of new cases of carcinoma rectum (C19-C20) worldwide is 7,32,210, with an incidence rate of 7.6/100,000 and 3,39,022 fatalities. In Bangladesh, there are 2463 new cases.¹ The male-female sex ratio in South Asia is 9.8 and 5.6 per 100,000 people, respectively.²

The gold standard for treating non-disseminated rectal or distal sigmoid cancer is anterior resection (AR) with resection margins as short as 2 cm are sufficient for oncologic safety.^{3,4} The widespread use of Total Mesorectal Excision (TME) and perioperative multimodal

***Correspondence:** Dr. Md. Abdul Munim Sarkar, Department of Surgery (Surgical Oncology), Rajshahi Medical College, Rajshahi. Res. Address: 150/2, Upashahar Housing Estate, Raj Cantonment, Rajshahi-6202
Email- doc.atonu45@gmail.com
ORCID ID: 0000-0002-6327-643X

therapy has considerably improved the oncological outcomes of rectal cancer in recent decades.³

The quality of life after such a sphincter-saving treatment should surely be superior to that following abdominoperineal excision with a permanent colostomy (17) but it frequently results in bowel dysfunction, which sometimes lowering quality of life. This bowel dysfunctions occurs due to a complex mechanism of neorectum, nerve injury, anorectal sensory and reflex mechanisms (RAIR), pelvic muscle injury, loss of anal tone and other factors.⁹ For many years, the focus has been on maintaining urinary and, more specifically, male sexual function. However, in recent anterior resection, nerve preservation has been studied in more depth to avoid bowel dysfunction.¹⁰⁻¹⁶ This Bowel dysfunctions complex has been summarized under the term anterior resection syndrome (ARS) which is characterized by a high frequency of bowel movements, clustering, incomplete bowel evacuation, incontinence for flatus and stool and urgency.¹⁸ ARS was identified in 64% of patients, with 41% having significant severity.¹⁹ It was also predicted that 50-90% of individuals who underwent low anterior resection experienced some level of intestinal impairment after surgery.²⁰ Another study found substantial low anterior resection syndrome (LARS) in 46% of patients after a median follow-up of 14.6 years.²¹ The severity of ARS following rectal cancer surgery is strongly linked to lower quality of life (QoL).^{22,23}

It is also reported that patients with high rectal resection might have functional abdominal complaints^{24,25} There are a number of scoring tools for measuring ARS, including the Wexner score, St. Marks score, FSFI scores, MSKCC-BFI score, and LARS score. Given the intricacy of other scoring systems, the LARS score is now preferred for first-line ARS evaluation²⁶ This internationally validated LARS score is a simple five-question tool that was developed in Denmark in 2012 and validated for an English translation in 2014. The LARS score produces numerical values ranging from 0 to 42. Scores of 0-20 are considered as “no LARS,” 21–29 as “minor LARS,” and 30-42 as “major LARS” (27).

Age of the patients, sex, the extent of rectal excision (TME/PME), the height of the anastomosis, neoadjuvant therapy, diverting stoma, post-operative complications are important predictive factors for gastrointestinal functional results^{28,18} The occurrence of functional abdominal complaints and the effect on

QoL has not been investigated extensively for rectal cancer. As a result, function after AR has been inconsistently assessed and insufficiently documented in both research and the daily clinical setting, resulting in a large variation in reported frequency and severity of symptoms.

Increasing knowledge about Anterior Resection Syndrome (ARS) among specialists may implement in systemic screening tools. Better preparation about the possible burden of ARS symptoms, active attitude towards discussing the impact of symptoms and to reinforce social support if possible. The findings of this study might be helpful to provide evidence-based information to the physician as well as patient groups for bowel dysfunction and treatment planning.

Materials and Methods

This prospective observational study was undertaken in the department of Surgical Oncology, National Institute of Cancer Research & Hospital (NICRH), Mohakhali, Dhaka with the patients who already underwent AR for carcinoma rectum from January 2020 to March 2021. Our objective was to evaluate pattern of bowel dysfunction after anterior resection by LARS score and to determine the severity of symptoms among the type of operation. Subjects were collected from OPD and indoor of NICRH and metastatic diseases were excluded from our study. The purpose and procedure of the study was discussed with the patient. All clinical information was recorded from patient's case record. Curative AR for rectal cancer usually performed according to the principle of TME/PME by open or laparoscopic method, and a straight (end-to-end) anastomosis done routinely using a stapling device. A diverting ileostomy was constructed where indicated and generally reversed 3 to 6 months after primary surgery. Data was collected on 3rd month and 6th month after definitive surgery (without covering ileostomy) or after ileostomy reversal. Written consent was taken from those who agreed to participate in this study. On receipt of the informed written consent, data were collected from the patients on variables of interest using the structured design interview, history, clinical examination and LARS score sheet by the principal investigator. Age, sex, type of surgery, extent of operative procedure: TME / PME and reversal of covering ileostomy were the independent variable. Evacuation difficulties (clustering), incontinence of flatus and / or feces, urgency, frequency of bowel movement were our outcome variables.

All data was collected using a structured questionnaire and LARS score sheet. Statistical analysis was done according to the objective of the study by using IBM SPSS (Statistical Package for Social Science) software version 27.0 for windows and graphs by MS Excel 2010. Continuous data was expressed as mean $\pm$ standard deviation (SD). The analysis was done by using independent sample t test for continuous variables and chi-square test for categorical variables. All reported *p*-values were two sided and value less than 0.05 taken significant and confidence interval at 95% level.

LARS score questionnaire

The aim of this questionnaire is to assess your bowel function. Please tick only one box for each question. It may be difficult to select only one answer, as we know that for some patients' symptoms vary from day to day. We would kindly ask you to choose one answer which best describes your daily life. If you have recently had an infection affecting your bowel function, please do not take this into account and focus on answering questions to reflect your usual daily bowel function.

Q. 1: Do you ever have occasions when you cannot control your flatus (wind)?

- | | |
|--------------------------------|---|
| • No, never | 0 |
| • Yes, less than once per week | 4 |
| • Yes, at least once per week | 7 |

Q. 2: Do you ever have any accidental leakage of liquid stool?

- | | |
|--------------------------------|---|
| • No, never | 0 |
| • Yes, less than once per week | 3 |
| • Yes, at least once per week | 3 |

Q. 3: How often do you open your bowels?

- | | |
|--|---|
| • More than 7 times per day (24 hours) | 4 |
| • 4 - 7 times per day (24 hours) | 2 |
| • 1 - 3 times per day (24 hours) | 0 |
| • Less than once per day (24 hours) | 5 |

Q. 4: Do you ever have to open your bowels again within one hour of the last bowel opening?

- | | |
|--------------------------------|----|
| • No, never | 0 |
| • Yes, less than once per week | 9 |
| • Yes, at least once per week | 11 |

Q. 5: Do you ever have such a strong urge to open your bowels that you have to rush to the toilet?

- | | |
|--------------------------------|----|
| • No, never | 0 |
| • Yes, less than once per week | 11 |
| • Yes, at least once per week | 16 |

The allocated points per question are indicated in the right-hand column, the score from each of the five answers is added together to give a final score between 0-- 42.

Interpretation: 0-20 = No LARS 21-29 = Minor LARS 30-42 = Major LARS

Result

In this study, a cohort of 35 cases undergoing anterior resection for carcinoma of rectal within a defined time frame was included and followed up meticulously after taking ethical approval and meeting enrollment criteria.

A notable observation is that approximately one-third of the patients (31.43%) were in the age range of 41-50 years, with the mean age being 45.83 ± 11.92 years and a broad age distribution ranging from 23 to 67 years. The gender distribution revealed a majority of male patients (60.0%), while 40% were female. (Table I)

It was observed that tumors were found in 22.86% of cases at the high rectum, 62.86% at the mid rectum (8-11 cm), and 14.29% at the low rectum. In our study 68.57% of cases were underwent for low anterior resection, 17.14% cases for high anterior resection, and 14.29% cases for ultra-low anterior resection. (Table I)

In this study, incontinence, clustering, urgency and frequency at 6th month were following as 26(61.90%), 27(64.29%), 25(59.52%), and 29(69.05%), respectively. This result was not statistically significant ($p > 0.05$) from 3rd month (Table II).

The mean LARS score was found 28.6 ± 7.2 (Range- 4-42) in 3rd month and 25.7 ± 6.8 (Range- 2-39) in 6th month. The difference was statistically significant ($p < 0.05$) between two groups (Table III).

At sixth month of follow up, different age group did not show any statistically significant ($p < 0.05$) result for ARS. In case of high anterior resection group, 2 (100.0%) patients were in no LARS, 3 (15.0%) in minor LARS and 1(7.7%) in major LARS group. Five (38.5%) patients were found of Ultra-low anterior resection in major LARS but not found in no LARS and minor LARS groups. In low anterior resection group 0 (0.00%) patients was found in no LARS group, 17 (85%) were in minor LARS and 7(53.8%) in major LARS group. The difference was statistically significant ($p < 0.05$). (Table IV)

Table I: Baseline characteristics of the study population (n=35)

	Number of patients	Percentage
Age (years)		
≤30	5	14.29
31-40	7	20.00
41-50	11	31.43
51-60	7	20.00
>60	5	14.29
Mean ±SD	45.83(±11.92)	Range 23-67 years
Sex		
Male	21	60.0
Female	14	40.0
Tumor distance from anal verge		
Low (d<7 cm)	5	14.29
Mid (8-11 cm)	22	62.86
High (>11 cm)	8	22.86
Type of anterior resection		
High anterior resection	6	17.14
Low anterior resection	24	68.57
Ultra-low anterior resection	5	14.29
Mesorectal excision		
TME	27	77.14
PME	8	22.86
Stoma closure		
>24 weeks	23	65.71
≤24 weeks	08	22.86
No	04	11.43

Note. LARS: Low Anterior Resection Syndrome; TME: Total Mesorectal Excision; PME: Partial Mesorectal Excision;

Table II: Anterior resection syndrome in different follow up (n=35)

Anterior resection syndrome	At 3 rd months(n=43)		At 6 th months (n=43)		p-value
	n	%	n	%	
Incontinence	29	82.86	26	74.29	0.382 ^{ns}
Clustering	27	77.14	27	77.14	1.0 ^{ns}
Urgency	26	74.29	25	71.43	0.788 ^{ns}
Frequency	31	88.57	29	82.86	0.495 ^{ns}

(Multiple responses considered); ns= not significant
p-value reached chi square test

Table III: LARS score in different follow up (n=35)

LARS score	LARS	At 3 rd months(n=43)		At 6 th months (n=43)		p-value
		n	%	n	%	
No LARS (0-20).	No syndrome	2	5.71	2	5.71	
Minor LARS (21-29)	Minor	15	42.86	20	57.14	
Major LARS (30-42)	Major	18	51.43	13	37.14	
Mean ± SD		28.6 ± 7.2		25.7 ± 6.8		0.001 ^s
Range (min-max)		4.0 - 42.0		2.0 - 39.0		

s= significant; p-value reached paired t-test

Note. LARS: Low Anterior Resection Syndrome

Table IV: Type of resection with LARS score at 6th months (n=35)

	LARS at 6 th months					
	No (0-20) (n=3)		Minor (21-29) (n=24)		Major (30-42) (n=16)	
	n	%	n	%	n	%
Type of anterior resection						
High anterior resection	2	100.0	3	15.0	1	7.7
Low anterior resection	0	0.0	17	85.0	7	53.8
Ultra-low anterior resection	0	0.0	0	0.0	5	38.5
Mesorectal excision						
TME	0	0.0	16	80.0	11	84.6
PME	2	100.0	4	20.0	2	15.4

Note. LARS: Low Anterior Resection Syndrome; TME: Total Mesorectal Excision; PME: Partial Mesorectal Excision

TME procedures also demonstrated noteworthy associations with LARS outcomes. After TME, 80% had minor LARS and 84.6% had major LARS, with no patients in the no LARS group, and this difference was statistically significant ($p < 0.05$). (Table IV)

Discussion

The present study aimed to investigate the patterns of bowel dysfunction after anterior resection in patients with rectal cancer. The findings shed light on the prevalence and severity of various forms of bowel dysfunction, as well as surgical procedure that may contribute.

In terms of patient demographics, one-third of the patients fell within the 41-50 age range, which may suggest that this age group is particularly susceptible to rectal cancer. The male-to-female ratio was approximately 3:2, consistent with previous studies.³⁰

As the tumour were more in mid and lower rectum, patients underwent for TME and more the LARS was found. This distribution differs from the findings of other study, who observed a higher proportion of tumors in the upper rectum.³¹ The discrepancy may be attributed to delayed visits to specialists, resulting in the disease being more advanced at the time of diagnosis.

Regarding surgical procedures, after 6th month, in the case of LAR, and ULAR group, LARS results were more. A recent study (19) observed that LARS was more significant in low anterior resection 20(93.5%) and in ultra-low anterior resection 19 (86%). This observation is similar to our study.

Almost all patients (94%) experiencing some degree of bowel dysfunction in our study. Major dysfunction

was observed in 37% of patients, while 57% experienced minor dysfunction. These rates align with the observations made by Pieniowski³² and Van Heinsbergen,³³ indicating a high burden of bowel dysfunction following anterior resection. The mean LARS score at the 6th month was significantly lower than at the 3rd month, suggesting some improvement over time.

Tables II, and III provide a comprehensive overview of the postoperative outcomes and complications, highlighting factors such as incontinence, clustering, urgency, frequency, and the LARS score at the 3rd and 6th months post-surgery. Notably, the mean LARS score demonstrated a statistically significant difference between the two time points. In comparison with other study, the most frequently occurring symptoms were urgency (35%), incontinence of flatus or liquid (51% and 19%), and clustering (58%).²² This dissimilarity may be explained by our patients presented with more advanced stage, CCRT was given to almost all patients and more radical surgery was done.

It is our limitation that initial bowel dysfunction due to position of cancer just after surgery were not included in our study. It may affect the results. Association with other factors that may also affect the result like neoadjuvant chemoradiation, Prognostic stage, TNM category, Stoma closure were not included in our study as we had small number of study population.

Conclusion

This study provides valuable insights into the patterns of bowel dysfunction following anterior resection for rectal cancer. The high prevalence of bowel dysfunction, including incontinence, clustering,

urgency, and frequency found in this study; underscores the long-term impact on patients' quality of life. Factors such as tumor location, surgical approach, neoadjuvant therapy, timing of stoma closure, and disease may influence the development and severity of bowel dysfunction.

Recommendation:

Large scale multicenter studies, are warranted to further explore these factors and identify strategies to minimize bowel dysfunction and improve patient outcomes after anterior resection in carcinoma rectum. The high prevalence of ARS emphasizes the importance of evaluating bowel dysfunction for rectal cancer treatment and for counseling these patients in hospital and daily clinical practice.

Acknowledgement

The authors thanks to all co-authors and the department of surgical oncology, NICRH. Special thanks to Bangladesh Medical Research Council (BMRC) for kind co-operation with financial support.

Conflict of interest: No conflict of interest

Funding: BMRC

Ethical Clearance: National Cancer Research Institute & Hospital (NICRH), Mohakhali, Dhaka

Submit Date: 06 February, 2024

Accepted: 23 February, 2025

Final Revision Received: 27 February, 2025

Publication: July 2025

References:

1. WHO-IARC (2020) , *The Global Cancer Observatory*. Available at: <https://gco.iarc.fr/today/data/factsheets/cancers/9-Rectum-fact-sheet.pdf> (cited on: April 20,2020).
2. WHO (2020) *No Title, world cancer report*. Available at: <https://www.who.int/news-room/fact-sheets/detail/cancer> (cited on: April 20,2020).
3. van der Heijden JA, Thomas G, Caers F, van Dijk WA, Slooter GD, Maaskant-Braat AJ. What you should know about the low anterior resection syndrome—clinical recommendations from a patient perspective. *European Journal of Surgical Oncology*. 2018;44:1331-7. DOI: 10.1016/j.ejso.2018.05.010
4. Lakkis Z, Manceau G, Bridoux V, Brouquet A, Kirzin S, Maggiori L, de Chaisemartin C, Lefevre JH, Panis Y, French Research Group of Rectal Cancer Surgery (GRECCAR) and the French National Society of Coloproctology (SNFCP). Management of rectal cancer: the 2016 French guidelines. *Colorectal Disease*. 2017;19:115-22. DOI: 10.1111/codi.13550
5. Rödel C, Graeven U, Fietkau R, Hohenberger W, Hothorn T, Arnold D, Hofheinz RD, Ghadimi M, Wolff HA, Lang-Welzenbach M, Raab HR. Oxaliplatin added to fluorouracil-based preoperative chemoradiotherapy and postoperative chemotherapy of locally advanced rectal cancer (the German CAO/ARO/AIO-04 study): final results of the multicentre, open-label, randomised, phase 3 trial. *The Lancet Oncology*. 2015;16:979-89. DOI: 10.1016/S1470-2045(15)00159-X
6. Marwan K, Staples MP, Thursfield V, Bell SW. The rate of abdominoperineal resections for rectal cancer in the state of Victoria, Australia: a population-based study. *Diseases of the colon & rectum*. 2010;53:1645-51. DOI: 10.1007/DCR.0b013e3181f46485
7. Sauer R, Becker H, Hohenberger W, Rödel C, Wittekind C, Fietkau R, Martus P, Tschmelitsch J, Hager E, Hess CF, Karstens JH. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *New England Journal of Medicine*. 2004;351:1731-40. DOI: 10.1056/NEJMoa040694
8. van Gijn W, Marijnen CA, Nagtegaal ID, Kranenbarg EM, Putter H, Wiggers T, Rutten HJ, Pahlman L, Glimelius B, van de Velde CJ, Dutch Colorectal Cancer Group. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer: 12-year follow-up of the multicentre, randomised controlled TME trial. *The lancet oncology*. 2011;12:575-82. DOI: 10.1016/S1470-2045(11)70097-3
9. PEMBERTON JH. Anatomy and physiology of the anus and rectum. *Fundamentals of anorectal surgery-Beck DE*. 1992:1-24.
10. Moriya Y. Function preservation in rectal cancer surgery. *International journal of clinical oncology*. 2006; 11:339-43. DOI: 10.1007/s10147-006-0608-z
11. Runkel N, Reiser H. Nerve-oriented mesorectal excision (NOME): autonomic nerves as landmarks for laparoscopic rectal resection. *International journal of colorectal disease*. 2013;28:1367-75. DOI: 10.1007/s00384-013-1705-x
12. Maurer CA, Z'graggen K, Renzulli P, Schilling MK, Netzer P, Büchler MW. Total mesorectal excision preserves male genital function compared with conventional rectal cancer surgery. *British journal of surgery*. 2001;88:1501-5. DOI: 10.1046/j.0007-1323.2001.01904.x
13. Lange MM, Maas CP, Marijnen CA, Wiggers T, Rutten HJ, Kranenbarg EK, Van De Velde CJ. Urinary dysfunction after rectal cancer treatment is mainly caused by surgery. *Journal of British Surgery*. 2008;95:1020-8. DOI: 10.1002/bjs.6126
14. Kinugasa Y, Arakawa T, Murakami G, Fujimiya M, Sugihara K. Nerve supply to the internal anal sphincter differs from that to the distal rectum: an immunohistochemical study of cadavers. *International journal of colorectal disease*. 2014;29:429-36. DOI: 10.1007/s00384-013-1811-9

15. Stelzner S, Wedel T. Anatomic principles of nerve-sparing rectal surgery. *Coloproctology*. 2015;37:240-7. DOI: 10.1007/s00053-015-0030-y
16. Stelzner S, Böttner M, Kupsch J, Kneist W, Quirke P, West NP, Witzigmann H, Wedel T. Internal anal sphincter nerves—a macroanatomical and microscopic description of the extrinsic autonomic nerve supply of the internal anal sphincter. *Colorectal Disease*. 2018 ;20:07-16 DOI: 10.1111/codi.13942
17. Bryant CL, Lunniss PJ, Knowles CH, Thaha MA, Chan CL. Anterior resection syndrome. *The lancet oncology*. 2012;13:e403-8. DOI: 10.1016/S1470-2045(12)70236-X
18. Sarcher T, Dupont B, Alves A, Menahem B. Anterior resection syndrome: what should we tell practitioners and patients in 2018?. *Journal of visceral surgery*. 2018;155:383-91. DOI: 10.1016/j.jvisc Surg.2018.03.006
19. Sturiale A, Martellucci J, Zurli L, Vaccaro C, Bruscianno L, Limongelli P, Docimo L, Valeri A. Long-term functional follow-up after anterior rectal resection for cancer. *International journal of colorectal disease*. 2017;32:83-8. DOI: 10.1007/s00384-016-2659-6
20. Juul T, Ahlberg M, Biondo S, Espin E, Jimenez LM, Matzel KE, Palmer GJ, Sauermann A, Trenti L, Zhang W, Laurberg S. Low anterior resection syndrome and quality of life: an international multicenter study. *Diseases of the Colon & Rectum*. 2014;57:585-91. DOI: 10.1097/DCR.0000000000000116
21. Chen TY, Wiltink LM, Nout RA, Kranenbarg EM, Laurberg S, Marijnen CA, van de Velde CJ. Bowel function 14 years after preoperative short-course radiotherapy and total mesorectal excision for rectal cancer: report of a multicenter randomized trial. *Clinical colorectal cancer*. 2015;14:106-14. DOI: 10.1016/j.clcc.2014.12.007
22. Battersby NJ, Juul T, Christensen P, Janjua AZ, Branagan G, Emmertsen KJ, Norton C, Hughes R, Laurberg S, Moran BJ. Predicting the risk of bowel-related quality-of-life impairment after restorative resection for rectal cancer: a multicenter cross-sectional study. *Diseases of the Colon & Rectum*. 2016;59:270-80. DOI: 10.1097/DCR.0000000000000552
23. Juul T, Ahlberg M, Biondo S, Emmertsen KJ, Espin E, Jimenez LM, Matzel KE, Palmer G, Sauermann A, Trenti L, Zhang W. International validation of the low anterior resection syndrome score. *Annals of surgery*. 2014;259:728-34. DOI: 10.1097/SLA.0b013e31828fac0b
24. Ekkarat P, Boonpipattanapong T, Tantiphlachiva K, Sangkhathat S. Factors determining low anterior resection syndrome after rectal cancer resection: a study in Thai patients. *Asian journal of surgery*. 2016;39:225-31. DOI: 10.1016/j.asjsur.2015.07.003
25. Ihn MH, Kang SB, Kim DW, Oh HK, Lee SY, Hong SM. Risk factors for bowel dysfunction after sphincter-preserving rectal cancer surgery: a prospective study using the Memorial Sloan Kettering Cancer Center bowel function instrument. *Diseases of the colon & rectum*. 2014;57:958-66. DOI: 10.1097/DCR.0000000000000163
26. Chen TY, Emmertsen KJ, Laurberg S. What are the best questionnaires to capture anorectal function after surgery in rectal cancer?. *Current colorectal cancer reports*. 2015;11:37-43. DOI: 10.1007/s11888-014-0217-6
27. Emmertsen KJ, Laurberg S. Low anterior resection syndrome score: development and validation of a symptom-based scoring system for bowel dysfunction after low anterior resection for rectal cancer. *Annals of surgery*. 2012;255:922-8. DOI: 10.1097/SLA.0b013e31824f1c21
28. Bondeven P, Emmertsen KJ, Laurberg S, Pedersen BG. Neoadjuvant therapy abolishes the functional benefits of a larger rectal remnant, as measured by magnetic resonance imaging after restorative rectal cancer surgery. *European Journal of Surgical Oncology (EJSO)*. 2015;41:1493-9. DOI: 10.1016/j.ejso.2015.07.003
29. Talukder MH, Islam M, Kamal M, Laila N, Shihabuddin M. Hospital Cancer Registry Report, NICRH, Dhaka 2015-2017, National Institute of Cancer Research and Hospital, Dhaka, Bangladesh, 2020. DOI:10.13140/RG.2.2.10796.08320
30. Kupsch J, Jackisch T, Matzel KE, Zimmer J, Schreiber A, Sims A, Witzigmann H, Stelzner S. Outcome of bowel function following anterior resection for rectal cancer—an analysis using the low anterior resection syndrome (LARS) score. *International journal of colorectal disease*. 2018;33:787-98. DOI: 10.1007/s00384-018-3006-x
31. Bregendahl S, Emmertsen KJ, Lous J, Laurberg S. Bowel dysfunction after low anterior resection with and without neoadjuvant therapy for rectal cancer: a population based cross sectional study. *Colorectal disease*. 2013;15:1130-9. DOI: 10.1111/codi.12244
32. Pieniowski EH, Palmer GJ, Juul T, Lagergren P, Johar A, Emmertsen KJ, Nordenvall C, Abraham-Nordling M. Low anterior resection syndrome and quality of life after sphincter-sparing rectal cancer surgery: a long-term longitudinal follow-up. *Diseases of the Colon & Rectum*. 2019;62:14-20. DOI: 10.1097/DCR.0000000000001228
33. Van Heinsbergen M, Den Haan N, Maaskant Braat AJ, Melenhorst J, Belgers EH, Leijtens JW, Bloemen JG, Rutten HJ, Bouvy ND, Janssen Heijnen ML, Konsten JL. Functional bowel complaints and quality of life after surgery for colon cancer: prevalence and predictive factors. *Colorectal Disease*. 2020;22:136-45. DOI: 10.1111/codi.14818