

Residual Ventricular Septal Defect after Surgical Repair of TOF—A Case Report

MK Hassan¹, KA Hasan¹, AMA Rahim¹, KN Mahmood¹, SN Hossain²

¹ Department of Cardiac Surgery, NICVD, ² Department of Cardiac Surgery, CMCH, Chittagong

Abstract:

Residual Ventricular septal defect after surgical repair for Tetralogy of fallot(TOF) can occasionally be hemodynamically important requiring re-intervention. Closed observation and followup make this defect hemodynamically insignificant, required no medication and no endocarditis. We describe one patient having residual defect after surgical repair of TOF.

(Cardiovasc. j. 2011; 3(2): 233-234)

Keywords

TOF,
Residual VSD

Introduction:

A ventricular septal defect is a consistent component of tetralogy of fallot(TOF) and is routinely closed during surgical correction of this more complex defect. Small residual defects are frequently described on postoperative transthoracic echocardiography (TTE) but the rate of possible spontaneous closure or the hemodynamic and clinical significance in case of persistent residual shunt, are sparsely documented.¹ Here we evaluated a patient of intracardiac repair of TOF with regard to residual shunt and looked at the rate of spontaneous closure in time with its eventual implications for the patient in midterm follow-up.

Case History:

A seven years old boy admitted in National Institute of Cardiovascular Diseases and Hospital(NICVD) Dhaka, with a diagnosis of tetralogy of fallot(TOF) for intracardiac repair. He looked cyanosed with clubbing, 120 mm of hg systolic pressure, 75 beats per minute of heart rate, and 20 breaths per minute of respiratory rate. On auscultation, revealed features suggestive of severe pulmonary stenosis. Chest skiagram showed typical boot shaped heart with oligemic lung fields, echocardiographic finding was tetralogy of Fallot with relatively small sized branch of pulmonary arteries and a 15mm subaortic ventricular septal defect(VSD) with right to left shunt. Cardiac Catheterization showed 120 mm of hg right ventricular pressure and severe infundibular & valvular stenosis with a malaligned VSD with Right to Left shunt (R—L). This patient underwent surgery for closure of malaligned VSD, resection of infundibular muscle bands, standard

surgical technique using cardiopulmonary bypass(CPB) with moderate hypothermia, cross clamping and cold blood cardioplegia were used.



Fi.-1: *Echocardiography showing Tetralogy of Fallot.*

Patch material include Polytetrafluoroethylene (PTFE) and a running suture technique with a Nonabsorbable suture (prolene) was employed. Exposure for VSD closure was through the right atrium. During operation VSD closure was tested and checked by vulsalva maneuver by anesthetist. patient was gradually weaned off from inotropes and mechanical ventilation and was discharged from hospital after postoperative TTE procedure. A residual VSD measuring about 4 mm was detected with two dimensional echocardiographic imaging. the patient was symptom free at one month and at 3 months follow-up. At last followup the R—L shunt was hemodynamically insignificant, as documented by TTE and electrocardiogram (Neither enlargement of cardiac chambers nor signs of ventricular strain respectively) the patient required no medication and no endocarditis was noted.

Address of Correspondence: Dr. Md Kamrul Hassan, Assistant Professor, Department of Cardiac Surgery, NICVD, Dhaka, Bangladesh.




Fig.-2: Echocardiography showing residual VSD

Discussion:

Haemodynamically significant residual VSD following repair for TOF imposes significant morbidity, mortality and economic burden on the family of the patient.¹ This is specially relevant in developing countries. These defects could be closed by AMVSO (Amplatzer membranous ventricular septal occluder) has been proved to be safe alternative but cost more than repeat redosurgery. The ADO (Amplatzer Duct Occluder) could be an alternative for closure of residual VSD in perimembranous region after surgical repair of tetralogy of Fallot.¹

When a VSD is part of more complex congenital heart disease such as tetralogy of Fallot, generally they are routinely closed with surgical mortalities ranging from 1% to 5%.^{2,4,5}

The focus on post-surgical history of these defects lies more in the long term quality of life and functional status of growing children and young adult, which will be influenced by eventual residual lesions, one of which is a VSD shunt.²

In TOF with small residual defect at the superior aspect of the VSD patch should be followed up closely for the defect enlargement and a late significant shunt.³

Ali Dodge Khatami and colleagues² looked at 7 patients who have residual VSD (2mm in size) in 52 patients of TOF which will have remained open after median follow-up of 3.1 years (Range 0.5–9.7).

Similar to above study and findings Yang and colleagues³ looked at the frequency and significance of residual defects in 294 patients undergoing significant closure of simple VSD and that with associated complex congenital heart disease. They detected a residual VSD in 96 of 294 patients (33%). This is slightly higher in patients with an isolated

VSD (41%) as compared with that of more complex defect with VSD component (33%).

Yang and colleagues recommended residual defect longer than 4 mm should undergo immediate surgical revision, and those with tetralogy of Fallot with a small residual defect at the superior aspect of the VSD patch should be followed closely for defect enlargement and late significant shunt.³

Residual shunt remained in our patient is in the lower aspect of the VSD patch with significant haemodynamic instability. Patient was asymptomatic, medication free and had no episodes of endocarditis at last follow-up, his quality of life was judged to normal.

Bol-Raap and colleagues studied 188 consecutive patients undergoing surgical closure of an isolated VSD among the 73 Residual VSD shunt, 37 disappeared spontaneously at a median time of 3.9 years.⁶

Residual shunt closed spontaneously were frequently after complete repair of AVSD (Atrioventricular septal defect), as compared with after correction of TOF and closure of an isolated VSD.⁵

Conclusion:

Residual shunt greater than 2 mm are unlikely to close spontaneously, but are haemodynamically and clinically irrelevant for the patient. Even after a perfect repair of TOF lifelong endocarditis prophylaxis is indicated.

References:

1. B. Vaidyanathan, BRI Kannan, R. Krishna Kumar. Device closure of residual ventricular septal defect after repair of tetralogy of Fallot using the Amplatzer duct occluder. *Indian Heart J* 2005;57:164-166.
2. Ali Dodge Khatami, W. Kuirisch, M. Tonarke, et al. Spontaneous closure of small residual ventricular septal defects after surgical repair. *Ann. Thorac. Surg* 2007;83:902-906.
3. Yang S.G, Novellor R, Nicolson S et al. Evaluation of ventricular septal defect repair using intraoperative transesophageal echocardiography; frequency and significance of residual defect in infants and children. *Echocardiography* 2000;17:681-684.
4. Hirschfeld, Bwe El. Tetralogy of Fallot. In: Mavondis C, Backer Cl. eds, *Pediatric Cardiac Surgery*, 3rd ed. Philadelphia PA: Mosby:2005: 383-397.
5. Backer CL, Mavondis C. Atrioventricular canal defects. In: Mavondis C, Backer Cl. eds, *Paediatric cardiac surgery*, 3rd ed. Philadelphia PA: Mosby:2005: 321-338.
6. Bol-Raap G, Werheim J, Kappetein AP, Witsenburg M, Boggs AJJC. Follow-up after surgical closure of congenital ventricular septal defect. *Euro. J. Cardio Thorac Surg* 2003; 24: 511-515.