

***BIPOLARIS HAWAIIENSIS* (BUGNICOURT EX M.B. ELLIS) UCHIDA & ARAGAKI - A NEW RECORD OF HUMAN PATHOGENIC SPECIES OF *BIPOLARIS* IN BANGLADESH**

SHAMIM SHAMSI* AND FATEMA YEASMIN

Department of Botany, University of Dhaka, Dhaka- 1000. Bangladesh

Key words: Bipolaris hawaiiensis, New record, Hyphomycetes, Bangladesh

Bipolaris hawaiiensis (Bugnicourt ex M.B. Ellis) Uchida & Aragaki is a human pathogenic fungus under the class Hyphomycetes. It is usually found in plants and soil. The fungus is responsible for allergic bronchopulmonary mycoses. It has been reported to cause subcutaneous infections, nasal pheohyphomycoses, corneal ulcers and allergic fungal sinusitis⁽¹⁻³⁾. Occurrence of *B. hawaiiensis* in Bangladesh has not yet been reported. However, several phytopathogenic species of *Bipolaris* occur in the country. *Bipolaris spicifera* (Bainier) Subrum. was reported on rice, *Bipolaris austaliensis* (M. B. Ellis) Tsuda & Ueyama was isolated from painted wood^(4,5). *Bipolaris oryzae* (Breda de Haan) Shoemaker causes brown spot of rice⁽⁶⁾ and *B. sorokiniana* (Sacc.) Shoem causes Bipolaris leaf blight of wheat⁽⁷⁾ in Bangladesh.

A study was conducted to identify the fungi associated with phylloplane of *Gerbera* spp. belongs to the family Asteraceae. During the period of this study *B. hawaiiensis* was isolated from healthy leaves of *Gerbera aurantiaca* (red flower). Leaf samples of *G. aurantiaca* were collected from Botanic garden, University of Dhaka during the month of February 2012. "Tissue Planting" method was followed for isolation of fungi using PDA medium⁽⁸⁾. *Alternaria citrii* Ellis & Pierce. Apud Pierce, *Aspergillus flavus* Link, *Aspergillus niger* van Teighem, *Aspergillus terreus* Thom, *Cladosporium cladosporioides* (Fresen). De Vries, *Colletotrichum dematium* (Pers. Ex Fr.) Grove, *Colletotrichum dematium* (Pers. Ex Fr.) Grove, and *Pestalotia* de Not were found to be associated with the healthy leaf samples of *G. aurantiaca*. Along with the above fungi an unknown species of *Bipolaris* was isolated from the *Gerbera* leaves. Morphological characters of the fungus were recorded on PDA .

Colonies of the fungus was black velvety. Hyphae pale to mid brown, smooth septate, 1 - 3 μm diameters. Conidiophore was solitary, flexuous, septate, pale to mid brown, up to 120 μm long, but usually much shorter, 2 - 7 μm diameter. Conidia straight, ellipsoidal, oblong, or cylindrical, rounded at the ends, pale to mid brown, 2 - 7 (mostly 5) pseudoseptate, 12 - 37 (24.5) \times 5 - 11 (8.2) μm . Conidia germinate from both poles (bipolar). A flattened hilum or point of attachment is seen on the basal cell of conidia (Figs 1. a-h and 2a-b)).

*Author for correspondence: <Prof.shamsi@gmail.com>.




Fig. 1. *Bipolaris hawaiiensis*: A. 7 days PDA colony. B. 10 days PDA colony. C-H. Conidiophore and conidia. (Bar = 50 μm).

The morphological characters recorded in the present study were compared with those reported^(9,10) and the fungus was identified as *Bipolaris hawaiiensis*. This is the first report on the association of fungi with *Gerbera* plant and occurrence of a human pathogenic species of *Bipolaris* in Bangladesh.




Fig. 2. *Bipolaris hawaiiensis* on *Gerbera auratiaca* (red flower) plant: a. conidiophore and b. Conidia ($\times 2500$)

Bipolaris hawaiiensis (Bugnicourt ex M.B. Ellis) Uchida & Aragaki com. nov. *Specimen examined*: Isolated from healthy leaves of *Gerbera auratiaca*, Botanic Garden, University of Dhaka, Dhaka, S. Shamsi 3061, 10 February 2012.

References

1. Maskin SL, RJ Fetchick, CD Jr Leone, PK Shakey, MG Rinaldi 1989. *Bipolaris hawaiiensis*- caused phaeohyphomycotic orbitopathy. A devastating fungal sinusitis in an apparently immunocompetent host. *Ophthalmology* **96**(2): 175-179.
2. Chowdhury A, HS Randhawa, V Sing, ZU Khan, S Ahmed, S Kathuria, P Roy, G Khanna and J Chandra 2011. *Bipolaris hawaiiensis* as etiologic agent of allergic bronchopulmonary mycoses: first case in paediatric patient. *Med Mycol.* **49**(7): 760-765.
3. Anonymous 2013. URL:<http://Moldlibrary.ca/bopolaris>. Opened on 30 March 2013.
4. Shamsi S 1999. Investigations into the sheath rot disease of rice (*Oryza sativa* L.) in Bangladesh. Ph. D. Thesis. Department of Botany, University of Dhaka, pp. i-xii + 1-127.
5. Shamsi, S. and Z. Yasmin. 2009. *Bipolaris australiensis* (M.B. Ellis) Tsuda & Ueyama - a new dematiaceous hyphomycetes record for Bangladesh. *Bangladesh J. Plant Taxon.* **16** (1): 91-93.

6. Haque SMA, MA Begum, MS Rahman and MAT Mia 2002. Seedborne fungi of Boro rice as affected by farmer's seed processing activities. *Bangladesh J. Plant Pathol.* **18**(1&2):83-89.
7. Hyder, MMA and Fakir GA 1992. Fungi associated with wheat grains in Bangladesh and their pathogenic significance. *Bangladesh J. Bot.* **21**(2): 173-180.
8. Anonymous . 1968. *Plant Pathologist's pocket Book*. Commonwealth Agricultural Bureau (CAB), Commonwealth Mycological Institute, Kew, Surrey, England, pp. 1-267.
9. Shoemaker RA 1959. Nomenclature of *Drechslera* and *Bipolaris* segregated from *Helminthosporium*. *Can. J. Bot.* **37**: 879-887.
10. Unchida JY and M Aragaki. 1979. Etiology of Necrotic Flecks on *Dendrobium* Blossoms. *Phytopathology* **69**: 1115.

(Manuscript received on 23 March, 2013; revised on 27 April, 2013)