

Effects of Ashwagandha (*Withania somnifera*) Root Extract Against Gentamicin Induced Changes of Serum Electrolytes in Rats

Sadia Choudhury Shimmi¹, Nasim Jahan², Nayma Sultana³

Abstract

Background: Regulation of electrolytes and body fluids are essential for maintaining the body homeostasis. Kidney plays an important role for these regulations. Higher doses of drugs, toxins, infectious agents, chemicals etc. can causes kidney damage and ultimately electrolytes disturbances can be occurred. Ashwagandha (*Withania somnifera*) is an herbal plant may have some role on serum electrolytes balance. **Objective:** To observe the effects of ashwagandha (*Withania somnifera*) root on serum electrolytes against gentamicin induced nephrotoxicity in Wistar albino rats. **Methods:** This experimental study was carried out in the Department of Physiology, Sir Salimullah Medical College (SSMC), Dhaka from 1st July 2010 to 30th June 2011. A total number of 35 Wistar albino rats, age from 90 to 120 days, weighing between 150 to 200 grams were selected for the study. After acclimatization for 14 days, they were divided into control group and experimental group. Control group was again subdivided into baseline control, (10 rats) and gentamicin treated control group, (10 rats). Again, experimental group (gentamicin treated group after ashwagandha treatment) consisted of 15 rats. All groups of animals received basal diet for 22 consecutive days. In addition to this, gentamicin treated control group also received gentamicin subcutaneously (100mg /kg body weight/day) for the last eight (15th to 22nd day) consecutive days. Again, gentamicin treated group after ashwagandha treatment received ashwagandha root extract (500mg/kg body weight/day, orally) for 22 consecutive days and gentamicin subcutaneously (100mg/kg body weight /day) for last eight (15th to 22nd day) days. All the animals were sacrificed on 23rd day. Then blood samples were collected and kidney weight was measured. For assessment of kidney function, some serum electrolyte levels e.g. serum sodium, potassium and chloride ion levels were estimated by ion selective electrode (ISE) electrolyte auto analyzer method, by using Biolyte 2000 auto analyzer . However, body weight and kidney weight of the animals were measured to assess the nephrotoxicity in these groups of animals. All these tests were done in the laboratory of Department of Physiology and Biochemistry, SSMC. Statistical analysis was done by one way ANOVA and Bonferroni tests as applicable. **Results:** The serum sodium and chloride ion levels were almost similar in all the groups and the differences were not statistically significant. The mean serum levels of potassium ion were significantly ($p<0.001$) lower in gentamicin treated group and ($p<0.05$) in gentamicin treated group after ashwagandha treatment in comparison to that of baseline control group. But this level of gentamicin treated group after ashwagandha treatment was significantly ($p<0.01$) higher than that of gentamicin treated group. Initial body weight was almost similar and no significant difference of this value was observed among the groups. Whereas, the final body weight was significantly ($p<0.001$) lower in gentamicin treated control group and in gentamicin treated group after ashwagandha treatment than that of baseline control group. Again this level of gentamicin treated group after ashwagandha treatment was significantly ($p<0.05$) higher in comparison to that of gentamicin treated control group. The kidney weight was significantly ($p<0.01$) higher in gentamicin treated control group when compared to that of baseline control and gentamicin treated group after ashwagandha treatment. Whereas, kidney weight of gentamicin treated group after ashwagandha treatment and of baseline control group was almost similar and showed no statistically significant difference of this value between this two groups. **Conclusion:** Ashwagandha (*Withania somnifera*) root extract may have some role in maintaining some of the serum electrolyte levels within normal limit, which indicates its nephroprotective effects against gentamicin induced toxicity.

Key words: Electrolytes, Ashwagandha, Gentamicin

J Bangladesh Soc Physiol. 2012 June; 7(1): 29-35
For Authors Affiliation, see end of text.

Received January 2012; Accepted April 2012

<http://www.banglajol.info/index.php/JBSP>

Introduction

Acids, bases and salts are collectively called electrolytes¹. Electrolytes are involved in many metabolic and homeostatic functions, of our body². Abnormal electrolytes concentrations reflect altered metabolic status³. Electrolyte imbalance can lead to serious consequences as it affects the homeostasis of the body. Commonly measured electrolytes are good indicators of kidney function¹. The kidney is an essential organ plays an important role in regulation of electrolytes and body fluids⁴. The test of electrolytes includes the measurement of sodium, chloride, potassium, bicarbonate ions for both diagnosis and management of renal, endocrine, acid-base, water balance and many other condition⁵. Altered levels of these electrolytes commonly occurs in renal dysfunction and may be life threatening⁶.

The amino glycoside antibiotic gentamicin is widely used in the treatment of infection caused by gram negative bacteria⁷. Drug metabolite may have higher activity and greater toxicity than the original drug. Metabolites of the drugs that are excreted from kidney may also cause cellular damage leading to kidney dysfunction⁸. Gentamicin is a well recognized nephrotoxic drugs produce renal tubular necrosis⁹. A less well documented effect of gentamicin is a disturbance of electrolyte homeostasis.¹⁰ Gentamicin causes electrolytes disturbance particularly serum potassium, sodium, magnesium, calcium etc¹¹.

Ashwagandha (*Withania somnifera*,) belongs to the family of Solanaceae, is the traditional Ayurvedic medicine¹². The roots of ashwagandha contains withaferin A and withanolides, the active ingredients contribute to the most of the biological actions¹³. Roots of this plant may have some therapeutic effects¹². Till today, no side effects have been found in ashwagandha¹⁴.

Again, different researchers from different countries have been studied the effect of ashwagandha (*Withania somnifera*) on electrolytes. Some researchers observed that

ashwagandha significantly decreases the serum electrolytes¹⁵ which may be due to its diuretic effect. Moreover, other investigators found that root of this plant causes non significant change of serum sodium ion but significantly increase serum potassium ion¹⁶.

Renal failure with electrolyte disturbance is potentially serious and causes high morbidity and mortality in our country. In recent years, great effort has been focused on traditional and herbal medicine for the treatment of renal disease along with electrolyte balance. But little is known on this aspect in Bangladesh. Therefore, the present study was carried out to observe the effects of ashwagandha (*Withania somnifera*) root extract on serum electrolyte levels in gentamicin intoxicated rats. It is also expected that the result of this study would make the ashwagandha acceptable among the people for maintaining the electrolyte concentration in the body.

Methods

This experimental study was conducted between July 2010 to June 2011 in the Department of Physiology, SSMC, Mitford, Dhaka. A total number of 35 apparently healthy Wistar albino male rats, weighing between 150 to 200 grams, age from 90 to 120 days were used. The animals were purchased from animal house of Bangladesh Council of Scientific and Industrial Research (BCSIR), Dhaka. Ethical permission was taken from the Institutional Ethics Committee (IEC) of SSMC, Dhaka. Prior conducting the study, the animals were acclimatized for 14 days at $23\pm 2^{\circ}\text{C}$ room temperature under 12 hour dark- light cycle. During this period, they had free access to food and water ad libitum. After 14 days of acclimatization, body weights were measured (initial bw), again, final body weights (final bw) of rats before anaesthetized on 23rd day were taken. The rats were divided into control group and experimental group. Control group was again subdivided into baseline control, consisted of 10 rats) and gentamicin treated control group (consisted of 10 rats). Again, experimental group

(gentamicin treated group after ashwagandha treatment) consisted of 15 rats. All groups of animals received basal diet for 22 consecutive days. In addition to this, gentamicin treated control group also received gentamicin subcutaneously (100mg/kg body weight/day) for the last eight (15th to 22nd day) consecutive days. Again, gentamicin treated group after ashwagandha treatment received ashwagandha root extract (500mg/kg body weight/day, orally) for 22 consecutive days and gentamicin subcutaneously (100mg/kg body weight/day) for last eight (15th to 22nd day) days. After giving gentamicin and ashwagandha all the animals including baseline control rats, were anaesthetized with the help of chloroform and sacrificed on 23rd day. Their blood samples were collected and kidney was removed and was washed in ice cold saline. Then it was wiped in tissue paper, weighed by electric balance analyzer. Blood was centrifuged at the rate of 4000 rpm for 5 minutes and supernatant serum was collected. Out of 35 serum samples, four (4) were hemolyzed. So, thirty one (31) samples were estimated. For assessment of kidney function, some serum electrolyte levels e.g. serum sodium, potassium and chloride ion levels were estimated

by ion selective electrode (ISE) electrolyte auto analyzer method, by using Biolyte 2000 auto analyzer¹⁷. However, body weight and kidney weight of the animals were measured to assess the nephrotoxicity in these groups of animals. All these tests were done in the laboratory of Department of Physiology and Biochemistry, SSMC. Statistical analysis were done by one way ANOVA and Bonferroni test as applicable by using SPSS windows, version 16.

Preparation of root extract of ashwagandha (*Withania somnifera*):

Ashwagandha is cultivated and harvested in the Ayurvedic nursery of Hamdard Laboratories, Meghna. After collecting the roots of this plant, it was dried in sunlight for 2 days, crushed in an electrical grinder into powder. Then the powder was extracted in methanol, filtered, evaporated by rotatory evaporator and dried. After that, the dried root extract of ashwagandha was dissolved by propylene glycol (2ml/kg body weight) and finally mixed with distilled water for feeding.

Results

Body weight and kidney weight are shown in (Table I).

Table I : Body weight and kidney weight in different groups (n=31)

Groups	Body weight (g)		Kidney weight (g)
	Initial (Day 1)	Final (Day 23)	
A ₁ (n=9)	181.13 ± 2.64 (177.12-185.24)	187.18 ± 3.66 (183.10-194.09)	0.57 ± 0.03 (0.54-0.61)
A ₂ (n=9)	182.30 ± 2.80 (178.04-185.92)	176.83 ± 3.69 (169.67-180.28)	1.20 ± 0.14 (1.07-1.38)
B(n=19)	181.85 ± 3.13 (177.10-186.92)	180.46 ± 2.59 (175.92-184.26)	0.58 ± 0.01 (0.55-0.59)
Statistical analysis			
		P value	
A ₁ vs A ₂	0.189 ^{ns}	0.000 ^{***}	0.003 ^{**}
A ₁ vs B	0.285 ^{ns}	0.000 ^{***}	0.188 ^{ns}
A ₂ vs B	0.363 ^{ns}	0.012 [*]	0.007 ^{**}

Results are expressed as Mean ± SD. Statistical analysis was done by ANOVA test and then Bonferroni test was performed to compare between groups. Figures in parentheses indicate maximum & minimum values.

Group A₁=Baseline control group

Group A₂=Gentamicin treated control group

Group B= gentamicin treated group after ashwagandha treatment

n: total number of animal

ns: non-significant * : p < 0.05

** : < 0.01

***: p < 0.001

Initial body weights of all the animals were almost similar and showed no significant difference of this value among the groups. Whereas, the final body weight was significantly ($p < 0.001$) lower in gentamicin treated control group and in gentamicin treated group after ashwagandha treatment than that of baseline control group. Again this level of gentamicin treated group after ashwagandha treatment was significantly ($p < 0.05$) higher in comparison to that of gentamicin treated control group.

The kidney weight was significantly ($p < 0.01$) higher in gentamicin treated control group when compared to that of baseline control and gentamicin treated group after ashwagandha treatment. Whereas, kidney weight of gentamicin treated group after ashwagandha treatment and of baseline control group was almost similar and showed no significant difference of this value between these two groups.

The mean serum sodium and chloride ion levels were slightly higher in gentamicin treated control group and gentamicin treated group after ashwagandha treatment in comparison to those of baseline control group. Whereas, these levels were slightly lower in gentamicin treated group after ashwagandha treatment than those of gentamicin treated control group. Although, these differences were not statistically significant (Figure 1). Again, the mean serum potassium ion level was significantly lower in gentamicin treated control group ($p < 0.001$) and gentamicin treated group after ashwagandha treatment ($p < 0.05$) in comparison to that of baseline control group. Again, this level was significantly ($p < 0.01$) higher in gentamicin treated group after ashwagandha treatment than that of gentamicin treated control group. (Figure 2).




Figure 1: Serum sodium and chloride ion levels in different groups of rats ($n=31$). Group A₁: Baseline control group, Group A₂: Gentamicin treated control, Group B: Gentamicin treated group after ashwagandha treatment




Figure 2: Serum potassium ion level in different groups of rats ($n=31$). Group A₁: Baseline control group, Group A₂: Gentamicin treated control group, Group B (Gentamicin treated group after ashwagandha treatment)

Discussion

In this study lower body weight was found both in gentamicin treated control rats and also in gentamicin treated group after ashwagandha treatment. Again, higher kidney weight was found in gentamicin treated control rats, whereas kidney weight was almost similar in gentamicin treated

group after ashwagandha treatment and baseline control group. This finding is in consistent with that of some other investigators¹².

In the present study, non-significant changes of serum sodium and chloride ion levels were observed among the groups. This finding is in agreement with that of some other researchers^{3,16,18}. Again, some other researchers observed significantly lower level of serum sodium ion in gentamicin treated group after ashwagandha treatment in comparison to that of baseline control group¹⁵. The inconsistent of the result may be due to that ashwagandha was used for 30 days to potentiate diuretic effect in that study in contrast to 22 days administration of ashwagandha in the animals of present study.

Moreover, in this study, hypokalemia was observed in gentamicin treated control group which was improved in gentamicin treated group after ashwagandha treatment towards baseline control. Similar finding was also reported by some other researchers^{9,16,18,19}.

It has been suggested that, toxic dose of gentamicin may decrease the plasma magnesium concentration. This hypomagnesaemia may reduce the body weight²⁰. Again, some other investigators suggested that gentamicin toxicity causes development of uremia, which in turn causes loss of appetite and decrease in body weight¹⁶. Moreover, gentamicin is injurious to renal epithelial cells, thus may decrease water reabsorption and ultimately decrease the body weight²¹.

Again, Gentamicin decreases serum potassium ion level, may be due its stimulatory effect on aldosterone secretion²². Furthermore, glomerular dysfunction and altered tubular reabsorption causes slightly increased level of sodium and decreased level of potassium and magnesium in gentamicin treated rats^{18,23}.

However, some researchers suggested that, presence of active ingredients like withanolides, may be responsible for diuretic activity of

ashwagandha by inhibiting the sodium ion reabsorption in kidney tubules²⁴.

Again, some other researchers suggested that, ashwagandha root contained sitoindosides VII-X and withaferin A, have antioxidant activity by enhancing the free radical scavenging enzymes such as, superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx)¹². Therefore, causes improvement of body weight¹⁶. Again, anti-inflammatory action of its active ingredients may be helpful in decreasing kidney weight¹⁶.

In the present study, nephrotoxicity was observed in rats treated with gentamicin as evidenced by some of the electrolyte imbalance to some extent. This is further supported by an increase in kidney weight in nephrotoxic rats of present study which may be due to increased production of free radicals, which initiate lipid peroxidation and subsequent cellular damage. Significant decrease final body weight of this group of animals is also in favor of this finding.

Again, in this study slightly lower level of serum sodium and chloride ions along with significantly higher level of potassium ion and lower in kidney weight, that comes to almost normal level in the gentamicin treated group after ashwagandha treatment provides an evidence that, the root extract of this ashwagandha plant may have nephroprotective effect against gentamicin toxicity. These effects are most likely due to presence of some active ingredients in ashwagandha root which have antioxidant property. However, the exact mechanism involved cannot be found out from this type of study due to time and financial constrain.

Conclusion

From this study, it can be concluded that ashwagandha (*Withania somnifera*) root extract can maintain some of the serum electrolyte levels to some extent, e.g. serum sodium, potassium and chloride ion levels in gentamicin intoxicated rats, may be due to its antioxidant property. This

effect of ashwagandha root reflects its nephroprotective effects against gentamicin toxicity.

Acknowledgement

Authors of this study are thankful to the authority of Pharmacy Department of Dhaka University, Hamdard Laboratories and BCSIR, Dhaka for the cooperation they provided. I would also express my sincere thanks to the DGHS for providing the partial financial support for completing the research work.

Author Affiliations

1. * Dr. Sadia Choudhury Shimmi, Assistant Professor, Department of Physiology, Delta Medical College and Hospital, Dhaka. E-mail: shimmi_cmc40@yahoo.com
2. Professor Nasim Jahan, professor and Head, Department of Physiology, Sir Salimullah Medical College, Dhaka. E-mail: prof.dr.nasimjahan@gmail.com
3. Nayma Sultana, Associate Professor, Department of Physiology, Sir Salimullah Medical College, Dhaka. Email: nayma_sultana@yahoo.com

**for correspondence*

References

1. Springate JE, Christensen SL, Feld LG. Serum creatinine level and renal function in children. *Am J Dis Child* 1992;146:1232-5.
2. Kraft MD, Btaiche IF, Sacks GS, Kudsk KA. Treatment of electrolyte disorders in adult patients in the intensive care unit. *Am J Health-Syst Pharm* 2005; 62: 1663-1682.
3. Parsons PP, Garland HO, Harpur ES, Old S. Acute gentamicin induced hypercalciuria and hypermagnesiuria in the rat: dose- response relationship and role of renal tubular injury. *Br J Pharmacol* 1997; 122(3): 570-576.
4. Hall JE. Urine formation by the kidneys. In: Guyton and Hall Text Book of Medical Physiology, Philadelphia: Saunders Elsevier; 2011. 307-326 p.
5. Gowda S, Desai PB, Kulkarni SS, Hull VV, Math AAK, Vernekar SN. Markers of renal function tests. *North Amm J Med Sci* 2010; 2:170-173.
6. Langston C. Managing fluid and electrolyte disorders in renal failure. *Vet Clin North Am small anim pract* 2008; 38(3): 677-697.
7. Gingell JC, Water worth PM. Dose of gentamicin in patients with normal renal function and renal impairment. *Brit Med J* 1968; 2: 19-22.
8. Sumathi T, Devaraj SN. Effect of Bacopa monniera on Liver and Kidney toxicity in chronic use of opioids. 2009; Available from: URL: <http://www.thefreelibrary.com/>
9. Stojiljkovic N, Veljkovic S, Mihailovic D, Stojiljkovic M, Radovanovic D, Randelovic P. The effect of calcium channel blocker Verapamil on gentamicin nephrotoxicity in rats. *Bosn J Basic Med Sci* 2008; 8(2): 170-176.
10. Foster JE, Harpur ES, Garland HO. An investigation of the acute effect of gentamicin on the renal handling of electrolytes in rat. *J Pharmacol Exp Ther* 1992; 261(1): 38-43.
11. Chien CL, Chen YS, Shih MT, Lee Ck, Huang HW. Gentamicin induced serum electrolytes abnormalities. *Med Sci* 1984; 5(3): 1997-2002.
12. Mishra LC. Scientific Basis for the therapeutic use of Withania somnifera (Ashwagandha): A Review. *Altern Med Rev* 2000; 5(4): 334-346.
13. Singh G, Sharma PK, Singh S. Biological activities of Withania somnifera. *Ann of Biol res* 2010; 1(3): .56-63.
14. Das K, Samanta TT, Samanta P, Nandi DK. Effect of extract of Withania somnifera on dehydration –induced oxidative stress- related uremia in male rats. *Saudi J of Kidney dis Transpl* 2010; 21(1): 75-80.
15. Andallu B, Radhika B. Hypoglycemic, diuretic and hypocholesterolemic effect of winter cherry (Withania somnifera, Dunal) root. *Indian J Exp Biol* 2000; 38: 607-609.
16. Jeyanthi T, Subramanian P. Nephroprotective effect of Withania somnifera: A dose- dependent study. *Renal failure* 2009; 31(9): 814-821.
17. Eisenman G. Glass Electrodes for hydrogen and other cations. Marcel Dekker. 1967; New York: 235-246.
18. Silan C, Uzun O, Comunoglu NU, Gokcen S, Bedirhan S, Cengiz M. Ameliorated and healing effects of Resveratrol in gentamicin –induced nephrotoxicity in rats. *Biol Pharm Bull* 2007; 30(3): 79-83.

19. Kalaiselvan A, Anand T, Soundararajan M. Renoprotective activity of *Ipomoea digitata* in gentamicin induced kidney dysfunction. *Journal of Ecobiotechnology* 2010; 2(2): 57-62.
20. Whang R. Magnesium deficiency, causes and clinical implications. *Drugs* 1984; 1: 143-150.
21. Muthuraman A, Singla SK, Rana A, Singh A, Sood S. Reno protective Role of Flunarizine (mitochondrial permeability Transition pore Inactivation) against Gentamicin Induced Nephrotoxicity in Rats. *Yakugaku Zasshi* 2011; 131(3): 437-443.
22. Salem CB, Hmouda H, Bouraoui K. Drug induced hypoklaemia. *Current Drug Safety* 2009; 4(1): 55-61.
23. Chou CL, Chen YH, Chau T, Lin, SH. Acquired batter- like syndrome associated with gentamicin administration. *Am J Med Sci* 2005; 329: 144-149.
24. Mirjalili MH, Moyano E, Bonfill M, Cusido RM, Palazón J. Steroidal lactones from *Withania somnifera*, an ancient plant for novel medicine. *Molecules* 2009; 14(7): 2373-2393.