



ORIGINAL ARTICLE

Relationship of Haemoglobin and C Reactive Protein level with Different Types of Heart Failure Patients

Mahboob Morshed¹, Md. Joynul Islam², Fariha Haseen³, Mohammad Ahtashamul Haque⁴, ABM Ruhul Amin⁵

¹Assistant Professor, Department of Cardiology, Shaheed Syed Nazrul islam medical College, Kishoregonj, Bangladesh; ²Associate Professor, Department of Clinical Neurosurgery, National Institute of Neurosciences & Hospital, Dhaka, Bangladesh; ³Associate Professor, Department of Public Health & Informatics, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh; ⁴Assistant Professor, Department of Oral and Maxillofacial Surgery, Dhaka Medical College, Dhaka, Bangladesh; ⁵Senior Consultant (orthopaedics), Jatio Sangsad Medical Centre, Dhaka, Bangladesh

[Received on: 2 February 2019; Reviewed on: 12 April 2019; Accepted on: 1 May 2019; Published on: 1 July 2019]

Abstract

Background: Initial level of haemoglobin and CRP level is important issue among the different type of heart failure patients for their in-hospital outcomes. **Objective:** The purpose of the present study was to see the relationship of haemoglobin (Hb) and CRP level with different types of heart failure patients. **Methodology:** This cross-sectional study was conducted in the Department of Cardiology at Mymensingh Medical College, Mymensingh, Bangladesh from December 2010 to November 2011 for a period of two (2) years. Patients of ACS who were presented within 12 hours of chest pain were included as study population. Study population were categorized in four groups according to the level of hemoglobin and C-reactive protein. Blood sample was collected for baseline laboratory investigations like Troponin-I, random blood sugar (RBS), blood urea, serum creatinine, lipid profile, Hemoglobin and CRP level. Sample were then send to standard laboratory/Biochemistry department of MMCH. **Result:** The mean age of the population was 52.18±8.88 years. Statistically non-significant mean age difference among the groups. Significant difference of mean left ventricular (LV) ejection fraction was found among the groups (p=0.005). Maximum patient in group 1 (48.5%) had ejection fraction in the range of 40 to 50%, compare to group 2(32.0%), group 3 (11.1%) and group 4 (5.8%). The heart failure was statistically significant in Killip class I (p=0.000) and Killip class II (p=0.000). **Conclusion:** In conclusion haemoglobin and CRP level of acute coronary syndrome patients is statistically significantly associated with different types of heart failure. [Journal of Current and Advance Medical Research 2019;6(2): 101-105]

Keywords: Relationship; haemoglobin level; CRP level; Heart Failure

Correspondence: Dr. Mahboob Morshed, Assistant Professor, Department of Cardiology, Shaheed Syed Nazrul islam medical College, Kishoregonj, Bangladesh; Email: mahboob3030@gmail.com; Cell no.: +8801819296183

Cite this article as: Morshed M, Islam MJ, Haseen F, Haque MA, Amin ABMR. Relationship of Haemoglobin and C Reactive Protein level with Different Types of Heart Failure Patients. J Curr Adv Med Res 2019;6(2): 101-105

Funding: This study has been performed without any funding from outside else.

Conflict of Interest: There was no conflict of interest to any of the authors.

Contributions to authors: Morshed M, Islam MJ involved in protocol preparation, data collection; statistical analysis Morshed M; Haseen F, Haque MA and Amin ABMR had involved in manuscript writing & revision of the manuscript.

Copyright: ©2019. Morshed et al. Published by Journal of Current and Advance Medical Research. This article is published under the Creative Commons CC BY-NC License (<https://creativecommons.org/licenses/by-nc/4.0/>). This license permits use, distribution and reproduction in any medium, provided the original work is properly cited, and is not used for commercial purposes.

Introduction

Lower hemoglobin level is associated with higher mortality in part with myocardial infarction¹. Statistically significant increase in mortality was found in patient presenting with hemoglobin concentration lower than 10 gm/dL². Clinical data suggest that C- reactive protein may represent a valuable marker of risk in the short term as well³. There is rapidly growing evidence suggesting that atherosclerosis is in part an inflammatory condition. CRP is a serum marker of inflammation which is easily and inexpensively measured⁴.

Hemoglobin and C-reactive protein can be measured in low cost. In our country there are small number of study with hemoglobin and C - reactive protein. This study may help in management to reduce the morbidity and mortality in ACS. With decline in mortality due to infectious and diarrhoeal disease and death from perinatal cause, cardiovascular diseases particularly CHD will be major health problem and number one killer diseases in developing countries⁵.

Anemia is common in heart failure and is associated with poor outcome a large cohort of community-dwelling patient with CHF anemia is a common and an independent prognostic factor for mortality⁶. Lower level of hemoglobin are associated with higher short term mortality in patient with acute myocardial infarction. Specific therapeutic strategies in anemic patient with MI should be further considered². Anemia causes hypoxia-induced vasodilatation leading to increased sympathetic activity and cardiac output. This mechanism decrease coronary reserve, which is already limited in patients with CAD because of high extraction rate of oxygen in cardiac circulation⁷. Manifestation of myocardial ischemia may occur with only mild anemia. Elevated sympathetic activity was also repeatedly associated with worse outcome after acute myocardial infarction⁸. In the long term there alterations lead to gradual development of cardiac enlargement and left ventricular hypertrophy. This present study was undertaken to see the relationship of haemoglobin and CRP level with different types of heart failure patients.

Methodology

This comparative cross-sectional study was conducted in the Department of Cardiology at Mymensingh Medical College, Mymensingh, Bangladesh from December 2010 to November

2011 for a period of two (02) years. Patients of ACS who were presented within 12 hours of chest pain were included as study population. Patient who were admitted in CCU with clinical features of ischaemic type of chest pain of both sexes within 12 hours of onset of chest pain and diagnosed as ACS were included in this study. Patients with iron therapy before admission, blood transfusion before admission, History of previous or current haemostatic disorder, patient with recent bleeding, renal insufficiency, patient admitted with Chronic disease or inflammatory condition, Patient with malignancy, history of PCI or history of CABG were excluded from this study.

All the data were recorded in a data collection sheet. Study population were categorized in four groups according to the level of hemoglobin and C-reactive protein. Patients who had less than 10 gm/dl haemoglobin level and more than or equal to 12 mg/L of CRP were designated as Group I. Patients who had more than 10 gm/dl haemoglobin level and less than 12 mg/L of CRP were designated as Group II. Patients who had more than 10 gm/dl haemoglobin level and more than or equal to 12 mg/L of CRP were designated as Group III. Patients who had less than 10 gm/dl haemoglobin level and less than 12 mg/L of CRP were designated as Group IV.

Age, sex, cardiovascular risks factor, history, family history of cardiovascular disease, treatment history and ECG were taken during admission. Blood sample was collected for baseline laboratory investigations like Troponin-I, Random Blood Sugar (RBS), Blood urea, Serum creatinine, lipid profile, Hemoglobin & CRP level. Sample were then send to standard laboratory /Biochemistry department of MMCH. The period of follow up was 5 days after admission. Permission from the ethical committee was taken. Continuous data were expressed as mean±SD.

Categorical data were analyzed with chi square test, Student's unpaired 't' test was used for analysis of continuous variables. Comparison between groups was done by unpaired t-test. Multivariate logistic regression analysis was done to determine the association of Hemoglobin & CRP Levels with in hospital outcome in Acute Coronary Syndrome. Avitex CRP (Omega Diagnostics Ltd. Scotland. UK) is a rapid latex agglutination test kit for detection of C-reactive protein in human serum. The detection limit is 6mg/L. Hemoglobin estimation was done by analyzer.

Result

Majority of the study population were in the age group of 41 to 50 Years and 51 to 60 Years of age group which were 73(36.3%) cases in each

followed by 61 to 70 Years which was 33(16.4%) cases. The mean age of the population was 52.18±8.88 years. Statistically non-significant mean age difference among the groups (Table 1).

Table 1: Age distribution of the study population

Age Group	Group Name				Total
	Group 1	Group 2	Group 3	Group 4	
30 to 40 Years	11(14.7%)	6(6.5%)	5(22.7%)	0(0.0%)	22(10.9%)
41 to 50 Years	23(30.7%)	38(41.3%)	7(31.8%)	5(41.7%)	73(36.3%)
51 to 60 Years	28(37.3%)	33(35.9%)	8(36.4%)	4(33.3%)	73(36.3%)
61 to 70 Years	13(17.3%)	15(16.3%)	2(9.1%)	3(25.0%)	33(16.4%)
Total	75(100.0%)	92(100.0%)	22(100.0%)	12(100.0%)	201(100.0%)

Mean±SD = 52.18±8.88; $\chi^2 = 9.56$, P = 0.39 NS; Group 1=Hb < 10 gm/dl CRP ≥ 12 mg/L; Group 2=Hb > 10 gm/dl CRP < 12 mg/L; Group 3=Hb > 10 gm/dl CRP > 12 mg/L; Group4= Hb < 10 gm/dl CRP < 12 mg/L

Significant difference of mean left ventricular (LV) ejection fraction was found among the groups. (p=0.005). Maximum patient in group 1 (48.5%)

had ejection fraction in the range of 40 to 50%, compare to group 2 (32%), group 3 (11.1%) and group 4 (5.8%) (Table 2).

Table 2: Distribution of the study population by Ejection fraction

Ejection fraction	Group Name				Total
	Group 1	Group 2	Group 3	Group 4	
< 40	4 (66.7%)	1(16.7%)	1(16.7%)	0(0.0%)	6(100.0%)
40 to 50	48(48.5%)	32(32.3%)	11(11.1%)	8(5.8%)	99(100.0%)
51 to 60	13(22.4%)	38(65.5%)	5(8.6%)	2(3.4%)	58(100.0%)
>60	5(20.8%)	14(58.3%)	4(16.7%)	1(4.2%)	24(100.0%)
Total	70(37.4%)	85(45.5%)	21(11.2%)	11(5.9%)	187(100.0%)

$X^2 = 23.736$; P = 0.005; Group 1=Hb < 10 gm/dl CRP ≥ 12 mg/L; Group 2=Hb > 10 gm/dl CRP < 12 mg/L; Group 3=Hb > 10 gm/dl CRP > 12 mg/L; Group4= Hb < 10 gm/dl CRP < 12 mg/L

The heart failure was statistically significant in Killip class I (p=0.000) and Killip class II

(p=0.000). Number of Killip class II was more in Group I (38.7%) than other groups (Table 3).

Table 3: Distribution of the study population by types of heart failure

Heart Failure	Group Name				Total	P value
	Group 1	Group 2	Group 3	Group 4		
No HF	26(34.7%)	70(76.1%)	13(59.1%)	8(66.7%)	117(58.2%)	0.357 ^{NS}
Killip I	1(1.3%)	0(0.0%)	0(0.0%)	0(0.0%)	1(0.5%)	0.000 ^S
Killip II	29(38.7%)	14(15.2%)	6(27.3%)	3(25.0%)	52(25.9%)	0.000 ^S
Killip III	12(16.0%)	5(6.5%)	2(9.1%)	1(8.3%)	21(10.4%)	0.99 ^{NS}
Killip IV	7(9.3%)	2(2.2%)	1(4.5%)	0(0.0%)	10(5.0%)	0.92 ^{NS}
Total	75(100.0%)	92(100.0%)	22(100.0%)	12(100.0%)	201(100.0)	

$X^2 = 31.406$; P = 0.002; Group 1=Hb < 10 gm/dl CRP ≥ 12 mg/L; Group 2=Hb > 10 gm/dl CRP < 12 mg/L; Group 3=Hb > 10 gm/dl CRP > 12 mg/L; Group4= Hb < 10 gm/dl CRP < 12 mg/L

Discussion

This prospective observational study was carried out in the department of cardiology, Mymensingh Medical College Hospital, Mymensingh during the period of December 2010 to November 2011. This study was done to find out the correlation of hemoglobin and C-reactive protein with acute coronary syndrome and in hospital outcome.

Among the admitted patient in coronary care unit Mymensingh Medical College Hospital, a total of 201 patient diagnosed as acute coronary syndrome including unstable angina, non-ST elevation myocardial infarction & ST elevation myocardial infarction were included in the study after considering the inclusion and exclusion criteria. The patients were divided in to 4 groups Group-I, Group-II, Group-III and Group-IV according to level of hemoglobin and C-reactive protein level. Among the 201 patient 75 patients were in Group-I (Hemoglobin <10 gm/dl, CRP \geq 12 mg/L), 92 patients in Group-II (Hemoglobin > 10 gm/dl, CRP <12 mg/L) 22 patients in Group-III (Hemoglobin > 10 gm/dl, CRP > 12 mg/L,) 12 patients included in Group-IV (Hemoglobin : <10 gm/dl, CRP < 12 mg/L).

There were no significant difference of age between these groups. The mean age of the patients were 52.18 ± 8.88 years. The highest percentage of Group-I were 51-60 years, and Group-II were 41-50 years, Islam and Majumder¹⁰ found the mean age of the ACS patients were 53.60 ± 8.5 in Bangladeshi population. Lee et al¹¹ found the mean age of ACS patients were 53.2 ± 10.6 years in Bangladeshi population which supported the finding of present study.

In this study maximum number of patients is Group-I had LV ejection fraction in the range of 40-50%, Group-I 48.5%, Group-II 32.3%, Group-III-11.1% and Group-IV 5.8%, respectively which was statistically significant.

Heart failure (any Killip class) was significantly more in Group-I than Group-II, III, IV. Though the incidence of Killip class III and cardiogenic shock/Killip IV were not significant among the groups, but Killip class-II, Killip class- III and Killip class- IV heart failure were more in the patient with low hemoglobin and high CRP level.

Al-Ahmad et al¹² found that highly statistically significant and independent association between low hemoglobin concentration and adverse

cardiovascular outcome among the patient with ACS. Their finding was worsening degrees of anemia were associated with progressively higher rates of hypotension tachycardia and heart failure. Franklin et al¹³ found that the patient group with lower hemoglobin concentration on admission more likely suffered from cardiogenic shock, CCF and post infarction angina than the groups with higher hemoglobin level on admission.

Conclusion

In conclusion haemoglobin and CRP level of acute coronary syndrome patients is statistically significantly associated with different types of heart failure. Patient with ACS has variable levels of hemoglobin and C-reactive protein. It has been found that patient with lower hemoglobin and higher CRP level have adverse in hospital outcome including heart failure and death. So categorizations of patient with ACS on basis of hemoglobin and CRP level may help for risk stratification and management.

References

1. Sarnak MJ, Tighiouart H, Manjunath G, MacLeod B, Griffith J, Salem D, Levey AS. Anemia as a risk factor for cardiovascular disease in The Atherosclerosis Risk in Communities (ARIC) study. *Journal of the American College of Cardiology*. 2002;40(1):27-33
2. Lipšic E, van der Horst IC, Voors AA, van der Meer P, Nijsten MW, van Gilst WH, van Veldhuisen DJ, Zijlstra F. Hemoglobin levels and 30-day mortality in patients after myocardial infarction. *International journal of cardiology*. 2005;100(2):289-92
3. Biasucci LM. CDC/AHA workshop on markers of inflammation and cardiovascular disease: application to clinical and public health practice: clinical use of inflammatory markers in patients with cardiovascular diseases: a background paper. *Circulation*. 2004;110(25):e560-7
4. Franklin K, Goldberg RJ, Spencer F, Klein W, Budaj A, Brieger D, Marre M, Steg PG, Gowda N, Gore JM. Implications of diabetes in patients with acute coronary syndromes: the Global Registry of Acute Coronary Events. *Archives of Internal Medicine*. 2004;164(13):1457-63
5. Wu WC, Rathore SS, Wang Y, Radford MJ, Krumholz HM. Blood transfusion in elderly patients with acute myocardial infarction. *New England Journal of Medicine*. 2001;345(17):1230-6.
6. Ezekowitz JA, McAlister FA, Armstrong PW. Anemia is common in heart failure and is associated with poor outcomes: insights from a cohort of 12 065 patients with new-onset heart failure. *Circulation*. 2003;107(2):223-5.
7. McKechnie RS, Smith D, Montoye C, Kline-Rogers E, O'Donnell MJ, DeFranco AC, Meengs WL, McNamara R, McGinnity JG, Patel K, Share D. Prognostic implication of anemia on in-hospital outcomes after percutaneous coronary intervention. *Circulation*. 2004;110(3):271-7.

8. Al Falluji N, Lawrence-Nelson J, Kostis JB, Lacy CR, Ranjan R, Wilson AC. Effect of anemia on 1-year mortality in patients with acute myocardial infarction. *American heart journal*. 2002;144(4):636-41.
9. Goncalves PA, Ferreira J, Aguiar C, Trabulo M, Silva JA, Seabra-Gomes R. Prognostic value of baseline hemoglobin in acute coronary syndromes. *Circulation* 2002;106(19):402-402
10. Islam AM, Majumder AA. Coronary artery disease in Bangladesh: A review. *Indian heart journal*. 2013;65(4):424-35
11. Lee PC, Kini AS, Ahsan C, Fisher E, Sharma SK. Anemia is an independent predictor of mortality after percutaneous coronary intervention. *Journal of the American College of Cardiology*. 2004;44(3):541-6
12. Al-Ahmad A, Rand WM, Manjunath G, Konstam MA, Salem DN, Levey AS, Sarnak MJ. Reduced kidney function and anemia as risk factors for mortality in patients with left ventricular dysfunction. *Journal of the American College of Cardiology*. 2001;38(4):955-62
13. Franklin K, Goldberg RJ, Spencer F, Klein W, Budaj A, Brieger D, et al. Implications of diabetes in patients with acute coronary syndromes: the Global Registry of Acute Coronary Events. *Archives of Internal Medicine*. 2004;164(13):1457-63.