

ROLE OF *GARDNERELLA VAGINALIS* AMONG SYMPTOMATIC BACTERIAL VAGINOSIS PATIENTS ATTENDING AT CHITTAGONG MEDICAL COLLEGE HOSPITAL

Nura Nasrin Rowshan Ara^{1*} Md Anwar Husain² Md Mostafizur Rahman⁴
Abu Hena Md Saiful Karim Chowdhury³

Abstract

Background : Now a day's bacterial vaginosis is an extremely common health problem for women in the world which causes many complications both in the pregnancy and non-pregnancy states. *G. vaginalis* is most important cause of bacterial vaginosis. **Materials and methods** : A cross sectional study was conducted to assess the role of *Gardnerella vaginalis* as an etiological agent of bacterial vaginosis. This study was undertaken to assess the patients attending at the outpatient department of Gynaecology and Obstetrics of Chittagong Medical College Hospital, Chittagong. A total of 170 sexually active female in the age group of 15-45 years, with abnormal vaginal discharge were selected for the study. A detailed history and a thorough clinical examination of all the cases were done. pH of the vaginal discharge was measured and three high vaginal swabs were collected. Bacterial vaginosis was diagnosed using Amsel's criteria and Nugent's method. *Gardnerella vaginalis* was isolated and identified with their drug sensitivity test by standard methods. **Results** : In this study 38(22.35%) *Gardnerella vaginalis* were isolated by culture and bacterial vaginosis was detected by Amsel clinical criteria (Clinical method) 43(25.30%), Gram

stain Nugent criteria (Gold Standard) 45(26.47%). Our study showed a relatively high prevalence of bacterial vaginosis and high isolation rate of *Gardnerella vaginalis* within the positive bacterial vaginosis patients. **Conclusion** : This result helps proper management and treatment.

Key words

Gardnerella vaginalis, Amsel criteria; Nugent criteria; Human blood bilayer Tween 80 (HBT) agar media; Intermediate flora.

Introduction

Bacterial Vaginosis (BV) is a clinical syndrome characterized by shift of protective resident micro organisms as *Lactobacillus* spp. by opportunistic pathogenic bacteria such as *Gardnerella vaginalis* and other anaerobic bacteria. It is a polymicrobial condition and it involves various organisms such as *Gardnerella vaginalis*, *Mycoplasma hominis*, *Mobiluncus* species, and other anaerobic bacteria, *i.e.*, *Peptostreptococcus* sp., *Prevotella* sp., *Porphyromonas* and *Bacteroids*¹. In most cases of BV, the predominant bacterial species found is *Gardnerella vaginalis*. Historically, *G. vaginalis* was thought to be the sole causative agent of this condition. But its role in the aetiology of BV was downgraded over the years. The biofilm-forming potential and cytotoxic activity of *G. vaginalis* have renewed interest in the virulence of this organism². So bacterial vaginosis is mostly caused by the synergistic interaction of *G. vaginalis* with obligate anaerobes. Recent evidence has once again placed *G. vaginalis* in the spot light and has indicated that *G. vaginalis* is equipped with a number of virulence properties and consequently the idea that it is the aetiological agent of BV is being revisited³.

Bacterial vaginosis is associated with gynecologic complications, such as cervicitis, salpingitis, endometritis, post-operative infections and pelvic inflammatory disease and many obstetric complications, such as premature rupture of the membranes, preterm deliveries, chorioamnionitis

1. Assistant Professor of Microbiology
Army Medical College, Chittagong.
2. Professor of Microbiology (Former)
Chittagong Medical College, Chittagong.
3. Lecturer of Microbiology
Chittagong Medical College, Chittagong.
4. Assistant Professor of ENT
Cox's Bazar Medical College, Cox's Bazar.

***Correspondence:** Dr. Nura Nasrin Rowshan Ara
E-mail: nasrinsumi@yahoo.com
Cell : 01819850606

Received on : 23.10.2017

Accepted on : 12.11.2017

and postpartum endometritis. Despite the fact that Bacterial Vaginosis (BV) is the leading vaginal disorder globally, very little is known about its etiology or pathogenesis.

This study was designed to isolate the causative agent *G. vaginalis* from bacterial vaginosis patients with their antibiotic sensitivity pattern and showed the role of *G. vaginalis* in bacterial vaginosis which would guide clinicians and microbiologists for proper handling of this pathogen & prevent unnecessary use of antibiotics.

Materials and methods

This was a cross-sectional comparative study carried out in the Department of Microbiology, Chittagong Medical College, Chittagong, during the period of July 2011 to June 2012. Approval from ethical review committee of Chittagong Medical College was duly taken. A total of 170 women, 50 pregnant and 120 non-pregnant, in the age group of 15-45 years patients attending the Gynae out-patient department of Chittagong Medical College Hospital was enrolled for this study. The results of the experiments were recorded systematically and Data was statistically analysed for significance of association of *Gardnerella vaginalis* with bacterial vaginosis diagnosed by Amsel criteria and Nugent criteria using Chi-square test. Statistical analysis was done by standard statistical procedure Statistical Package for Social Sciences (SPSS) and $p < 0.05$ was taken as significant.

Inclusion criteria

- i) Women of reproductive (15-45 years) age group.
- ii) Patient having history of abnormal vaginal discharge.
- iii) Patient with or without mild vulver itching or burning.
- iv) Patient included pregnant and non-pregnant women.

Exclusion criteria

- i) Patient below 15 yrs & over 45 yrs.
- ii) Patient associated with known case of malignancy or AIDS.
- iii) History of taking antimicrobial agents or vaginal medication for vaginitis within the last one month.
- iv) Patient having history of vaginal douche on the day of examination.
- v) Menstruating women.

Three vaginal swab samples were collected with all aseptic precaution after taking informed consent from patient or her legal attendant. Samples were collected from each patient by standard technique. First swab sample used for amine test and wet mount preparation. Second swab sample was collected from left lateral vaginal wall for culture of *Gardnerella vaginalis*. Third swab sample collected from vaginal fornix and used for making Gram's stain for Nugent criteria.

Detection of bacterial vaginosis by –

- Amsel criteria
- Nugent criteria

Isolation and Identification of *Gardnerella vaginalis*: Culture in Human blood bilayer Tween 80 (HBT) agar media and other biochemical test.

Procedure of Amsel Criteria: It should require the presence of at least three of the following four criteria.

- i) **Physical Examination of Vaginal Discharge:**
Presence of thin, gray, homogenous, malodorous, adherent vaginal discharge.
- ii) **pH Measurement of Vaginal Fluid:**
In BV patients vaginal fluid with a pH > 4.5 .
- iii) **Whiff Test or Amine Odour Test:**
At first one or two drops of 10% KOH was added in vaginal secretion on a glass slide and smelled for fishy odour (amine odour).
- iv) **Wet Mount Preparation:**
Presence of clue cell on saline wet mount in BV patients.

Procedure of Nugent Criteria: Third swab sample collected from right lateral vaginal wall and was rolled on a glass slides, the smear were air dried and then fixed with methanol for Gram's stain. Then fixed smears were stained Koploff's modification of Gram's for detection of clue cells and evaluation of bacterial morphotypes under light microscope (At x1000) according to scoring system (Score 0-10) of interpretation by Nugent et al.

The amount of each morphotype detected on the smear was graded and allocated a score as below (Table I).

Table I : Nugent criteria Gram stain interpretation (Score 0 to10)

Score	Lactobacillus morphotypes	Gardnerella & Bacteroides morphotypes	Curved Gram-variable rods
0	4+	0	0
1	3+	1+	1+ or 2+
2	2+	2+	3+ or 4+
3	1+	3+	
4	0	4+	

The score of each morphotype is to be added together to get a total score.

Total score = *Lactobacilli* + *G. vaginalis* and *Bacteroides spp.* + curved rods (In each slide).

By using the scoring system, the study cases were grouped into three groups ie, Bacterial Vaginosis (BV) group, intermediate group, normal flora group.

*A slide with a total score of ≥ 7 is interpreted as "BV".

* A slide with a total score of 4 to 6 is interpreted as "intermediate group".

* A slide with a total score of 0 to 3 is interpreted as "normal flora".

Culture for Isolation of Gardnerella vaginalis: The second swab inoculated into a selective and differential Human blood bilayer Tween 80 (HBT) agar media for isolation of *G. vaginalis*.

Procedure of Culture: Collected vaginal swab was inoculated and the plate was placed immediately in the candle extinction jar containing water soaked cotton. All plates are incubated in 5% CO_2 with increased humidity at 37⁰ C for 48 - 72 hrs for primary isolation of *G. vaginalis* and read at 48 hours and rechecked at 72 hours before discarded. The plates were examined by oblique lighting after 24 hrs, 48 hrs, and 72 hrs.

Colonies on HBT agar media were identified as round opaque, smooth colonies that were pinpoint in size after 24 hrs of incubation and 0.5 mm in diameter at 48 hrs, produce β hemolysis after 48 or 72 hrs of incubation.

The β -haemolytic colonies from HBT agar were examined by Gram's staining to see Gram negative coccobacilli. Subcultures were done on Human blood Columbia agar media and Sheep blood agar media by using β -haemolytic colony for pure isolation and to see the haemolytic character. Colonies were also used for catalase test, oxidase test and fermentation of different sugar.

The Identification of Gardnerella vaginalis, Based on

- i. Colonial morphology: Colonies on HBT agar were identified as small white colonies with β -hemolysis after 48 to 72 hours of incubation.
- ii. Clear β -hemolysis with diffuse edges on HBT media, but no hemolysis on sheep blood agar. The zone of hemolysis was 1 to 2 mm wide around the isolated colonies on HBT agar after 48 hours of incubation.
- iii. Gram stained smear from a colony: Gram variable or Gram negative coccobacilli or small rods.
- iv. Catalase and oxidase test negative.
- v. Fermentation of different sugar: Maltose, mannitol, lactose, sucrose.
- vi. Susceptibility to different antimicrobial agents.

Antimicrobial Susceptibility: All the isolates of *G. vaginalis* obtained by culture were tested for antimicrobial susceptibility by the single disc diffusion method against different antimicrobial agents. The organisms were tested against Metronidazole (MTZ) Clindamycine (CD) Ampicillin (AMP) Ceftriaxone (CRO) Erythromycin (E) Ciprofloxacin (CIP) Vancomycin (VA) Cotrimoxazole (SXT) Chloramphenicol (C) and Tetracycline (TE).

Results

A total of 170 clinically suspected cases of Bacterial Vaginosis (BV) aged between 15-45 years were included in this study. Among the study cases, 120 (70.59%) were non-pregnant and 50 (29.41%) were pregnant with gestational age ranging from 6 to 32 weeks.

Table I : Distribution of study population on the basis of Amsel criteria

Pregnancy Status	Total	Amsel Criteria	
		Bacterial Vaginosis	Other than B V
Pregnant	n=50	13 (26.00)	37 (74.00)
Non-pregnant	n=120	30 (25.00)	90 (75.00)
Total	170(100.00)	43 (25.30)	127 (74.70)

- Figures within parentheses indicate percentage

Table II : Frequency distribution of Amsel's criteria

Diagnostic criteria (Amsel's)	Women with bacterial vaginosis (n=43)	Women without bacterial vaginosis (n=127)	χ^2 test Significance
Homogeneous vaginal discharge	Present	17	p = 0.000 Highly significant
	Absent	110	
pH	≥4.5	40	25 p = 0.000 Highly significant
	<4.5	03	
Amine test	Positive	15	p = 0.000 Highly significant
	Negative	112	
Clue cells	Present	05	p = 0.000 Highly significant
	Absent	122	




Fig 1 : Distribution of study population on the basis of Nugent criteria. (Bar chart showing distribution of Nugent criteria)




Fig 2 : Distribution of study population on the basis of culture of *G. vaginalis* (Pie Chart : Distribution of isolated *G. vaginalis* result)

Table III : Association of Amsel clinical criteria and culture of *G. vaginalis* (n=170)

Culture of <i>G. vaginalis</i>	Amsel clinical criteria		Total
	Positive	Negative	
Positive	38 (22.35)	00 (0.00)	38 (22.35)
Negative	05 (2.95)	127 (74.70)	132 (77.65)
Total	43(25.30)	127 (74.70)	170 (100.00)

- Figures within parentheses indicate percentages
- $\chi^2 = 139.495$, p = 0.000, Highly significant (p < 0.001)

Table IV : Association of culture of *G. vaginalis* and Nugent criteria (n=170)

Nugent Criteria	Culture of <i>G. vaginalis</i>		Total
	Positive	Negative	
Bacterial vaginosis	38 (22.35)	07 (4.12)	45 (26.47)
Intermediate	00 (00)	58 (34.12)	58 (34.12)
Normal Flora	00 (0.0)	67 (39.41)	67 (39.41)
Total	38 (22.35)	132 (77.65)	170 (100.00)

- Figures within parentheses indicate percentages
- $\chi^2 = 154.869$, p = 0.000, Highly significant (p < 0.001)

Table I : On the basis of Amsel criteria (Clinical criteria) 43(25.30%) cases were Bacterial vaginosis (BV) positive and 127(74.70%) BV negative.

Table II: Shows frequency of component of Amsel criteria. Here within 170 patients 58 cases presented homogenous vaginal discharge, 65 cases showed pH 4.5, amine test positive 57 cases and clue cell present only 48 cases. Within these 43(25.30%) cases are BV positive, because we included the presence of at least three of the four criteria. But we included clue cell must. p<0.0001 was taken as significant.

Fig 1: On the basis of Nugent criteria by Gram-staining, study cases were categorized into three groups, Positive bacterial vaginosis (BV) were 45(26.47%) intermediate group were 58 (34.12%) and normal flora were 67(39.41%).

Fig 2: Shows that culture of vaginal fluid yielded growth of *G. vaginalis*. In which 38(22.35%) cases of *G. vaginalis* were isolated and 132(77.65%) cases were culture negative.

Table III shows the association of culture of *G. vaginalis* with Amsel criteria. The Amsel criteria were positive in all 38(22.35%) culture-positive cases. Additional 05(2.95%) cases were positive among culture-negative cases. The difference was highly significant ($p < 0.001$) when Amsel criteria was compared with culture.

Table IV shows the association of culture of *G. vaginalis* with Nugent criteria (Gram stain). Culture was positive in all 38(22.35%) cases out of 45(26.47%) BV positive cases by Nugent criteria. No more cases were positive in intermediate group and no case in normal flora group of Nugent criteria. The difference was highly significant ($p < 0.001$), when culture of *G. vaginalis* and Nugent criteria were compared (Here intermediate group and normal flora group were considered as negative).

Discussion

Bacterial vaginosis is considered as a common vaginal disorder in women of reproductive age. The interest in bacterial vaginosis has increased lately because of the evidence of adverse sequel to this disorder, such as amniotic fluid infection, clinical chorioamnionitis, Premature Rupture of Membranes (PROM) preterm delivery, low birth weight and postpartum endometritis. Non-pregnant women with bacterial vaginosis have been reported to get post-abortion pelvic inflammatory disease, post hysterectomy vaginal cuff cellulitis and plasma cell endometritis. Several publications have also reported an altered vaginal micro flora being linked to an increased susceptibility to the acquisition of HIV and other sexually transmitted infectious agents such as *Neisseria gonorrhoeae* and *Chlamydia trachomatis*^{4,5}.

In this study out of 170 study cases, 43(25.30%) cases were diagnosed as bacterial vaginosis by Amsel criteria, 45(26.47%) cases were positive by Nugent criteria.

On the basis of Amsel clinical criteria, a total of 43(25.30%) cases had been identified to have BV which is slightly lower than that of Navarrate P et al., Rangari et al. and Neelam et al. who reported 31.1%, 58% and 38.55% cases of BV respectively^{6,7,8}. This slightly lower incidence in our study may be due to mandatory inclusion of clue cells on saline wet mount as a marker of BV

for every case, which makes the Amsel criteria more specific. Higher prevalence rates have been reported in previous studies too^{9,10}. Factors responsible for higher prevalence of bacterial vaginosis among the study population were lower socio-economic status, improper sanitation, poor hygiene, malnutrition¹¹.

Among the individual criteria used to diagnose bacterial vaginosis, raised pH is recognized as the most sensitive but least specific criteria¹⁰. In the present study, the pH of the vaginal fluid was also found to be significantly associated with bacterial vaginosis. Majority of the patients with bacterial vaginosis had a pH between 5.0-5.5. Amsel et al also found a pH of more than 4.5 in 81 % cases of bacterial vaginosis¹². Errors in pH measurement may be made by sampling cervical mucus rather than vaginal discharge which has a higher pH or due to presence of cervical infection which increases the pH by increasing the flow of cervical secretions into the vaginal canal¹¹. Amine test is both highly sensitive and specific. Association between amine test and bacterial vaginosis was found to be statistically significant in this study. Detection of amine odour is observer dependant with wide person to person variability. The amine test is easily performed, rapid, inexpensive diagnostic test with good sensitivity and specificity which, as suggested by previous studies, is ideally suited to clinical setting where microscopy is not available¹³.

Significant association was found between clue cells and bacterial vaginosis which was in confirmation with earlier studies¹¹. According to a previous study, the sensitivity and specificity of clue cells on wet mount for diagnosis of bacterial vaginosis is 81% and 99%. However, recognition of clue cells in wet mount which is an excellent denominator of bacterial vaginosis is subjected to variability, depending on the quality of microscope, the adequacy of specimen and the skill of observer¹¹. In our study, among BV cases diagnosed by Amsel criteria, 100% had clue cells positive on vaginal wet smear. That's why we have got the accurate results but the incidence of BV was slightly low.

In our study, the study cases were categorized into three groups according to Nugent criteria. Out of 170 study cases, 45 (26.47%) cases were diagnosed as BV, 58 (34.12%) cases as intermediate

group and 67 (39.41%) cases as normal flora group. The Nugent criteria with mandatory inclusion of clue cells in Gram's smear make the diagnosis reliable and specific. Nugent criteria with inclusion of clue cells had been able to identify 45(26.47%) BV patients. This is slightly higher than that of Udayalaxmi et al and Devi et al who reported 19% and 20.5% in India^{14,15}.

According to Rosenstein et al, the intermediate stage is considered a transitional phase and the patients may go on to frank bacterial vaginosis¹⁶. Gram staining of vaginal secretions is more reliable with sensitivity of 89-93% and specificity of 70-83%. This technique is least expensive, requires the least time to perform, is more widely available than other laboratory methods and is the most interpretative of the laboratory methods¹⁷.

In this study vaginal specimen from study cases were subjected to culture in Human Blood Bilayer Tween (HBT) agar media a highly selective media, yielded growth of *G. vaginalis* from 22.35% of study cases. The isolation was higher than that of Devi et al and Udayalaxmi in India who reported 17.42% and 16.7% respectively^{14,15}. Slightly higher rate reported by Gupta et al and Nahar et al might be due to the use of three or more media that were either non selective or enriched for primary isolation of *G. vaginalis* and variable methods for their identification^{18,19}.

In our study the antibiotic sensitivity pattern of *G. vaginalis* showed extreme variation. Gardnerella vaginalis showed high (52.63%) resistant to most commonly used metronidazole and 100% sensitive to clindamycin.

The association between incidences (22.35%) *G. vaginalis* by culture with bacterial vaginosis by amsel criteria (25.30%) and Nugent criteria (26.47%) are highly significant. Begum et al and Akhter et al from BSSMU in Bangladesh reported similar findings and association^{17,20}. But different study showed very low isolation rate of *G. vaginalis* 6% and 10.2%. Those studies showed slightly association between Gardnerella vaginalis with bacterial vaginosis. But Gupta et al in India, Nahar et al in Bangladesh and Pheirfer et al in UK who reported 54.1%, 38.98% and 91.44% bacterial vaginosis respectively^{18,19,21}. They isolated 50%-90% *G. vaginalis* within positive bacterial vaginosis. They showed *G. vaginalis* was

the main causative organism of bacterial vaginosis. In our study positive bacterial vaginosis were 43(25.30%) or 45(26.47%). Within this 38(22.35%) cases were *G. vaginalis*. Only 5/7 cases were *G. vaginalis* negative. This is the great association between *G. vaginalis* with bacterial vaginosis. This study showed *G. vaginalis* had significant role in the bacterial vaginosis.

Conclusion

Early detection of causative agent and treatment of bacterial vaginosis appear to have a role in reducing the complications associated with this infection. Hence, it may be important to explore primary preventive strategies which target the risk factors or behaviours for bacterial vaginosis. Our study showed *G. vaginalis* is the principle causative agent of bacterial vaginosis. Though *G. vaginalis* is very fastidious bacteria and our laboratory setting is limited, it needs further trial for diagnosis and evaluation in clinical and laboratory settings. So detection of causative agent of bacterial vaginosis and proper antibiotic treatment will prevent recurrent bacterial vaginosis which will reduce drug resistance.

Disclosure

All the authors declared no competing interest.

References

1. Sobel Jack D. Bacterial vaginosis. Annu Rev M. 2000; 51:349-356.
2. Patterson JL, Stull-lane A, Girerd PH & Jefferson KK. Analysis of adherence, biofilm formation and cytotoxicity suggests a greater virulence potential of Gardnerella vaginalis relative to other bacterial-vaginosis-associated anaerobes. Microbiology.2010; 156(2): 392-399.
3. Gelber SE, Aguilar JL, Lewis KL & Ratner AJ. Functional and phylogenetic characterization of vaginolysin, the human-specific cytolysin from Gardnerella vaginalis. J Bacteriol.2008; 190: 3896-3903.
4. Eschenbach DA. Bacterial vaginosis and anaerobes in obstetric-gynecologic infection. Clin Infect Dis. 1993; 16: 282-287.
5. Joesoef MR, Wiknjastro G, Norojono W, Sumampouw H, Linnan M, Hansell MJ. Coinfection with chlamydia and gonorrhoea among pregnant women with bacterial vaginosis. Int J STD AIDS. 1996; 7:61-64.

6. Navarrete P, Dominguez M, Castro E & Zemelman R. Evaluation of Nugent and Amsel criteria for the diagnosis of bacterial vaginosis. *Rev Med Chil.* 2000; 128(7):767-771.
7. Rangari AA, Singh P & Sharma VK. Comparison of the Amsel's composite clinical criteria and Nugent's criteria for diagnosis of Bacterial vaginosis: A step towards preventing mis-diagnosis. *J of Advance Researches in Biological Science.* 2013; 5(1):37-44.
8. Neelam S, Sohail I. Rapid Clinical Diagnostic Tests for Bacterial Vaginosis and its Predictive Value. *Int J Pathol.* 2010;8:50-52.
9. Bhalla P, Kaushika A. Epidemiological and microbiological correlates of bacterial vaginosis. *Indian J Dermatol Venereol Leprol.* 1994; 60: 8-14.
10. Mohadani JW, Dekate RR, Shrikhande AV. Cytodiagnosis of discharge per vaginum. *Indian J Pathol Microbiol.* 1998; 41:403-411.
11. Baruah FK, Sharma A, Das C, Hazarika NK, Hussain JH. Role of Gardnerella vaginalis as an etiological agent of bacterial vaginosis. *IRAN. J. MICROBIOL.* 2014; 6(6)409-414.
12. Amsel R, Totten PA, Spiegel CA, Eschenbach DA, Holmes KK. Nonspecific vaginitis: Diagnostic criteria and microbial and epidemiologic associations. *Am J Me.* 1983; 74:14-22.
13. Hay PE. Recurrent bacterial vaginosis. *Dermatol Clin.* 1998; 16:769-773.
14. Udayalaxmi G, Subbannayya K & Shalini S. Comparison of the methods of bacterial diagnosis of bacterial vaginosis. *J of Clinical and Diagnostic Research.* 2011; 5: 498-501.
15. Devi PS, Rao Ashriyan, M Rajaram, K & Jagadish C. Diagnosis of Bacterial vaginosis in a rural setup. *Indian J of Medical Microbiology.* 2004; 22(1): 47-50.
16. Rosenstein IJ, Morgan DJ, Sheehan M, Lamont RF, Taylor RD. Bacterial vaginosis in pregnancy: Distribution of bacterial species in different Gram stain categories of the vaginal flora. *J Med Microbiol.* 1996; 45: 120-126.
17. Begum N, Muazzam N, Shamsuzzaman SM, Islam MDU, Chowdhury AK & Rashid A. Diagnosis of bacterial vaginosis by acridine orange staining and its comparison to conventional methods and association of Gardnerella vaginalis with bacterial vaginosis. *Bangladesh J Med Microbiol.* 2010; 04(01):37-42.
18. Gupta BK, Kumar R, Sofat R, Khurana S & Deepinder. The role of Gardnerella vaginalis in non-specific vaginitis in intrauterine contraceptive device user. *Indian J of pathology and microbiology.* 1998; 4(1): 67-70.
19. Nahar A, Rahman KM, Hassan MM & Haq JA 1994. 'Studied on the anaerobic vaginosis among the Bangladeshi women of reproductive age group,' *Bangladesh Med Res Counc Bull .* 1994; 20: 129-138.
20. Akhter S, Satter H, Tarafder S, Miah RA, Sharmin S, Ahmed S. *Bangladesh J Med Microbiol.* 2010; 04 (01): 24-27.
21. Pheifer TA, Forsyth PS, Durfee Ma, Pollock HM & Holmes KK. Non specific vaginitis, role of Haemophilus vaginalis and treatment with metronidazole. *N Engl J Med.* 1978; 298: 1429-1434.