

Prevalence of Childhood Obesity in Dhaka City: Identifying Risk Factors

Khan MAI¹, M A Muhit MA², Silvana K³, Afrooz F⁴, Mahamuduzzaman ASM⁵

Conflict of Interest: None

Received: 19.12.2023

Accepted: 10.01.2024

www.banglajol.info/index.php/JSSMC

Key Words:

childhood obesity, risk factors dietary habits, physical activities and life style risk factors.

ABSTRACT:

The human race is facing a novel and enormous health challenge due to the rapidly unfolding global epidemic of obesity. Childhood obesity continues to increase significantly throughout the world with possible future consequences of developing insulin resistance, dyslipidemia, hypertension, metabolic syndrome, impaired glucose tolerance or Type 2 diabetes etc. To fight obesity, it is most important to identify the risk factors of childhood obesity. Therefore, it was planned to conduct this study, to explore the prevalence of childhood obesity in the studied population and to identify the risk factors contributing to the problem among these obese children that may help to design effective local prevention efforts and to develop recommendations for the prevention and management of obesity for a developing country like Bangladesh for a healthy tomorrow.

[J Shaheed Suhrawardy Med Coll 2024; 16(1): 22-27]

DOI: <https://doi.org/10.3329/jssmc.v16i1.85261>

Authors:

1. Dr. Md. Aynul Islam Khan; Associate Professor, Department of Paediatrics, Shaheed Suhrawardy Medical College, Dhaka.
2. Prof. Dr. M. A. Muhit; Public health and disability expert, University of South Asia, President, CSF Global.
3. Dr. Kiswar Silvana; Assistant Professor, Department of Paediatrics, Manikgonj Medical College, Manikgonj.
4. Dr. Farzana Afrooz; Assistant Professor, Department of Paediatrics, Shaheed Suhrawardy Medical College, Dhaka.
5. Dr. Abu Shoyeb Md. Mahamuduzzaman; Assistant Professor, Department of Paediatrics, Shaheed Suhrawardy Medical College, Dhaka.

Correspondence:

Dr. Md. Aynul Islam Khan; Associate Professor & Head of department, Department of Paediatrics. Shaheed Suhrawardy Medical college, Dhaka.

Mobile: 01716665566; Email: aynulislamkhan@yahoo.com

Introduction:

The human race is facing a novel and enormous health challenge due to the rapidly unfolding global epidemic of obesity. The reasons behind this may include changes in the nature of our diet, transformation of agricultural production, the restructuring of the food supply and distribution mechanisms, urbanization and the development of sedentary modes of work and leisure. (James et al. 2006). Childhood obesity causes "bow-legged" children, pseudo tumor cerebri, asthma, obstructive sleep apnea syndrome, gallstones and fatty liver, insulin resistance, metabolic syndrome, hyperandrogenemia, impaired glucose tolerance, Type 2 diabetes, abnormal menstrual cycle, hirsutism, acne, acanthosis nigricans etc

are frequently seen in obese children. (Must and Strauss, 1999). Socio-economic and socio-demographic risk-factors include family size, birth order, family income, family expenditure, care-giving etc. are responsible for the development of childhood obesity (Ravelli et al 1979). There is no study on analysis of risk factors for development of obesity in our population. Thus, this study was designed to focus on a selected population of school children to identify the prevalence of obesity among them and try to find out the presence of some selected risk factors for the development of obesity. The identification of these risk factors is the key for prevention of childhood obesity. Therefore, this study was conducted to find out the prevalence of childhood obesity

in the studied population and to identify the risk factors contributing to create this problem among these obese children and also help to design effective local prevention efforts and to develop recommendations for the prevention and management of childhood obesity for a developing country like Bangladesh for a healthy tomorrow.

Methodology

Materials and Methods

A case-control study was conducted at eight (8) schools in Dhaka city in collaboration with the School of Public Health & Life Sciences, University of South Asia, Banani, Dhaka during the period of January, 2012 to December 31, 2012 (12 months).

Study population and Sampling technique

All school going children aged 6-9 yrs in 8 purposively selected schools (equal no of Bengali-medium & English medium schools) of Dhaka city were selected. Written consent were obtained from school authorities initially and then from the students or their guardians to conduct the study. Height and weight of all children in this age group were measured. Thereafter, the BMI were calculated for age and sex of all students as per CDC growth chart. Children whose BMI for age were $\geq$ 95th percentile of CDC growth chart were selected as obese. Finally, 24 obese students from each school were selected by lottery method from four age groups. From each age group equal no. of boys and girls were enrolled. Thereafter, for each obese child one non-obese age and sex matched child was also selected. In total, 384 students (8 X 24 192 obese and 192 non-obese) were selected for the study purpose.

Data collection

Data regarding different risk factors were collected from the participants using structured questionnaire by the researcher. The risk factors were classified into 3 groups - dietary habits, physical activities and life style risk factors. Information on socio-economic and socio-demographic status like position of the index case, family members, family structure, income level, expenditure on food and educational status of parents were also included in the questionnaire.

Data analysis plan

Data were analyzed by computer based program SPSS 12.0 version. Confidence interval was set at 95% level. P value <0.05 was considered as significant. Chi-square test was done for making comparison between qualitative data. Odds ratios were calculated manually to see the associations between the childhood obesity and risk factors.

Ethics

The Ethical Committee of University of South Asia, approved the protocol. Informed written consent was taken from all the mothers/parents/guardians after full explanations of the nature and purpose of the procedure used for the study. Anonymity was maintained throughout the study, and none of the names were used in the database.

Result

Total 4,527 students were screened (English medium school = 2183 and Bangla medium school 2369), out of them 485 students were obese (10.7%). The bivariate analysis among Bengali and English medium obese and non-obese children were found significant different ($p < 0.00001$)

Table 1: Distribution of obese child according to their medium of schools (4-9 yrs)

	Bengali	English	Total
Obese	198(8.4%)	287 (13.2%)	485 (10.7%)
Non Obese	2171	1896	4067
Total	2369	2183	4552

df-1, Chi-square (χ^2) Test; p value < 0.00001 .

No significant differences were found between the number of issue of a family, family structure, birth order, father's and mother's educational level, father's and mother's occupation with childhood obesity. Family income per month and expenditure of money on food per month were significantly related with the development of childhood obesity ($P < 0.00001$) Majority of the mother of both groups were the main care giver and no added risk was detected in relation to care giver or working mother.

Table 2: Distribution of monthly family income & expenditure on food of cases and controls:

Monthly family income (Tk.)	Case (n=192)	Control (n=192)	p-value
≥25,001-50,000	4	41	
50,001-80,000	56	122	<0.00001
80,001 and above	132	29	<0.00001

Monthly family expenditure on food (Tk.)	Case (n=192)	Control (n=192)	p-value
≥25,001-50,000	4	41	
50,001-80,000	56	122	<0.00001
80,001 and above	132	29	<0.00001

Physical inactivity like watching TV, Video game and computer watching and also the availability of playing ground at school and at or near home were significantly ($P < 0.00001$) related with the development of childhood obesity. Watching some socio-risk factors were analyzed. Family income >80,000/- per month and spending money >50,000/- per month on food were associated with the development of childhood obesity with Odds Ratio of 12.36 and 6.42 respectively.

TV watching for more than 4 hrs and time spend for Video game or computer for more than were found to be risk in development of obesity with the OR of 6.1 and 4.96 respectively. Physical activity more than 2 hrs, presence of play ground at or near home and school were negatively related to the development of childhood obesity with OR of .2, 0.09 and .75 respectively.

Table 3: Socio-demographic risk factors for development of childhood obesity

Socio-demographic Risk-factors	Case (192) (n = 96)	Case (192) (n = 96)	Odds ratio
Family income >50,000/- taka	132	29	12.36
Expenditure on food >30,000/- taka	108	32	6.42

Table 4: Activity-related risk factors for development of obesity in studied children.

Activity related risk factors	Case (n=192)	Control (n=192)	Odds ratio
Watching TV < 4 hrs	94	164	0.09
Watching TV 4-8 hrs	98	28	6.1
Watching Videogame and computer < 2 hrs	45	58	0.70
Watching Videogame and computer 2-4 hrs	84	26	4.96
Physical activity <2 hr	145	60	6.78
Physical activity ≥ 2 - 4 hr	52	125	0.20
Relationship with Play ground at or near home	55	135	0.09
Relationship with Play ground at School	135	145	0.75

Dietary risk factors were analyzed for the development of childhood obesity considering the amount of diet and the frequency of intake of foods were analyzed. Multiple food habits were considered like amount of rice, no. of bread, the method of egg preparation (boiled/ fried), no. of egg intake daily/weekly, vegetables, fish, mutton, chicken and chicken roast and some fast food items etc to see whether these items were responsible for the development of childhood obesity. Taking more rice (>2.5 plate), preparation of egg (boiled/ fried), increased frequency of taking egg were found to be risk factors for the development of obesity with ORs of 2.04, 2.01 and 4.8 respectively. Excess intake of mutton and chicken (three times/weekly) and noodles (daily) were found to be risk for the development of childhood obesity with odds ratios were 3.2, 2.37 and 4.9 respectively. Odds ratios were not significant in case of fruits (OR 0.95) and vegetables (OR 0.58). Intake of fish (three times/weekly) was found healthy for the children and odds ratio was 1.2 and daily intake was found to be negatively related with the development of childhood obesity with OR of 0.56, but on the contrary if the intake decreased to once weekly, the chance of development of obesity was more (3.19). The intake of fast foods like chips (OR 6.44), burger (OR 7.1), pizza (OR 6.5), chicken roast (OR 5.5), ice cream (OR 4.3) were significantly working as risk factors in the development of childhood obesity.

Table 5: Dietary habit - related risk factors for development of childhood obesity

Risk factors		Case (192)	Contro (192)I	Odds Ratio
Considering the amount of Rice	Two and Half Plate	130	58	2.04
Considering the preparation of Egg	Omelet/Fried egg	140	110	2.01
Considering the frequency of taking egg only	Once daily	130	58	4.8
	3 times weekly	62	76	0.72
Relationship with Vegetables intake	Once daily	40	60	0.58
	3 times weekly	96	96	1
Relationship with Fruit intake	Once daily	45	90	0.27
	3 times weekly	115	120	0.95
Considering the frequency of taking Mutton	3 times weekly	120	60	3.2
	Once weekly	54	65	0.73
Considering the frequency of taking Chicken only	3 times weekly	130	95	2.37
	Once weekly	44	130	0.16
Considering the frequency of taking Noodles	Once daily	72	21	4.9
	3 times weekly	102	75	1.7
Considering the frequency of taking Fish Only	Once daily	88	115	0.56
	3 times weekly	52	45	1.2
	Once weekly	52	20	3.19
Considering the frequency of taking Chips only	Once daily	92	24	6.44
	3 times weekly	80	84	0.91
Considering the frequency of taking Burger only	3 times weekly	92	22	7.1
	Once weekly	52	30	2.05
Considering the frequency of taking Pizza only	Once weekly	18	3	6.5
	Once fortnightly	39	8	5.86
Considering the frequency of taking Fried Chicken only	3 times weekly	45	10	5.5
	Once weekly	64	35	2.16
Considering the frequency of taking Ice Cream only	3 times weekly	27	07	4.3
	Once weekly	55	40	1.5

Discussion:

This study was conducted in 8 schools of Dhaka city among 6-9 years of age group of children and the schools were purposively selected. The prevalence of obesity was found 10.7%. Obesity was found more among the English medium school boys of 8-9 yrs of age group. Some studies were done in India, Umesh et al. (2002) and Kaushik et al. (2007) showed that obesity was found 6%-8% and 5.10% respectively in Kolkata, but Anoop Misra et al. (2008) showed in his study that in urban area in New Delhi obesity was increasing from 16% in 2002 to 24% in 2006-2007. There was a study which was done in Dhaka city in 2000-2001 by Rahman et al. where obesity was found 7.6%. Our study is showing the increasing trend of childhood obesity in Dhaka city but did not matched with any of the Indian studies because of

the socio-cultural status, economical status and geographical variations etc as well as year of study time.

A study was conducted by Rahman et al. in 2000-2001 in Dhaka city where childhood obesity was found 7.6% among the studied population which became one and half time more in our study in last one decade which matched with the findings in developed country like USA and Australia. The prevalence of obesity quadrupled over 25 years among the children of USA (Ogden et al. 2002) and in Australia, obesity was increased by 4.6 fold among girls and 3.4 folds among boys (Magarey, 2001).

Socio-demographic risk factors like family size, family structure, birth order, care giver, family income and expenditure on food etc were analyzed. Wilkinson et al. 1997 showed in his study that the prevalence of obesity was high among the nuclear family and in single children family and the prevalence declined with increasing the size of the family but our study failed to show any significant association with childhood obesity. This study was done in urban area and where the family size was usually small with only one or two children and also the unavailability of joint family in urban area showed no

significant associations with childhood obesity which was consistent with that of Revali et al. study (1979).

This study was performed only in urban area with majority of the parents were literate and educated, so, their educational status had no significant association with the development of childhood obesity. So, it was not possible to see the association of childhood obesity with uneducated parents.

Socio-demographic risk factors like family income and family expenditure on food were analyzed to see the association with childhood obesity. World Health Organization (1998) considered income, expenditure, education, occupation and place of residence (urban/rural) to

evaluate socio-economic status in relation to obesity. Our study showed that 68.75% of children were from high income family in obese group. High family income (>80,001/-per month) with OR of 12.36 and spending on foods (50,001/- per month) with OR of 6.02 were associated with the risk of development of childhood obesity.

Children's daily life is full of activities and movements, for example, walking, sports, gardening, running or jogging, household chores, active games and play, and stretching. But due to urbanization, life style modification and digitalization of world, most of the children are becoming habituated to sedentary behavior like watching TV and Videos and using computers etc. Absence of these daily activities they are becoming obese. Also due to lack of playing and outdoor games facilities, children are habituated to sedentary activities and become obese. We observed in our study that in most of the schools had small playing ground and in some schools there was no playing ground and interestingly children were not allowed to play in these playing ground except weekly for 30-45 minutes. No significant association was found between schools having play ground and childhood obesity. On the contrary, who had play ground at or near home, were less prone to develop obesity. Timperio et al. (2004) showed in his study that without safe places near home might keep their children almost inactive with indoor games and allow them to develop sedentary behaviors and to acquire childhood obesity which matched with our study report.

TV watching (4-8 hrs) and computer game (2-4 hrs) playing showed significant association ($p=0.00001$) with the development of childhood obesity. The result of our study matched with a cross-sectional study done by Mendoza et al. (2007) in which we found that watching TV / Video > 2 hours/day was associated with a high risk of being overweight or at risk for overweight and higher adiposity. Though the study was done in urban area and most of the children (53.6%) had no access to computer game. Total no. of children who were engaged in playing computer game < 2 hrs in a day was forty nine, (49), of them, twenty two (22) were obese and twenty seven (27) were non-obese. No significant association (<.44) was found in the development of childhood obesity. Robinson et al. (1999) showed in his study that the effect of reducing TV/Video watching and Video game use on changes in adiposity, physical activities and dietary intake and

also revealed that reducing these activities help to prevent childhood obesity.

We studied dietary habit related risk factors for the development of childhood obesity. The intake of amount of rice and also the method preparation of eggs and also the frequency of taking egg daily or weekly taken into account, taking more rice (>2.5 plate) daily and having egg daily were found as risk factors for development of childhood obesity with higher odds ratios. Excess intake of mutton and chicken (three times /weekly) and noodles (daily) were found to be risk for the development of childhood obesity with odds ratios were 3.2, 2.37 and 4.9 respectively. Odds ratios were not significant in case of fruits (OR 0.95) and vegetables (OR 0.58). Intake of fish (three times/weekly) was found healthy for the children and odds ratio was 1.2 and daily intake was found to be negatively related with the development of childhood obesity with OR of 0.56, but on the contrary if the intake decreased to once weekly, the chance of development of obesity was more (3.19). The intake of fast foods like chips (OR 6.44), burger (OR 7.1), pizza (OR 6.5), chicken roast (OR 5.5), ice cream (OR 4.3) were significantly working as risk factors in the development of childhood obesity.

The study showed the prevalence of childhood obesity was 10.7% amongst 6-9 yrs of age children in the studied population which clearly indicating that the childhood obesity is an emergency health problem among the school going children in Dhaka city. Multiple risks factors as for example, family income, family expenditure on food, dietary habits, sedentary behaviors and physical activities were identified for the development of childhood obesity in the study. Further study needs to confirm these findings which are responsible for the development of childhood obesity in Dhaka city as well as for all the children of the country.

Conclusion

The study findings showed that the prevalence of childhood obesity in the mentioned age group was 10.7% in Dhaka city indicating that the childhood obesity is an emerging health problem of school going students of Dhaka city.

Multiple risk factors like family income, family expenditure on food, dietary habits, watching TV, watching video

and computer game and physical activities were identified in this study, further study is needed to confirm the finding in Dhaka city as well as for all children of the country.

Limitation

The limitations of the study were the followings: Study was limited to Dhaka city only among the nearly similar socio-economic background. The sample size was quite inadequate to represent this emerging health problem, and the schools were selected purposively.

Acknowledgements:

The authors thank all the personnel of the School of Public Health & Life Sciences, University of South Asia, Banani & the teachers, staffs and other personnel of the schools for their constant cooperation and inspiration. I wish to express my heartfelt thanks to Mr. Shahjahan, Senior researcher of Child Sight Foundation. I must express my thanks and indebtedness to all the head of the schools who helped me and allowed me to do the job at their school premises. I also thank my wife and my deep love to my son Saqeeb and daughter Bushra, for all patience, sacrifice and continuous help extended to me.

References

1. American Academy of Paediatrics 2003, 'Prevention of Pediatric Overweight and Obesity'. Paediatrics, Vol. 112 No. 2, pp. 424-430.
2. Anderson, P, Butcher, K and Levine, PB 2003, 'Maternal employment and overweight children', Journal of Health Economics, Vol. 22 pp. 477-504.
3. Anoop Misra et al. 2008, 'Childhood obesity in Asian Indians: a burgeoning cause of insulin resistance, diabetes and sub-clinical inflammation', Asia Pacific Journal of Clinical Nutrition, Vol. 17 no. S1, pp. 172-175.
4. Dietz, WH, and Gortmaker, SL 1985, 'Do we fatten our children at the television set? Obesity and television viewing in children and adolescents', Paediatrics, Vol. 75, pp. 807-12.
5. Dietz, WH 1996, 'The role of lifestyle in health: the epidemiology and consequences of inactivity', Proceedings of the Nutrition Society, Vol. 55, pp. 829-840.
6. Dietz, WH 1994, 'Critical periods in childhood for the development of obesity', American Journal of Clinical Nutrition, Vol. 59, pp. 955-959.
7. Flegal, KM, Ogden, CL, and We, i R, et al. 2001, 'Prevalence of overweight in US children: comparison of US growth charts from the Centers for Disease Control and Prevention with other reference values for body mass index', American Journal of Clinical Nutrition, Vol. 73, no. 6, pp. 1086-1093.
8. Gupta, AK and Ahmed, AJ 1990, 'Childhood obesity and hypertension', Indian Paediatrics, vol. 27, pp. 333-337.
9. James, WPT, Rigby, NJ, Leach, RJ, Kumanyika, S, Lobstein, T, Swinburn, B 2006, 'Global strategies to prevent childhood obesity: forging a societal plan that works', International Obesity Task Force, International Association for the Study of Obesity.
10. Kaushik, B, Samiran, B, Ashish, M, and Mithu, B 2007, 'Overweight and obesity among affluent Bengalee schoolgirls of Lake Town, Kolkata, India', Maternal and Child Nutrition, Vol. 3, no. 2, pp. 141-145.
11. Magarey, A. M, Daniels, LA, and Boulton, TJ 2001, 'Prevalence of overweight and obesity in Australian children and adolescents: reassessment of 1985 and 1995 data against new standard international definitions', The Medical Journal of Australia, Vol. 174, no. 11, pp. 561-564.
12. Must, A, and Strauss, RS 1999, 'Risks and consequences of childhood and adolescent obesity', International Journal of Obesity, Vol. 23, pp. S2-S11.
13. Mendoza, JA, Zimmerman, FJ, and Dimitri AC 2007, 'Television viewing, computer use, obesity, and adiposity in US preschool children', International Journal of Behavioral Nutrition and Physical Activity, Vol. 4, pp. 44.
14. Monteiro, CA, Moura, EC, Conde, WL and Popkin BM 2004, 'Socioeconomic status and obesity in adult populations developing countries: a review', Bulletin World Health Organization, Vol. 82, no. 12, pp. 940-946.
15. Nancy, F K, John, HH, Dawn, J, Theresa, A, Nicklas, PG, and Dennis S 2007, 'Assessment of Child and Adolescent Overweight and Obesity', Paediatrics, Vol. 120, pp. S193-S228.
16. Ogden, CL et al. 2002, 'Prevalence and trends in overweight among U.S. children and adolescents, 1999-2000', The Journal of the American Medical Association, Vol. 288, no. 14, pp. 1728-1732.
17. Reilly, JJ, Armstrong, J, and Dorosty, AR et al. 2005, 'Early life risk factors for obesity in childhood: cohort study. British Medical Journal, Vol. 330, pp. 1357-1363.
18. Ravelli GP, and Belmont, L 1979, 'Obesity in nineteen year old men: family size and birth order associations', American Journal of Epidemiology, Vol. 109, pp. 66-70.
19. Robinson, TN 1999, 'Reducing children's television viewing to prevent obesity: a randomized controlled trial', The Journal of the American Medical Association, Vol. 282, pp. 1561-1567.
20. Rahman, SMM, Akter, BMD, Siddiqui, MZA and Rashid, M. 1998' Prevalence of childhood obesity in Dhaka city', Mymensingh Medical Journal, Vol. 7, pp. 3-6.
21. Rahman, SMM, Kabir, I, Khaled, MA, Bhuyan, MAH, Harun AR, Malek, MA and Khan, MR, 'Prevalence and Determinants of Childhood Obesity in Dhaka City', 10 ASCON Abstract no: 173, pp. 33.
22. Salmon, A, Crawford, BD, A Timperio, A, and Owen, N 2000, 'The association between television viewing and overweight among Australian adults participating in varying levels of leisure-time physical activity', International Journal of Obesity, Vol. 24, pp. 600-606.
23. The Asia-Pacific Perspective: Redefining obesity and its treatment', 2000, World Health Organisation, International Association for the Study of Obesity, International Obesity Task Force, Sydney.
24. Timperio, A, Crawford, D, Telford, A, and Salmon, J2004, 'Perceptions about the local neighborhood and walking and cycling among children. Preventive Medicine, Vol. 38, no. 1, pp. 39-47.
25. Tremblay, MS and Willms, JD 2003, 'Is the Canadian childhood obesity epidemic related to physical inactivity?', International Journal of Obesity, Vol. 27, no. 99, pp. 1100-1105.
26. Umesh, K, Preeti, S, Priyali, P, Sada ND, Sanjiv, B et al. 2002, 'Prevalence of Obesity Amongst Affluent Adolescent School Children in Delhi', Indian Paediatrics, Vol. 39, pp. 449-452.
27. World Health Organization 1998, 'Obesity: Preventing and Managing the Global Epidemic: Report of a WHO Consultation on Obesity', Geneva: WHO. page 1-258.
28. Wilkinson, PW, Parkin, JM, Pearson, J, Philips, PR and Sykes, P 1977, 'Obesity in childhood: A community study in Newcastle upon Tyne', Lancet, Vol. 1, pp. 350-352.