

Original Article

Antibiotic Sensitivity Pattern of Uropathogens at Khwaja Yunus Ali Medical College Hospital

Ahmed AA¹, Alam AN², Fatmi R³, Rahman MM⁴, Munia MS⁵, Juyee NA⁶

Abstract

Increasing resistance among urinary tract pathogens to conventional drugs is found alarming worldwide. The aim of this study was to obtain data on susceptibility patterns of isolated uropathogens from urine samples of patients attending at Khwaja Yunus Ali Medical College Hospital, Enayetpur, Sirajgonj, Bangladesh a tertiary health care center of remote area to currently used antimicrobial agents.

A total of 656 urine samples were studied of which 163 (24.8%) were culture positive. Among 163 culture positive cases all uropathogenic isolates were identified. Among the isolates, E. coli was the most predominant 116 (71.1%) followed by Pseudomonas aeruginosa 29 (17.8%). Staphylococcus saprophyticus 12 (7.4%) and Klebsiella pneumoniae 4 (2.4%) and Acenatobacter sp. 2 (1.2%). The majority of isolated E. coli were sensitive to Meropenam (58.6%) and Amikacin (57.7%) followed by gentamicin (52.6%), amoxiclav (43.1%), ciprofloxacin (42.2%). Sensitivity & resistance rate in Staphylococcus saprophyticus were same (50%) in Cephadrine. Where complete (100%) resistance was shown to Ampicillin in Pseudomonas aeruginosa, Klebsiella pneumoniae and Acenatobacter sp. However most of isolates were poorly sensitive to ampicillin (1.7%), erythromycin (2.5%), cefuroxin (3.4%) and amoxicillin (9.5%).

This area-specific monitoring studies aimed to gain knowledge about the type of pathogens responsible for UTIs and their resistance patterns may help the clinician to choose the right empirical treatment. Meropenem, amoxiclav, amikacin, gentamicin, second-generation of both cephalosporins and quinilones are found the most sensitive against the common uropathogens which might be used in the treatment of UTI.

Key Words: Urinary tract, E. coli, Pseudomonas aeruginosa, Staphylococcus, Klebsiella

Introduction

The increasing trend of emerging bacterial resistance to a large number of antibiotics undermines the ability of antibiotics to control infections¹⁻³. Urinary tract infections (UTIs) are one of the most common bacterial infections both in the community and hospital setting.

An estimate of patients suffering from UTI is around 150 million per annum across the Globe, which may rise to 75% in the female population by the age of 24, and 15-25% of this group will suffer from a relapse of this disease⁴⁻⁶. By considering above facts this study is aimed to explore the trends of the antibiotic-resistant uropathogens and their susceptibility toward various

1. Dr. Abdullah Akhtar Ahmed, Professor and HOD, Department of Microbiology, KYAMC.
2. Dr. Ahmed Nawsher Alam, Associate Professor, Department of Microbiology, KYAMC.
3. Dr. Rejwana Fatmi, Lecturer, Department of Microbiology, KYAMC.
4. Dr. Md. Mashiur Rahman, Lecturer, Department of Microbiology, KYAMC.
5. Dr. Marufa Sultana Munia, Lecturer, Department of Microbiology, KYAMC.
6. Dr. Nusrat Akhtar Juyee, Lecturer, Department of Pathology, Ibrahim Medical College, Dhaka.

Place of Study: Department of Laboratory Services, Khwaja Yunus Ali Medical College Hospital, Enayetpur Sharif, Sirajgonj.

Correspondance: Dr. Abdullah Akhtar Ahmed, Professor and HOD, Department of Microbiology, KYAMC.

The present study will further help in formulating the guide- lines for establishing a proper empirical therapy in Khwaja Yunus Ali Medical College Hospital, Enayetpur, Sirajgonj for UTIs while awaiting culture sensitivity reports. Meropenem, amikacin, amoxiclav, gentamycin and ciprofloxacin, the third and fourth generation cephalosporins were found the most commonly sensitive antimicrobial drugs which might be prescribed in the treatment of UTI.

Materials and Methods

This study was done in the Microbiology Laboratory Services department of Khwaja Yunus Ali Medical College Hospital in 2013. The patients comprised both sexes and all age groups. Urine samples were collected by using sterile, dry, wide mouthed leak proof plastic container and bacteriological tests were done for diagnosis of etiological agents of UTI. Standard method for isolation and identification of the organisms were carried out with all urine samples. Blood agar and MacConkey agar media were plated using a calibrated loop withdrawing 0.01 ml of urine sample. The bacterial colonies were counted and multiplied by 100 to give an estimate of the number of bacteria present per milliliter of urine. Significant growth was determined as >10⁵ colony forming units (CFU)/ ml of midstream urine samples and >10² CFU/ml of a catheter specimen⁷. Gram negative bacteria were identified by morphological study, oxidase test, routine biochemical tests such as motility test, indole and urease production and Triple Sugar Iron reaction. Gram positive bacteria were identified by catalase, coagulase and Novobiocin tests. Samples which showed significant colony count were taken into consideration and sensitivity pattern of the isolated organisms was determined by modified Kirby-Bauer technique using Mueller-Hinton agar⁸. Interpretation of results was done measuring the sizes of zones of inhibition and reported according to "The Clinical Laboratory Standard Institute" (CLSI) guidelines⁹. Antibiotics for uropathogens were tested include amoxiclav, amikacin, cotrimoxazole, gentamicin, imipenem, meropenem, cephradine, ceftazidime, ceftriaxone, cefixitin, cefuroxin, ciprofloxacin, erythromycin, tetracycline, ceftriaxone, and azithromycin.

Results

Prevalence of Uropathogens:

In this study, a total of 656 urine samples were tested

during from patients of all age groups in both sexes. Among them 163 (24.85%) were culture positive (Table 1).

Table-1 Distribution of specimens with or without culture positive.

| Urine culture | Frequency | Percent |
|------------------|-----------|---------|
| Culture positive | 163 | 24.85 |
| Culture negative | 493 | 75.15 |
| Total | 656 | 100 |

The most common isolated uropathogens were *E. coli* (71.1%) and *Pseudomonas aeruginosa* (17.8%). *Staphylococcus saprophyticus* (7.4%). Other bacterial species named *Klebsiella pneumoniae* and *Acenatobacter sp.* were also found in patients with UTI although they were least frequent (Table 2).

Table 2 Frequency of bacterial agents isolated from urine specimens

| Bacterial Isolates | Number | Percentage |
|-------------------------------|--------|------------|
| <i>Escherichia coli</i> | 116 | 71.1 |
| <i>Pseudomonas aeruginosa</i> | 29 | 17.8 |
| <i>S. saprophyticus</i> | 12 | 7.4 |
| <i>Klebsiella pneumoniae</i> | 4 | 2.5 |
| <i>Acenatobacter sp.</i> | 2 | 1.2 |
| Total | 163 | 100 |

This study shows, more male patients (58.3%) suffered from UTI than female (41.7%). The incidence of UTI was ranged in patients between 1-65 years old. The highest number of male patients with UTI was found within the age range of 46-65 years and female patients by the age range 31 - 45 (Table 3).

Table 3: Distribution of UTI among 163 patients in different age level and their relation to sex.

| Age Level in years | Sex | | | |
|--------------------|-------------|---------|---------------|---------|
| | Male (n=95) | | Female (n=68) | |
| | Frequency | Percent | Frequency | Percent |
| 0-5 | 2 | 2.1 | 1 | 1.5 |
| 6-15 | 1 | 1 | 2 | 2.9 |
| 16-30 | 6 | 6.3 | 17 | 25 |
| 31-45 | 17 | 17.9 | 24 | 35.3 |
| 46-65 | 36 | 37.9 | 18 | 26.5 |
| Ø 65 | 33 | 34.7 | 6 | 8.8 |

Antibiotic Susceptibility:

The isolated bacteria showed wide range of differences in their susceptibility pattern to the tested antibiotics. As the result indicated, high proportions of the test organisms were sensitive to meropenem, amikacin, gentamicin, amoxiclav, ciprofloxacin, azithromycin. Again, a significant percentage of the test organisms also showed resistance to ampicillin, amoxicillin, erythromycin and carbenicillin (Table 4).

| Antibiotics | Isolated uropathogens | | | | |
|---------------|-----------------------|--------------|-----------|----------|------------|
| | E. coli | S. saprophyt | Ps. aerug | K. pneum | A. baumani |
| Ampicillin | 1.7 | 16.7 | | | |
| Amoxicillin | 9.5 | 25.0 | 13.8 | | |
| Amoxiclav | 43.1 | 33.3 | 31.0 | 50.0 | |
| Amikacin | 57.7 | 33.3 | 65.5 | | |
| Azithromycin | 42.2 | 33.3 | 51.7 | 50.0 | |
| Carbenicillin | | 25.0 | 3.4 | | |
| Ceftazidim | 19.0 | 8.3 | 27.6 | 50.0 | |
| Ceftriaxone | 29.3 | 25.0 | 27.6 | | |
| Cefixim | 17.2 | | 10.3 | 50.0 | |
| Cefuroxin | 9.5 | 8.3 | 3.4 | | |
| Cephadrine | 27.6 | 50.0 | 13.8 | 25.0 | 50.0 |
| Ciprofloxacin | 42.2 | 33.3 | 48.3 | 25.0 | 50.0 |
| Cloxacillin | | 41.7 | | | |
| Cotrimoxazole | 24.1 | 25.0 | 13.8 | 25.0 | |
| Doxycyclin | 20.7 | 33.3 | 10.3 | 25.0 | |
| Erythromycin | 2.5 | | | | |
| Gentamicin | 52.6 | 41.7 | 31.0 | 50.0 | 50.0 |
| Imipenem | 28.4 | 16.7 | 48.3 | 25.0 | 50.0 |
| Levofloxacin | 14.6 | | 13.8 | | |
| Meropenem | 58.6 | 25.0 | 55.2 | 25.0 | 50.0 |
| Piperacillin | 19.0 | 16.7 | 27.6 | | 50.0 |
| Tobramycin | 10.3 | 41.7 | 10.3 | 25.0 | 50.0 |

Table 4: Susceptibility pattern (%) of bacterial agents isolated from urine specimens

Discussions

This paper describes the first study undertaken at Khwaja Yunus Ali Medical College Hospital in 2013 to evaluate susceptibility patterns of bacterial strains isolated from UTIs. It provides valuable laboratory data concerning uropathogens and enables the situation of this hospital to be compared with that in other countries. Here we described the relationships between isolated bacterial agents and antibiotic susceptibility and resistance of UTIs. The sex distribution of patients in our study is analogous with those of other reported studies, showing a predominance of males (58.3%) with UTI. The elevated incidence of infection among males is related to the differences between male and female in economic solvency and privilege which prevails in the

underdeveloped countries⁸. The incidence and prevalence of UTI in both men and women was found to increase with age as seen in Sweden and Finland during 2005-2007⁹. The most common uropathogens in our study were *E. coli* (71.1%) and *Pseudomonas aeruginosa* (17.8%). Although the percentage of *E. coli* is much higher in our study, it supports another finding of Chittagong indicating that *E. coli* is the principal etiological agent of UTI, accounting for 82.6% of the screened cases¹⁰. In another study, it was reported that predominant uropathogens are *E. coli* followed by *Pseudomonas aeruginosa* which also partially support our study where the 2nd most common uropathogen was found *Klebsiella pneumoniae*¹¹. Infection frequency of *Staphylococcus saprophyticus* and *Acinetobacter* sp. were found to be very few in this study (Table 3) which is also affirmed by another work. The most effective antimicrobial agents in our study were meropenem, amikacin, gentamicin, amoxiclav, ciprofloxacin, azithromycin. Again, a significant percentage of the test organisms also showed resistance to ampicillin, amoxicillin, erythromycin and carbenicillin (Table 4). It has been reported that meropenem is the most effective antibiotic against *E. coli*¹². Our result was further supported by another study where the susceptibility rate of *E. coli* to amikacin remained 93-100%¹³. Again, most of the pathogens showed considerable resistance to ampicillin, amoxicillin, cefuroxin and erythromycin (Table 4). The widespread use, more often the misuse, of antimicrobial drugs has led to a general rise in the emergence of resistant bacteria. It is reported that in the United States there is a rise in the resistant strains to ampicillin due to its widespread use¹⁴. With this evidence from our study, we can suggest meropenem, amikacin, ceftazidime, gentamicin and imipenem to be prescribed as the empirical treatment for UTI.

Conclusion

By keeping the emerging antimicrobial resistance in mind, it is strongly recommended that antibiotic therapy should only be commenced after the culture and sensitivity report from the microbiology laboratory. This would not only help in the sensible use of antibiotics but also would restrain the spreading of antimicrobial resistant strains in the community as well as in the hospital.

Acknowledgment

We are grateful to the Chairman of the Trusty Board and Director of Khwaja Yunus Ali Medical College Hospital to conduct this study. We thank Mr. Md. Abdul Karim,

Medical Technologist of the department of Laboratory Services of Khwaja Yunus Ali Medical College & Hospital for assisting in laboratory work.

References

1. Bahadin J., Teo S. S. H., Mathew S. (2011). Aetiology of community-acquired urinary tract infection and antimicrobial susceptibility patterns of uropathogens isolated. *Singapore Med. J.* 52415-420
2. Nerurkar A., Solanky P., Naik S. S. (2012). Bacterial pathogens in urinary tract infection and antibiotic susceptibility pattern. *J. Pharm. Biomed. Sci.* 21 1-3
3. Sood S., Gupta R. (2012). Antibiotic resistance pattern of community acquired uropathogens at a tertiary care hospital in Jaipur, Rajasthan. *Indian J. Community Med.* 37 39-44
4. Hunstad D. A. (2010). Intracellular lifestyles and immune evasion strategies of uropathogenic *Escherichia coli*. *Annu. Rev. Microbiol.* 64 203-221
5. Mukherjee M., Basu S., Mukherjee S. K., Majumder M. (2013). Multidrug-resistance and extended spectrum beta-lactamase production in uropathogenic *E. Coli* which were isolated from hospitalized patients in Kolkata, India. *J. Clin. Diagn. Res.* 7 449-453
6. Pouwels K. B., Visser S. T., Hak E. (2013). Effect of pravastatin and fosinopril on recurrent urinary tract infections. *J. Antimicrob. Chemother.* 68 708-714
7. Forbes BA. Baily and Scott's diagnostic microbiology. 10th ed. St. Louis, Missouri: Mosby; 1998; 283-304.
8. Bauer AW, Kirby WM, Sherris JC, Tench JM. Antibiotic susceptibility method. *Am J Clin Path.* 1996; 45: 493- 496.
9. Clinical and Laboratory Standards Institute. 2006. Performance standards for antimicrobial susceptibility testing; 16th informational supplement. M100-S16. Clinical and Laboratory Standards Institute, Wayne, PA.
8. Men in Families and Family Policy in a Changing World. United Nations publication. New York, 2011. Sales No. E.11.IV.1. ISBN 978-92-1-130306-3
9. Irene Eriksson (2011) Urinary Tract Infection- a serious health problem in old women, Umeå University Medical Dissertations, New Series No 1410.
10. Chowdhury S and Parial R. (2015). Antibiotic Susceptibility Patterns of Bacteria among Urinary Tract Infection Patients in Chittagong, Bangladesh. *SMU Medical Journal*, 2(1): 114-127.
11. Olafsson M, Kristinsson KG, Sigurdsson JA (2000) Urinary tract infections, antibiotic resistance and sales of antimicrobial drugs--an observational study of uncomplicated urinary tract infections in Icelandic women. *Scand J Prim Health Care* 18:35-8.
12. Sultana S, Khatun M, Khoibar Ali M, Mawla N, Akhter N. (2015). Pattern of antibiotic sensitivity of bacteria causing urinary tract infection in a private Medical College Hospital in Dhaka. *Bangladesh Journal of Medical Science* 14 (1):70-74.
13. Shigemura K, Tanaka K, Okada H, Nakano Y, Kinoshita S, Gotoh A, Arakawa S, Fujisawa M (2005) Pathogen occurrence and antimicrobial susceptibility of urinary tract infection cases during a 20-year period (1983-2002) at a single institution in Japan. *Jpn J infects Dis* 58:303-8.
14. Sahm DF, Critchley IA, Kelly LJ, Karlowsky JA, Mayfield DC, Thornsberry C, Mauriz YR, Kahn J (2001) Evaluation of current activities of fluoroquinolones against gram-negative bacilli using centralized in vitro testing and electronic surveillance. *Antimicrob Agents Chemother* 45:267-74.