

# Perception about Pain Relief During Normal Labour Among Health Care Providers Conducting Delivery

Tasnim S<sup>1</sup>

## Abstract:

*Pain during labour is a physiological phenomenon that varies in intensity among women and subjected to many social and cultural modifiers.*

*The objective of the study was to explore the perception and practice of the primary level health care providers who conducts normal vaginal delivery regarding pain relief during labour.*

*A cross sectional questionnaire survey was done at institute of child and mother health during May and June 2009 among a group of senior staff nurses and family welfare visitors.*

*Among 97 respondents 75.3% were senior staff nurses and 24.7% were Family welfare visitors. Only 6.2% thought a women with labour pain should receive an analgesic, 7.4% gives an analgesic and 10.5% reported to receive such drug during their own childbirth. About 58.6% reported to use injection hyoscine butyl bromide and 6.9% uses injection Pethidine analgesia during labour. Forty percent reported to carry out some activity to comfort women in labour. Those were giving assurance (88.7%), explaining the mother about the process of labour (84.5%) and 77.3 % would allow companion in the labour room. About two thirds respondents thought that pain relief may delay progress of labour, 69.5% apprehend fetal distress while 60% are of the opinion that women should endure the natural pain.*

*Key words: Analgesia, labour pain, supportive care, health care provider*

## Introduction:

In 1591, eufane MacAyane of Edinburgh was dragged from her home, her pleas for mercy was ignored and she was buried alive into a pit because she asked for pain relief during her difficult labour of twin sons<sup>1</sup>. The church teachings of that day regarded the pain of childbirth as a punishment justly inflicted by God to women and asking for relief from pain was presumably against God's wishes and eufane was punished for that sin<sup>2</sup>.

Pain during labour is a physiological phenomenon. The evolution of pain during first stage of labour is associated with ischaemia of the uterus during contraction as well as effacement and dilation of Cervix<sup>3</sup>. In the second stage pain is caused by stretching of the vagina, perineum & compression of pelvic structures. However, pain sensation is a response of the total personality and a subjective phenomenon. In a study from Nizeria 70% women rated labour pain as moderate to severe, 32% said they would not want any pain relief and 33% would want it to be eliminated<sup>4</sup>. Severe pain is common affecting some 66-70% nulliparous and 35-40% multiparous labours<sup>5</sup>.

Pain relief in childbirth is subject to many social and cultural modifiers, which continue to change. There is no other circumstance in which it is considered acceptable for a person to experience untreated severe pain even while under a physician's care<sup>6</sup>. Control of pain rather than absolute amelioration is seen by many to provide greater satisfaction. The importance of analgesia as a contribution to overall satisfaction has been recognized increasingly in the last 50 years<sup>7</sup>. Sir James Simpson, professor of Obstetrics at the university of Edinburgh first advocated the use of ether for Obstetric analgesia in 1850's<sup>2</sup>. Queen Victoria in 1853 had the pains of childbirth relieved by intermittent whiffs of Chloroform and thereafter the opposition from churches regarding obstetric analgesia remained silent.

Analgesia is the loss or modulation of pain perception and its use during labour should produce optimum pain relief with minimum side effects to mother & baby<sup>3</sup>. There are both pharmacological and non pharmacological methods. Pharmacological methods include opioids, inhalation analgesia and regional analgesia. Pethidine (Demerol or Meperidine) is commonly recommended drug for relieving pain during labour. Pain relief is more satisfactory if hyoscine is given at same time and by using hyosine the dose of Pethidine can be reduced<sup>6,8</sup>. Lumbar epidural analgesia for painless labour have been introduced since 1990s in developed countries and the number of women receiving an anaesthetic procedure for pregnancy increased to more than 60%<sup>5</sup>. Among the non pharmacological methods there are Psychoprophylaxis, Hypnosis and natural child birth that emphasized the reduction of tension to induce relaxation<sup>4</sup>.

In Bangladesh use of analgesic drug to relieve pain during labour is not a routine practice. The current study was

1. Dr Saria Tasnim.  
FCPS. M MED. Diploma in Community epidemiology  
Associate Professor, Obstetrics and Gynaecology.  
Institute of Child & Mother Health, Dhaka, Bangladesh

carried with the purpose of exploring perception and practice of the primary level health care providers who conducts normal vaginal delivery regarding the pain relief during labour.

#### Method:

This was a cross sectional self administered questionnaire survey among the senior staff nurses and family welfare visitors who participated in different short courses of training at institute of child and mother health during May and June 2009.

#### Results

Among 97 respondents 75.3% were senior staff nurses and 24.7% were Family welfare visitors (Fig.1).




Figure 1: Profile of respondents (n= 97)

Eighty four percent works in the labour room. Only 6.2% thought a women with labour pain should receive an analgesic, 7.4% gives an analgesic to labour patient and 10.5% received such drug during their own childbirth (Table 1).

Table 1: Attitude regarding use of analgesic during labour

| Statement                                                           | Percentage |
|---------------------------------------------------------------------|------------|
| Every patient at normal labour should receive analgesic             | 6.2        |
| Gives analgesic to patients at labour                               | 7.4        |
| Received analgesic during her own labour                            | 10.5       |
| Use non pharmacological method for pain relief during normal labour | 40.6       |
| Use of analgesic will influence progress of normal labour           | 85.2       |

However, 58.6% reported to use injection hyoscine butyl bromide and 6.9% uses injection Pethidine (Table 2).

Table 2: Type of medicine used for pain relief during normal labour\*

| Name of medicine                | Percentage |
|---------------------------------|------------|
| Injection Pethidine             | 6.9%       |
| Injection Hyosine butyl bromide | 58.6%      |
| Inj Diclofenac                  | 3.4%       |
| Tablet Butapan                  | 60.9%      |
| Tablet Paracetamol              | 4.6%       |
| Others                          | 5.7%       |

Multiple response

Forty percent stated to employ some activity to comfort women in labour. Those were assurance (88.7%), explaining the mother about the process of labour (4.5%) and 77.3 % would allow companion in the labour room (Table 3).


Table 3: Type of non pharmacological method for pain relief during normal labour

| Method used                          | Percentage |
|--------------------------------------|------------|
| Give assurance                       | 88.7       |
| Massage the back                     | 53.6       |
| Allow patient to move freely         | 78.4       |
| Show the patient to take deep breath | 40.2       |
| Show the patient how to bear down    | 72.2       |
| Help to do labour exercises          | 74.2       |
| Explain the process of labour        | 84.5       |
| Allow companion of her choice        | 77.3       |
| Rebuke for screaming                 | 9.2        |

Multiple response

About two thirds respondents thought that pain relief may delay progress of labour, 69.5% feared fetal distress while 60% are of the opinion that women should endure the natural pain of labour (Fig-2).

Figure 2:Opinion regarding effect of analgesia on outcome of labour



#### Discussion

Common attitude of health care providers were restrictive towards using analgesic for normal labour, and they also thought it would interfere with normal progress of labour. However, they expressed positive attitude to help mothers to cope with labour pain through assurance and carry out some activities to comfort the women in labour. The drug Hyoscine butyl bromide was reported to be commonly used but there is lack of evidence to show any analgesic effect on labour pain. In developed countries there is an increasing trend of using epidural and other regional anesthesia for painless labour. In a study from Sweden it was found that among all women who had vaginal deliveries between 1983- 1986 lumbar epidural was used in 16%, paracervical

block in 12%, pethidine or morphine in 49% and pudendal block in 62%<sup>9</sup>.

American college of obstetrician and gynaecologist suggested that maternal request should be sufficient justification for pain relief during labour and should not be denied because of absence of other medical indications<sup>10</sup>. However, there are simultaneous movements to ensure natural childbirth as much as possible. Effective pain relief does not ensure a satisfactory birthing experience rather the interpersonal relationship established between patient and health care provider are more important in the determination of satisfaction with management<sup>11,7</sup>.

In most developing countries home birth is a norm. Typically in home birth the woman's mother, sister, aunts, and female friends remained with the parturient women throughout labour and offer her social support<sup>12</sup>. The social support includes emotional support and reassurance, comfort measures (e.g. bath or shower, massage, soothing touch, breathing and relaxation techniques), and encouraging the woman to change positions and work with her body<sup>13</sup>. On the other hand in most hospital setting the women in labour is alone and often frightened by the intermittent appearance and disappearance of unknown people, including obstetricians; midwives; nurses; and medical, nursing, or midwifery students<sup>14</sup>. Studies have demonstrated that labor nurses in some institutions spend as little as 6.1% of their time performing supportive activities for the laboring women in their care<sup>15, 16</sup>. Some negative experiences regarding hospital delivery was reported that in the labour room women are compelled to adopt birthing position according to the existing policy, privacy are not well maintained and health care providers are often rude, impatient and reluctant to their concerns<sup>17</sup>.

In Bangladesh majority of births takes place at home and 83% are conducted by traditional birth attendants<sup>18</sup>. Study has revealed that the preference for such births were that mothers felt at ease with their near ones at home, the traditional birth attendant was either a relative or like a keen to them accustomed with their customs and the women would enjoy more control over her own childbirth<sup>19</sup>.

Recognizing the importance of continuous human companionship during labour in some countries for example in USA 'Doulas' are being employed in hospital based labour and are part of the delivery team<sup>20</sup>. A doula is a woman who do not perform clinical tasks but has been professionally trained to provide continuous social support to the birthing family during labor and delivery and their role complements that of the medical care providers<sup>20</sup>. There is evidence that Doula care has resulted in significant reductions in requests for epidurals and delay in their administration<sup>21,22</sup>. Childbirth has historically been a deeply entrenched social event in which women bonded as they

shared a rite of passage. It is often stated that as culture advances the labour becomes painful, for in women of primitive races pain was absent.

### Conclusion

The health care providers in this study were conservative in recommending use of analgesia in normal labour and opined that the women should perceive pain of childbirth as a natural process that should be endured. However, they were in favour of supportive care to comfort the mother. In the spectrum of uninterrupted natural labour pain to intervened painless labour what is the choice of our mothers and what options can be made accessible in the present context needs to be explored.

### References

1. T E Keys. The history of surgical anesthesia ( New York: Schumans 1945)
2. Meyer Friedman, Gerald W Friedland. Medicine's 10 greatest discoveries. University press limited. India 1999.
3. Ralph W Yarnell and John S McDoland. Obstetric analgesia and anesthesia. In: Alan H Decherney & Lauren Nathan (Ed) current Obstetric & Gynaecologic diagnosis and treat 9th edition Lange medical books/McGraw -Hill. 2004
4. A Fola Faponide, Oluwafemi Kuti . Perception of labour pain by Pregnant Women in Southwestern Nigeria Trop J Obstet Gynaecol 2004; 21:153-155
5. John A Crowhurst. Analgesia and anaesthesia. In: Edmond D Keith (ed) Dewhurst's text book of Obstetrics Gynaecology. Blackwell publishing. USA. Seventh edition 2007.
6. Ian Donald. Practical Obstetric problems. PG publishing Pte limited. New Delhi. Fifth edition 1990.
7. Bailliere's clinical Obstetrics and Gynaecology 1998; 12:499-512
8. Hingson R A, Hellsman LM. Anaesthesia for obstetrics J B Lippincot CO, Philadelphia. 1956
9. Eva Gerdin, Sven Cnattingius. The use of obstetric analgesia in Sweden 1983-1986;97:789-796.
10. ACOG technical bulletin. Obstetric analgesia and anaesthesia. International journal of gynaecology and obstetrics 1996; 54 : 281-292
11. BarbaraM. Morgan, ChristopherJ. Bulpitt, Pamela Clifton and PeterJ. Lewis. Analgesia and satisfaction in childbirth (the queen charlotte's 1000 mother survey). 1982; 321-22: 480

12. Klaus M, Kennell J, Berkowitz G, Klaus P. Maternal assistance and support in labor: father, nurse, midwife, or doula? *Clin Consult Obstet Gynecol* 1992; 4:211–7.
13. Hodnett ED. Caregiver support for women during childbirth. *Cochrane Database Syst Rev* 2000;2:CD000199.
14. Enkin M, Keirse MJ, Neilson J, et al. A guide to effective care in pregnancy and childbirth. 3rd ed.:Oxford University Press; 2000.
15. Gagnon AJ, Waghorn K. Supportive care by maternity nurses: a work sampling study in an intrapartum unit. *Birth* 1996;23:1–6.
16. Gagnon AJ, Waghorn K, Covell C. A randomized trial of one-to-one nurse support of women in labor. *Birth* 1997;24:71–7.
17. Afsana K, Sabina S F. The challenges of meeting rural Bangladesh women's needs in delivery care . *Reproductive health matters* 2001; 9 ( 18) : 79-88.
18. Bangladesh Demographic and health survey 2007. National Institution of Population research and Training 2008. Dhaka.
19. Saria Tasnim , A K M Shahabuddin, S Chowdhury, P C Barua, AKMF Rahman, A Rahman et al. Community trial of skilled midwifery serices for improving safe motherhood in urban setting. Final report. 2001. Dhaka
20. Bruce A. Meyer, Jane A. Arnold, Debra Pascali-Bonaro, Social Support by Doulas during Labor and the Early Postpartum Period. *Hospital Physician* September 2001 57-65
21. Thorp JA. Weighing the benefits of epidural analgesia during labor. *Contemp Ob/Gyn* 1997; 42:95–106.
22. Lieberman E. No free lunch on labor day. The risks and benefits of epidural analgesia during labor. *J Nurse Midwifery* 1999;44:394–8.