



Original Article

Distribution of Bacterial Strains and their Antibiotic Sensitivity Profile in an Urban Referral Hospital

M. Bulbul Hasan¹, S Gul Nahar², M. Nawshad Ali³, Mst. Rokeya Khatun⁴, Chinmoy Kanti Das⁵

Abstract

The present study was done to assess the distribution of bacterial pathogens with their pattern of antibiotic susceptibility in an urban referral hospital in RMCH. A total of 393 bacteria strains were isolated from various specimens over a 10-months period. The majority of the organisms were *Escherichia coli* (33.33%) followed by *Klebsiella* species (27.48%), *Staphylococcus aureus* (17.05%), *Acinetobacter* species (8.14%), *Pseudomonas* species (7.12%), and others. The third-generation Cephalosporins like Ceftriaxone, Ceftazidime and Cefotaxime were sensitive to Ciprofloxacin of various Enterobacteriaceae was only between 33-40% compared to 52.8-37.9% against Gentamicin. Majority of the Enterobacteriaceae were resistant to Ampicillin, whereas almost all of the Enterobacteriaceae (94-100%) were sensitive to Imipenem. About 97.0% *Acinetobacter* species were susceptible to Imipenem. Sensitivity of the organism (*Acinetobacter*) to third-generation Cephalosporins ranged between 50-56%, whereas 40.6% were found sensitive to Ciprofloxacin. The sensitivity to Chloramphenicol, Co-trimixazole, Cephalexin and Ampicillin ranged between 9.3% to 34.3%. About 93.0% of *Pseudomonas* species were sensitive to Imipenem. The rate of susceptibility to Gentamicin and Netilmicin was higher than those of the Ciprofloxacin and Ceftriaxone (37.8% and 53.5% vs. 39.2%). About 70% of isolated *S. aureus* were resistant to Oxacillin but all were sensitive to Vancomycin. The result of this study would help the physicians to make a judicious choice of antibiotics for therapeutic purposes.

Key words: distribution, bacterial strains, antibiotic sensitivity profile, urban referral hospital, Enterobacteriaceae

TAJ 2012; 25: 72-76

Introduction

Infectious diseases still remain a major cause of morbidity and mortality in third-world countries including Bangladesh. Bacterial pathogens resistant commonly used antimicrobials are now creating a challenge to the clinicians and researchers. The multi-drug-resistant organisms are a serious medical problem that has significantly affected the treatment of infectious

diseases^{1,2} and has become a major clinical concern globally^{3,4}. Bacterial resistance pattern to antimicrobial agents can differ significantly from one country to another and within a country as evidenced by several recent surveillance studies^{5,6,7,8}.

Introduction of newer antimicrobial agents is usually followed sooner or later by emergence of bacterial resistance of these drugs for many

¹ Associate Professor, Department of Microbiology, Rajshahi Medical College, Rajshahi.

² Assistant Professor, Department of Microbiology, Rajshahi Medical College, Rajshahi.

³ Associate Professor, Department of Pediatric Surgery, Rajshahi Medical College, Rajshahi.

⁴ Assistant Professor, Department of Gynae and Obstetrics, Rajshahi Medical College, Rajshahi.

⁵ Associate Professor, Department of Community Medicine, Rajshahi Medical College, Rajshahi.

reasons⁹. Development of multi-drug resistance in clinical isolates like *Salmonella typhi*, *Pseudomonas* species and *Klebsiella* species has been reported in Bangladesh^{10,11,12}. The study on antibiotic susceptibility pattern is particularly important in developing countries that do not control antibiotic usage and maintain adequate epidemiological surveillance.

Therefore, the present study has been designed to find out the antibiotic resistance patterns of medically important bacteria in an urban hospital. The outcome of this study might enable to determine the trend of drug resistance prevailing in Bangladesh.

Methods

This was a prospective study carried out in RMCH. Samples were collected over a 10-month period during January to October, 2005 from both sexes and different age groups. A total of 393 clinical isolates were tested and the specimens included were Urine, Pus, Sputum, Vaginal swab, Throat swab and Conjunctival swab.

All samples were routinely cultured on MacConkey and Blood agar plates. In addition to these plates, Chocolate agar media were used for culturing Pus, Vaginal swab and Conjunctival swab specimens. After overnight incubation, plates were checked for the presence of suspected pathogens. All the suspected colonies were identified by colony characteristics, motility, Gram staining results, and biochemical reactions¹³.

Antimicrobial susceptibility test of the isolated organisms was done by disk diffusion method using the Kirby-Bauer technique¹⁴ and as per recommendations of the national Committee for clinical Laboratory Standards⁶. Antimicrobial agents used for determining antibiogram of isolated organisms were Penicillin (10 units), Ampicillin (10 µg), Cefuroxime (30 µg), Ceftazidime (30 µg), Cefotaxime (30 µg), Cotrimoxazole (25 µg), Imipenem (10 µg), Gentamicin (10 µg), Tetracycline (30 µg), Ciprofloxacin (5 µg), Vancomycin (30 µg), Chloramphenicol (30 µg), Netilmicin (30 µg),

Oxacillin (1 µg), Erythromycin (15 µg), Rifampicin (5 µg), Fusidic acid (10 µg), Cephalexin (30 µg), and Ceftriaxone (30 µg). All disks were obtained from Oxoid Ltd, Basinstoke, Hampshire, UK.

In order to monitor the quality of the test result, on each day of testing the reference ATCC strains *Escherichia coli* ATCC 25922 and *Staphylococcus aureus* ATCC 292213 were included. The zone of inhibition was compared with recommended standard values¹⁵.

Results

A total of 393 strains of bacteria were isolated from various specimens. The specimens were Urine (189, 48.09%), Sputum (88, 22.39%), Pus (71, 18.07%), Throat swab (20, 5.09%) and others (25, 6.36%), including Vaginal and Conjunctival swabs.

Out of 393 organisms isolated, majority were *Escherichia coli* (131, 33.33%) followed by *Klebsiella* species (108, 27.48%), other isolates included *Staphylococcus aureus* (67, 17.05%), *Acinetobacter* species (32, 8.14%), *Pseudomonas* species (28, 7.12%) and *Providencia* species (11, 2.80%). (Table 1)

Table 1: Pattern of organism isolated from various samples

Table 1: Pattern of organism isolated from various samples

Serial	Name of organisms	Number	Percentage
1.	<i>Escherichia coli</i>	131	33.33
2.	<i>Klebsiella</i> species	108	27.48
3.	<i>Staphylococcus aureus</i>	67	17.05
4.	<i>Acinetobacter</i> species	32	8.14
5.	<i>Pseudomonas</i> species	28	7.12
6.	<i>Providencia</i> species	11	2.80
7.	<i>Proteus</i> species	09	2.29
8.	<i>Enterobacter</i> species	01	0.25
9.	Others*	06	1.53
	Total	393	100

Others indicate *S. saprophyticus* (n=3), *S. epidermidis* (n=2) and beta hemolytic streptococci (n=1)

Among the total bacterial isolates, 260 (66.6%) were members of the Enterobacteriaceae family including *E. coli*, *Klebsiella* species, *Providencia* species, *Proteus* species, and *Enterobacter* species. Considering susceptibility pattern with different antimicrobials, it is found that almost all of the Enterobacteriaceae were sensitive Imipenem (94-

100%). The third-generation Cephalosporins like Ceftriaxone, Ceftazidime and Cefotaxime were sensitive against 45-66% isolated Enterobacteriaceae was only between 33-40%. Sensitivity to Gentamicin ranged between 52.8-67.9%. (Table II).

Table II: Sensitivity of the isolated Enterobacteriaceae (n=260)

Organisms	Number (%) sensitive to-												
	Amp	Cepl	CXM	CAZ	CRO Gen	CTX	Tet	Cip	Chl	Cot	Imp	Net	
<i>E.coli</i> (n=131)	10 (7.6)	28 (21.3)	49 (37.4)	74 (56.4)	76 (58.0)	80 (61.1)	43 (32.8)	89 (67.9)	53 (40.4)	83 (63.3)	31 (23.6)	129 (98.4)	78 (59.5)
<i>Klebsiella</i> spp (n=108)	6 (5.5)	19 (17.5)	38 (35.1)	60 (55.5)	56 (51.8)	62 (57.4)	44 (40.7)	57 (52.8)	39 (36.1)	56 (51.8)	33 (30.5)	102 (94.4)	67 (62.0)
<i>Providencia</i> spp (n=11)	01 (9.0)	02 (18.1)	05 (45.4)	06 (54.5)	06 (54.5)	05 (45.4)	06 (54.5)	07 (63.6)	04 (36.3)	05 (45.4)	04 (36.3)	11 (100)	08 (72.7)
<i>Proteus</i> spp (n=9)	01 (11.1)	02 (22.2)	04 (44.4)	05 (55.5)	05 (55.5)	06 (66.6)	04 (44.4)	06 (66.6)	03 (33.3)	04 (44.4)	02 (22.2)	09 (100)	05 (55.5)
<i>Enterobacter</i> spp (n=1)	00	00	00	00	00	00	00	00	00	00	00	00	00

Amp = Ampicillin, cepl= Cephalexin, CFM= Cefixime, CAZ= Ceftazidime, CRO= Ceftriaxone, CTX= Cefotaxime, Tet= Tetracycline, Gen= Gentamycin, Cip= Ciprofloxacin, Chl= Chloramphenicol, Cot= Co-trimoxazole, Imp= Imipenem, Net= Netilmicin

Almost all of *Acinetobacter* species (31, 96.8%) were susceptible to Imipenem, While 23 (71.8%) and 21 (65.6%) of the isolates were sensitive to Tetracycline and Gentamicin respectively. The sensitivity to third-generation Cephalosporins ranged between 50-56%, whereas 40.6% were found sensitive to Ciprofloxacin. The sensitive of the organism to Chloramphenicol, Co-trimoxazole, Cephalexin and Ampicillin were 34.3%, 31.2%, 15.6% and 9.3% respectively.

Out of 28 isolates of *Pseudomonas* species, 26 (92.8%) were sensitive to Imipenem. Only 01 (3.5%) isolate was susceptible to Ampicillin, while 12(42.8%) were sensitive to Carbenicillin and Cefotaxime. The sensitivity of the organism to Ciprofloxacin and Ceftriaxone was 39.2% each, while the rate of susceptibility to Gentamicin and Netilmicin was higher than the formers (67.8 % and 53.5%). Susceptibility to Co-trimoxazole, Chloramphenicol and Tetracycline were between 17.8%-32.1% respectively. (Table III)

Table III: Antibiogram of isolated *Acinetobacter* species and *Pseudomonas* species by disk diffusion method

Organisms	Number (%) sensitive to-												
	Amp	Cepl	CXM	CAZ	CRO Gen	CTX	Tet	Cip	Chl	Cot	Imp	Net	
<i>Acinetobacter</i> (n=32)	03 (9.3)	05 (15.3)	15 (46.8)	17 (53.1)	18 (56.2)	16 (50.0)	23 (71.8)	21 (65.6)	13 (40.6)	11 (34.3)	10 (31.2)	31 (96.8)	18 (56.2)
<i>Pseudomonas</i> spp (n=28)*	01 (3.5)	-	-	13 (46.4)	11 (39.2)	12 (42.8)	09 (32.1)	19 (67.8)	11 (39.2)	08 (28.5)	05 (17.8)	26 (92.8)	15 (53.5)

* Carbenicillin was tested in addition and 12 (42.8%) *Pseudomonas* were sensitive

Considering sensitivity of *Staphylococcus aureus*, It was found that only 05 (7.4%) and 08 (11.9%) of the isolates were sensitive to Penicillin and

Ampicillin respectively. Only 20 (29.8%) isolates were sensitive to 1 µg Oxacillin showing a calculated 70.2% of the isolates being Methicillin

resistant or MRSA. Only 23 (34.3%) of the isolates were sensitive to Ciprofloxacin. However, all of them were sensitive to Vancomycin and the

sensitivity to Rifampicin and Fusidic acid ranged between 85-88%. (Table IV)

Table IV: Antibiogram of isolated *S. aureus* by disk diffusion method

Organisms	Number (%) sensitive to-												
	Pen	Amp	Ox	Cepi	Cip	Ery Gen	Tet	Chl	Cot	Net	Van	Rif	
<i>S. aureus</i> (n=67)	05 (7.43)	08 (11.9)	20 (29.5)	25 (37.31)	23 (34.3)	15 (22.3)	24 (35.8)	39 (58.2)	44 (65.6)	18 (26.8)	53 (79.1)	67 (100)	57 (85.0)

Amp = Ampicillin, Cepi= Cephalexin, CXM= Cefuroxime, CAZ= Ceftazidime, CRO= Ceftriaxone, CTX= Cefotaxime, Tet= Tetracycline, Gen= Gentamicin, Cip= Ciprofloxacin, Chl= Chloramphenicol, Cot= Co-trimoxazole, Imp= Imipenem, Net= Netilmicin, Pen= Penicillin, Ox= Oxacillin, Ery= Erythromycin, Rif=Rifampicin, Van= Vancomycin and Fus= Fusidic acid

Discussion

Antibiotic resistance is a common phenomenon in developing countries where drugs are available freely without prescription. Now-a-days, antibiotics have been extensively and newer antibiotics are continuously being added for the treatment of various infections. Proper use of antibiotics is very important in reducing unnecessary expenses, development of resistance to useful and life-saving antibiotics as well as to minimize many side effects. The resistance pattern varies from one country to another.

In the present study, most of the Enterobacteriaceae were sensitive to Imipenem ranging between 94-100%. This high level of sensitivity to Imipenem could be due to its restricted and limited use in the clinical practice. The drug has only recently been introduced in Bangladesh and is very expensive which has further restricted its widespread use. Similar effectiveness of Imipenem has also been reported from other countries^{15,7,16}.

The third-generation cephalosporins like Ceftriaxone, Ceftazidime and Cefotaxime were sensitive against 45-66% isolated Enterobacteriaceae. The first-and second-generation Cephalosporins were less effective. In the United States, the frequency of resistance to Ceftazidime has increased from 1.5% to 3.6% from 1991 to 1997 as reported by the National Nosocomial Infections Surveillance system. A surveillance trial involving 102 medical centres in the United States detected 10.3% and 23.8% Ceftazidime-resistant *E.coli* and *K. pneumoniae* respectively¹⁶.

However, the sensitivity of various Enterobacteriaceae to Ciprofloxacin in the present study was only between 33-40%. This low-level of sensitivity to Quinolones and Cephalosporins was the result of very extensive use of these antibiotics in clinical practice. A large majority of patients were found prescribed by these drugs on their first contact with physicians¹⁶.

All the staphylococcus aureus isolated in this study were sensitive to Vancomycin whereas 70% were Methicillin resistant (MRSA). The prevalence of MRSA differs strongly between countries¹⁷. In the present study, very high isolation rate of MRSA was found as detected by 1 µg Oxacillin disk. Another study with wound specimens from diabetic patients in Bangladesh in 1994, reported an isolation rate of 37.2% and 21.6% MRSA amongst the hospitalized and non-hospitalized diabetics respectively¹⁸. The present findings show that the prevalence of MRSA in hospitalized patients has increased significantly over time.

This study would help the physicians to make judicious choice of antibiotics and would be helpful for formulation of an antibiotic policy.

References

1. Berger-Bachi B. Resistance mechanisms of gram-positive bacteria. *Int J Med Microbiol* 2002; 292:27-35.
2. Poole K. overcoming multidrug resistance in gram-negative bacteria. *Curr Opin investig Drugs* 2003;4:128-139
3. Jones R.N, Marshall SA. Antimicrobial activity of cefepime tested against Bush group I beta-

- lactamase producing strains resistant to ceftazidime. *Diagn Microbiol Infect Dis* 1994; 19: 33-38
4. Mocerling RC. Emerging resistance with Gram-positive aerobic infections. Where do we go from here? Introduction: Problems with antimicrobial resistance in Gram-positive cocci. *Clin Infect Dis* 1998; 26:1177-1178
 5. Jones RN, Salazar JC, Pfaller MA, Doern GV. Colombian Antimicrobial Resistance Study (CARS) Group. Multi center evaluation of the antimicrobial activity for six broad-spectrum beta-lactams in Colombia using the E-test method. *Diagn Microbiol Infect Dis* 1997; 29:265-272
 6. National Committee for Clinical Laboratory Standards. Tentative standard M2-A6: Performance Standards for Antimicrobial Disk Susceptibility Tests. NCCLS, Villanova, Pa; 1997.
 7. Yamaguchi K, Mathai D, Biedenbach DJ, Lewis M, Gales AC, Jones RN, Japan Antimicrobial Resistance Study (JARS) Group. Evaluation of the in vitro activity of six broad-spectrum beta-lactam antimicrobial agents tested against over 2000 clinical isolates from 22 medical centers in Japan. *Diagn Microbiol Infect Dis* 1999; 34:123-134.
 8. Sader HS, Mimica I, Rossi F, et al. Evaluation of the in vitro activity of cefepime compared to other broad spectrum cephalosporins against clinical isolates from eighteen Brazilian hospitals by using the E-test. *Diagn Microbiol Infect Dis* 1997; 28:87-92.
 9. Shanahan PMA, Thospson CJ, Amyes SGB. The global of antibiotic-resistant bacteria: their sources and reservoirs. *Rev Med Microbiol* 1994; 5:174-182.
 10. Asna SMZ, Haq JA. Decrease of antibiotic resistance in *Salmonella typhi* isolated from patients attending hospitals of Dhaka City over a 3-year period. *Intl J Antimicrob Agents* 2000; 3:249-251
 11. Bequi AAMA, Rahman KM. Transferable drug resistance in *Pseudomonas aeruginosa*, Bangladesh. *Med Res Council Bull* 1987; 2:61-68
 12. Ahmad AAA, Srinivas. Patterns of antibiotic resistance of *Klebsiella pneumoniae*. *Bangladesh Med Rev* 1987; 13:9-14.
 13. Baron EJ, Peterson LR, Finegold SM. Enterbacteriaceae. In Bailey and Scotts Diagnostic Microbiology, 9th edition. St. Louis: Mosby; 1994: pp.374-379
 14. Bauer AW, Kirby WMM, Sherris JC, Truck M. Antibiotic susceptibility testing by a standard single disk method. *Am J Clin Pathol* 1966; 45: 493-496.
 15. National Committee for Clinical Laboratory Standards. Tentative standard M100-S8: Performance Standards for Antimicrobial Susceptibility Testing. NCLS, Villanova, Pa; 1998.
 16. Johnson DM, Biedenbach DJ, Jones RN, The Philippines Antimicrobial Resistance Study (PARS) Group. In vitro evaluation of broad-spectrum beta-lactams in the Philippines medical centers: role of fourth-generation cephalosporins. *Diagn Microbiol Infect Dis* 1999; 35:291-297
 17. Christina MJE, Vandenbroucke G. Management of Methicillin resistant *Staphylococcus aureus* in the Netherlands. *Rev Med Microbiology* 1998; 9:109-116
 18. Jinnah F, Chowdhury K, Begum J, et al. Multidrug resistant *Staphylococcus aureus* isolated from the wound of Diabetic patients. *J Infect Dis Antimicrob Agents* 1998; 15:15-18.

All corresponds to
Bulbul Hasan
 Associate Professor
 Department of Microbiology
 Rajshahi Medical College
 Rajshahi, Bangladesh
 E-mail: bulbul010163@yahoo.com