



## Review Article

# Optimum Iron Provision Potentiate the Cognitive Functions in Children- A Review Article

Md Belal Uddin,<sup>1</sup> Laila Shamima Sharmin,<sup>2</sup> Khan Ishrat Jahan<sup>2</sup>  
Manirul Haque Tarafder,<sup>3</sup> Md Razikul Islam<sup>4</sup>

### Abstract

Micronutrient deficiencies, especially those related to iodine and iron, are linked to different cognitive impairments, as well as to potential long-term behavioral changes. Among the cognitive impairments caused by iron deficiency, those referring to attention span, intelligence, and sensory perception functions are mainly cited, as well as those associated with emotions and behavior, often directly related to the presence of iron deficiency anemia. In addition, iron deficiency without anemia may cause cognitive disturbances. At present, the prevalence of iron deficiency and iron deficiency anemia is 2%–6% among European children. Given the importance of iron deficiency relative to proper cognitive development and the alterations that can persist through adulthood as a result of this deficiency, the objective of this study was to review the current state of knowledge about this health problem. The relevance of iron deficiency and iron-deficiency anemia, the distinction between the cognitive consequences of iron deficiency and those affecting specifically cognitive development, and the debate about the utility of iron supplements are the most relevant and controversial topics. Despite there being methodological differences among studies, there is some evidence that iron supplementation improves cognitive functions. Nevertheless, this must be confirmed through adequate follow-up studies among different groups.

**Keywords:** Iron deficiency, anemia, cognitive functions, supplementation.

TAJ 2021; 34 (1): 115-122

### Introduction

The importance of iron in cognitive functions is highlighted in some early studies.<sup>1</sup> In those studies, some alterations with regard to psychomotor development and cognitive functions have been generally linked to the deficiencies of this mineral. Overall, micronutrient deficiencies are linked to different cognitive impairments, especially iodine and iron, as well as to potential long-term behavioral changes.<sup>2</sup> Generally, a lack of iron in the neonatal period and early childhood is

considered as a key to the development of disturbances in cognitive development.<sup>3</sup> At the same time, there is little consensus about the effectiveness of possible preventive programs. Cognitive disturbances in general and vasomotor development have been related to iron deficiency (ID), and more recently, a possible link between dieting, ID, and the possible presentation of cognitive alterations has been studied.<sup>4</sup>

Experimental studies show that ID is capable of causing cognitive impairment in humans, with

<sup>1</sup> Professor, Department of Pediatrics, Rajshahi Medical College, Rajshahi.

<sup>2</sup> Assistant Professor, Department of Pediatrics, Rajshahi Medical College, Rajshahi.

<sup>3</sup> Registrar, Department of Pediatrics, Rajshahi Medical College Hospital, Rajshahi.

<sup>4</sup> Assistant Registrar, Department of Paediatrics, Rajshahi Medical College Hospital.

brain mitochondrial damage as a basis for these alterations.<sup>5</sup> Among the cognitive impairments caused by ID, those related to attention span, intelligence, and sensory perception functions are mainly cited, as well as those related to emotions and behavior. Generally, these impairments have been related to iron deficiency anemia (IDA). In addition, it must be noted that ID without anemia may cause cognitive disturbances. Concerning ID and IDA, some concepts must be noted. Iron status is a continuum from IDA to ID without anemia to normal iron status with varying amounts of stored iron and finally to possible iron overload. ID is defined as a condition in which there are no mobilizable iron stores and signs of a compromised supply of iron to tissues, including the erythrocytes, are detected. When iron-deficient erythropoiesis occurs, hemoglobin concentrations are reduced to below optimal levels.

The existence of anemia is considered when individual hemoglobin levels are below two standard deviations of the distribution mean for hemoglobin in an otherwise normal population of the same sex and age who are living at the same altitude.<sup>6</sup> Twenty years ago, ID with anemia affected > 45% of children aged <5 years in developing countries and >7% in developed countries.<sup>7</sup>

Currently, it is estimated to affect 2%–6% of European children, with a significant reduction in prevalence.<sup>7</sup> Enriched foods and the use of supplements have contributed to that reduction, among other factors.<sup>19</sup> Despite this improvement in the monitoring of ID, it seems that some health problems are associated with long-term deficits in cognitive and motor function and achievement gaps in education (small to gestational age infants, too short breastfeeding period, ID, severe and moderate protein-energy malnutrition in early childhood, and intrauterine ID). In addition, behavioral differences have also been found in long-term follow-up studies of children who had protein-energy malnutrition.<sup>2</sup>

In this field of study, the relevance of ID and IDA with respect to cognitive function is the main topic in the literature. The distinction between the cognitive consequences of ID and those affecting

specifically cognitive development is another relevant point. The debate about the utility of iron supplements is also a topic that remains controversial.

Given the relevance of the problem of ID relative to proper cognitive development and the alterations that can persist through adulthood as a result of this deficiency, the objective of this study was to review the current state of knowledge about this health problem, trying to conclude with some practical conclusions.

## **Materials and methods**

Searching was carried out on references included in the papers, published reviews, and via hand searching. The literature search was mainly but not exclusively based on the last 15 years. Some former articles considered as pioneers and/or especially relevant were also included.

Studies meeting the following criteria were included in the review: 1) studies focused on cognitive disturbances and ID and 2) controlled trials, randomized controlled trials, and comparative studies. The exclusion criteria were 1) case reports, 2) populations with nonspecific ID-related problems, 3) participants with special conditions/pathologies other than ID (e.g., iodine deficiency), and 4) full text/abstract not available. Previous studies (reviews and meta-analysis) were considered as another source of articles, provided that they fitted the inclusion criteria. As a result, the following main different topics were obtained: anemia vs. ID, neurocognitive development and educational performance, and the controversy about treatment and supplementation.

An analysis was carried out about references included in the papers as well as via hand searching. After that, selected references were analyzed based on the full text. After applying the exclusion criteria, 89 studies were finally included.

## **Results**

### **Anemia versus iron deficiency**

In a study about the effects of children's IDA on executive functions at ten years, it has been found that those who had anemia showed slower reaction

times, less accuracy, and higher latency peak N2 and smaller amplitude of the P300 wave (which is related to working memory) in the electroencephalogram (EEG).<sup>8</sup>

Compared with a control group (non-IDA infants), IDA infants (aged 3–15 months) showed lower scores in cognition, fine motor, and social/emotional areas. In addition, they had higher delta/theta and lower alpha power in EEG activity. Children aged 8–10 years were studied in order to determine the cognitive effects of IDA by means of the EEG P300 wave and some psychometric tests. As a result, P300 latency in the anemic girls was delayed as compared with the control group, but no statistically significant difference was observed for P300 latency and P300 amplitude between the control group and the anemic group.<sup>9</sup>

Poorer object permanence and short-term memory encoding and/or retrieval have been shown in infants with IDA aged 9–12 months. These cognitive effects were attributable, in part, to IDA-related deficits in socioemotional function. Children with poor socioemotional performance seemed to be more vulnerable to the effects of IDA on cognitive function.<sup>10</sup>

If ID occurs during the first two years of life, it is associated with poor psychomotor performance and changes in behavior, such as reduced levels of responsivity to persons and stimuli, irritability, and inhibition. Different studies demonstrate that effects observed during infancy persist in the long term. It has been reported that formerly anemic preschool children are less active, more inhibited, and more timid than the corresponding controls.<sup>11</sup>

ID and IDA may have some prolonged effects on behavior and development.<sup>29</sup> In a review about the effects of ID in the first 2 years, it is concluded that the relationship between ID and IDA to cognitive and mental development in childhood is still unclear. Generally, follow-up studies have found poorer cognitive scores on measures of mental and cognitive functioning in the long run.<sup>12</sup>

Non-IDA predominantly due to anemia inflammation has been related to lower performance on tests of cognitive function, this

likely being due to decreased delivery of iron to host tissues, including the central nervous system. In fact, the cognitive achievement is strongly related to hemoglobin level (higher hemoglobin levels result in better central nervous system function) and could be expected in anemic patients.<sup>12</sup>

In a study focused on the effect of ID on cognitive development among children, the authors concluded that the effect of short-term iron treatment in anemic children aged <2 years is controversial.<sup>4</sup> With respect to anemic children aged >2 years, it seems that iron treatment would be more effective. Finally, they stated that preventing IDA could produce benefits to development but that they would be small and transient.

#### **COGNITIVE DEVELOPMENT AND EDUCATIONAL PERFORMANCE**

A consequence of ID relies on alterations in cognition that occur among iron-deficient individuals during the early parts of their life cycle and perhaps at later times as well. While ID was presumed to cause most of its effects only if anemia was present, it is clear that many organs show morphologic, physiologic, and biochemical changes before there is any significant drop in hemoglobin concentration. ID is associated with alterations in many metabolic processes that may impact brain functioning (e.g., mitochondria electron transport, neurotransmitter synthesis and degradation, protein synthesis, organogenesis). It is necessary to separate the developmental aspects of ID and neural functioning from the aspects of ID that could occur at any time in life.<sup>13</sup>

In a recent study, a chronic ID group showed significantly lower scores on language, environmental sound perception, and motor measures when compared with infants with normal nutritional iron status at six months and 14–18 months. The conclusion was that the development of language and motor skills and environmental sound perception appeared to be sensitive to the effects of chronic ID in infants.<sup>14</sup> In fact, IDA produces alterations in cognitive processes related to visual attention and concept acquisition in

children. Lower school achievement and poorer fine-hand movements have been described in children.<sup>15</sup>

For some authors, iron status does not play a major role in the educational performance and intelligence of school-going adolescents, stating that several factors (apart from ID) affect educational performance and intelligence.<sup>40</sup> Eden<sup>41</sup> highlights that children with ID and IDA, especially those in lower socioeconomic groups, are at high risk for neurodevelopmental impairment both from the ID itself as well as from central nervous system damage caused by the associated increased absorption of other metals like lead. With regards to the low socioeconomic groups, 26% of post-institutionalized children have been shown to be iron deficient. ID is associated with neurobehavioral alterations months after postadoption, mediated by the effect on attention and activity levels (hyperactivity).<sup>16</sup>

There is evidence that, despite iron repletion, ID during the brain growth spurt alters metabolism and neurotransmission, myelination, and gene and protein profiles. In humans, there is compelling evidence that 6 to 24-month-old infants with ID are at risk for poorer cognitive, motor, socioemotional, and neurophysiologic development in short and long-term outcomes. Alterations in the mesolimbic pathway, where dopamine plays a major role in behavioral activation and inhibition, positive affect, and inherent reward, may help explain altered socioemotional behavior in iron-deficient infants.<sup>17</sup>

On the other hand, IDA resistant to iron administration (with the consequent failure of iron therapy) has been observed in most trials of children aged <2 years but not in older children. For children aged >2 years and in adolescents with IDA, evidence suggests cognitive or behavioral deficits.<sup>17</sup>

### **The treatment and supplementation**

Several benefits of iron supplementation, independent of increasing hemoglobin, such as those on immune function, physical performance, thermoregulation, cognition, and restless leg syndrome, among others, have been described.

Concerning the behavioral problems associated with IDA, it seems that early iron supplementation of marginally low birth weight infants does not affect cognitive functions at age 3.5 years but significantly reduces the prevalence of behavioral problems. For other authors, iron seems to have a modest effect on linear growth in deficient populations.<sup>18</sup>

The question is, does it depend on the previous ID (without anemia), or is it necessary to suffer from IDA? In this regard, another study reported the benefits of iron supplementation in an urban population of nonanemic iron-deficient adolescent girls. That supplementation improved verbal learning and memory. Another study showed that in healthy formerly iron-deficient anemic children, the altered cognitive function might be prolonged despite the iron treatment.

The authors suggested that long-lasting changes in myelination and energy metabolism, perhaps especially in the hippocampus, might account for these long-term effects on an important aspect of human cognitive development. This study tested two groups of adolescents (nonanemic ID and iron-deficient anemic) after eight months of iron supplementation. This supplementation resulted in significant improvements in cognitive function and school achievements.<sup>19</sup>

Another study highlights that there seems to be a dose-response relationship between hemoglobin and cognitive function in children with ID, whereas no similar evidence has been found in iron-sufficient children.<sup>20</sup>

In a study, it was reported that fine motor functioning among offspring was positively associated with prenatal iron/folic acid supplementation in an area where ID was prevalent.<sup>21</sup>

In another study, authors gave an iron supplementation to term breastfed infants (aged 1–6 months). Concerning the development effects, that supplementation resulted in higher visual acuity and developmental indexes at age 13 months. After controlling for background factors, children who had severe, chronic ID in infancy showed poorer arithmetic achievement and written

expression, motor functioning, and some specific cognitive processes, and this occurred ten years after treatment.<sup>22</sup>

It was found that children aged between 16 months and 24 months improved their performance in neurodevelopmental measures when they had received iron in a timely manner. In preschool children, some improvements in cognitive functions have been reported after iron supplementation: e.g., in discrimination, specifically selective attention.<sup>24</sup>

In a former study, it was reported that the administration of heme iron-fortified cookies might improve the intellectual performance of low-income preschool children. The study has reported cognitive improvements after iron supplementation in children. With respect to cognitive development,<sup>24</sup> it has been reported that iron treatment of young children with IDA has a positive short-term effect on psychomotor development or cognitive function, but the effect of longer-term treatment remains unclear. Not only iron but also a mixture of iron and folic acid supplementation seems to produce beneficial effects on cognition in young adolescent girls.<sup>25</sup>

On the contrary, this study states that neither combined nor individual supplements improve the performance on tasks such as A-not-B or fixation duration in infants.<sup>26</sup> These authors did not find any improvements in cognitive performance after supplementation with multiple micronutrient-fortified drinks. The effect of several drinks containing different micronutrients and sugar seems to be positive when those beverages are given alone. In other cases, the combination attenuates that positive effect on cognition.<sup>27</sup>

It was found that, in children with insufficient iron and n-3 fatty acid status, iron supplementation improved verbal and nonverbal learning and memory, particularly in children with anemia. In contrast, docosahexaenoic acid/eicosapentaenoic acid supplementation had no benefits on cognition and impaired working memory in anemic children and long-term memory and retrieval in girls with ID.<sup>28</sup>

## Discussion

Previous reviews on this field of the study concluded that ID has a negative impact on cognition, behavior, and motor skills that can have long-term implications.<sup>4</sup>

Many studies highlight that the relationship between ID and cognitive consequences could be confounded by external factors such as the low socioeconomic background. Apart from socioeconomic factors, ID seems to impact areas such as the hippocampus, and the cognitive dysfunction might be due to mitochondrial damage. In addition, changes in brain dopamine metabolism are presumed to occur, as well as altered serotonergic neurotransmission<sup>5</sup> and alterations in the dopamine receptors. In fact, brain ID and/or IDA result in major changes in dopamine levels in crucial brain areas. These changes related to central (brain) ID might cause different effects than those caused by IDA.<sup>1,22,29</sup>

More recently, it was reported that alterations in the mesolimbic pathway, positive affect, and inherent reward might give support to explain the altered socioemotional behavior that has been described in children with ID.<sup>17</sup> This author states that poorer motor sequencing and bimanual coordination and lower spontaneous eye blink rate in children with IDA are consistent with impaired function in the nigrostriatal pathway. Generally, nutrition has been shown to affect the brain's macrostructure (e.g., development of brain areas such as the hippocampus), microstructure (e.g., myelination of neurons), and level and operation of neurotransmitters (e.g., dopamine levels or receptor numbers), all of which can have an impact on cognitive development. Specifically, nutrients such as iron and omega-3 polyunsaturated fatty acids, in particular, might have specific effects on frontal lobe functioning.<sup>3</sup>

There are works in which it is possible to obtain a clear idea about the effects of ID and IDA on behavior and development. Generally, in non-IDA, there seems to be lower performance in cognitive functions, so a question arises as to whether ID, IDA, and anemia (other than IDA) may cause the same cognitive deficits. This question has an

unclear response to date. Some studies correlate hemoglobin levels to cognitive function, while others report an improvement in cognitive functions after iron supplementation, which seems not to be correlated to those levels.<sup>30</sup>

Despite the specific effect of hemoglobin levels or iron status not being clear, IDA has proved to cause different cognitive alterations in all stages of life. A relevant effect of ID is caused during brain growth, as stated by a study.<sup>31</sup>

Along with the work of others seem to point out that there are critical periods in which ID might have long-term effects on cognition and behavior (eg, premature babies, infants aged <1 year, and preschool children).<sup>32</sup> Apart from those critical periods, it seems that there are not any substantial different effects of ID or IDA on cognitive functions depending on the age reported clearly in the literature. With regards to the sex, it seems that there are no differences between boys and girls with ID in cognitive test performance.<sup>28</sup>

There seem to be two tendencies considering the supplementation – on the one hand, the use of multimicronutrient supplements and, on the other hand, the use of specific micronutrients such as iron in this field of study. So supplementation, fortification, or treatment should be considered without distinction in order to explore the cognitive improvements in the case of ID and IDA.

Some authors report a behavioral improvement but not better results on cognitive function after iron supplementation, while others highlight an improvement in verbal learning/memory in ID (nonanemic) individuals. Despite years of treatment, there seem to be some cognitive alterations that persist. The controversial results focus not only on the long-term follow-up studies but on short-term ones too. For example, some authors state that an iron supplementation over a period of 8 months or 30 days is effective, while others point out that the supplementation during six months is not effective.<sup>19,24</sup>

Another relevant point highlights the moment to introduce the iron supplementation. When should it be introduced? There are a lot of studies

indicating the success of early iron supplementation: prenatal, age 1–6 months, age >27 months, age 16–24 months, and preschool, among others.<sup>21,22,23</sup>

Finally, there are controversial results with respect to the "mixtures" used as supplements. Thus, the beneficial effect of iron and folic acid supplementation has been reported. On the contrary, another two studies do not find positive results with combined supplements. For others, the combination of supplements might attenuate some positive effects.<sup>25,27</sup>

Iron supplements seem to be beneficial in iron-deficient children, but there is a risk of adverse effects in those who are iron-replete. It has been suggested that excessive iron supplementation of infants may have adverse effects on growth, risk of infections, and even on cognitive development. In addition, excessive iron intake can have negative effects on brain development. So, in populations with a low prevalence of ID, general supplementation should be avoided. The same applies to iron-fortified foods. In this regard, more studies are needed to better determine the risks and benefits of iron supplementation and iron-fortified foods given to iron-deficient and iron-sufficient children.<sup>33</sup>

Despite there being methodological differences among studies, there is some evidence that iron supplementation may improve some functions, such as attention, concentration, and intelligence. Nevertheless, these results require confirmation with well-powered, blinded, independently funded, randomized controlled trials of at least one year's duration in different age groups including children, adolescents, adults, and older people and across all levels of baseline iron status.<sup>34,35</sup>

Many studies differ in the characteristics of the study population, the definition of exposure, type of treatment, and confounders, so it is difficult to assess a causal relationship between ID and IDA and cognitive and mental development in childhood, and the same applies to the use of supplements.<sup>12</sup>

Maybe, an early-introduced correct (adequate) diet is better than the use of supplements. Thus, some

studies suggest that increased meat intake by breastfed infants aged >6 months would adequately support iron requirements.<sup>36</sup>

## Conclusion

There is a consensus on the fact that ID has a negative impact on cognition, behavior, and motor skills. With respect to this negative impact, its causal link with ID might be confounded by external variables such as socioeconomic status.

ID, IDA, and non-ID anemia may cause some cognitive deficits, but it remains unclear if those deficits are the same. These cognitive deficits may appear at any age. Levels of hemoglobin seem to correlate to cognitive performance, but iron supplementation improves cognitive functions regardless of the hemoglobin levels.

Despite the possible action of confounders, ID is clearly related to alterations in areas such as the hippocampus, mitochondrial damage, brain dopamine metabolism, and myelination.

The use of supplements (fortification, treatment) must be based on well-established indications, avoiding their use as a routine. The use of multi-supplements does not seem to add a plus with respect to the use of specific supplements. The success of the supplementation might be based on an early prescription after having diagnosed an iron deficiency. It remains controversial whether that supplementation is successful or not, depending on the timing of the therapy (e.g., critical periods).

More studies of appropriate duration in different age groups (infancy, toddler, children, adolescents) and across all levels of baseline iron status are required.

## References

1. Beard J. One person's view of iron deficiency, development, and cognitive function. *Am J Clin Nutr* 1995; 62:709–710.
2. Grantham-McGregor SM, Walker SP, Chang S. Nutritional deficiencies and later behavioural development. *Proc Nutr Soc* 2000; 59:47–54.
3. Bryan J, Osendarp S, Hughes D. Nutrients for cognitive development in school-aged children. *Nutr Rev* 2004; 62:295–306.
4. Grantham-McGregor S, Ani C. A review of studies on the effect of iron deficiency on cognitive development in children. *J Nutr* 2001; 131:649S–668S.
5. Atamna H, Walter PB, Ames BN. The role of heme and iron-sulfur clusters in mitochondrial biogenesis, maintenance, and decay with age. *Arch Biochem Biophys* 2002; 397:345–353.
6. World Health Organization. *Iron Deficiency Anaemia. Assessment, Prevention, and Control. A Guide for Programme Managers*. Geneva, Switzerland: World Health Organization; 2001.
7. Grantham-McGregor SM, Ani CC. The role of micronutrients in psychomotor and cognitive development. *Br Med Bull* 1999; 55:511–527.
8. Algarin C, Nelson CA, Peirano P. Iron-deficiency anemia in infancy and poorer cognitive inhibitory control at age 10 years. *Dev Med Child Neurol* 2013; 55:453–458.
9. Bandhu R, Shankar N, Tandon OP. Event related potentials in anemic school-going girls of age group 8 to 10 years. *Indian J Physiol Pharmacol* 2011; 55:234–240.
10. Carter RC, Jacobson JL, Burden MJ. Iron deficiency anemia and cognitive function in infancy. *Pediatrics* 2010; 126:e427–e434.
11. De AndracaOyarzun I, Salas Aliaga I, de la Parra Cieciva A. Mother-child interaction and child behavior in preschool children with a history of iron-deficiency anemia in infancy. *Arch Lat Nutr* 1993; 43:191–198.
12. Petranovic D, Batinac T, Petranovic D. Iron deficiency anaemia influences cognitive functions. *Med Hypotheses* 2008; 70:70–72.
13. Beard JL. Iron deficiency and neural development: an update. *Arch Lat Nutr*. 1999; 49:34S–39S.
14. Beltran-Navarro B, Matute E, Vasquez-Garibay E. Effect of chronic iron deficiency on neuropsychological domains in infants. *J Child Neurol* 2012; 27:297–303.
15. Walter T. Effect of iron-deficiency anemia on cognitive skills and neuromaturation in infancy and childhood. *Food Nutr Bull* 2003; 24:S104–S110.
16. Fuglestad AJ, Georgieff MK, Iverson SL. Iron deficiency after arrival is associated with general cognitive and behavioral impairment in post-institutionalized children adopted from Eastern Europe. *Matern Child Health J* 2013; 17:1080–1087.

17. Lozoff B. Early iron deficiency has brain and behavior effects consistent with dopaminergic dysfunction. *J Nutr* 2011; 141:740S–746S.
18. Bhandari N, Bahl R, Taneja S. Effect of micronutrient supplementation on linear growth of children. *Br J Nutr* 2001; 85: S131–S137.
19. Congdon EL, Westerlund A, Algarin CR, et al. Iron deficiency in infancy is associated with altered neural correlates of recognition memory at 10 years. *J Pediatr* 2012; 160:1027–1033.
20. Sungthong R, Mo-suwan L, Chongsuvivatwong V. Once-weekly and 5-days a week iron supplementation differentially affect cognitive function but not school performance in Thai children. *J Nutr* 2004; 134:2349–2354.
21. Christian P, Murray-Kolb LE, Khatry SK, et al. Prenatal micronutrient supplementation and intellectual and motor function in early school-aged children in Nepal. *JAMA* 2010; 304:2716–2723.
22. Lozoff B, Jimenez E, Hagen J, et al. Poorer behavioral and developmental outcome more than 10 years after treatment for iron deficiency in infancy. *Pediatrics* 2000; 105: E51.
23. Matiashvili K, Manjavidze N, Ghonghadze T. Influence of ferrotherapy on psychomotor development of children of yearly age with iron deficiency anemia. *Georgian Med News* 2012; 206:38–41.
24. Idjradinata P, Pollitt E. Reversal of developmental delays in iron-deficient anaemic infants treated with iron. *Lancet* 1993;341(8836):1–4.
25. Sen A, Kanani SJ. Impact of iron-folic acid supplementation on cognitive abilities of school girls in Vadodara. *Indian Pediatrics* 2009; 46:137–143.
26. Siegel EH, Kordas K, Stoltzfus RJ. Inconsistent effects of iron-folic acid and/or zinc supplementation on the cognitive development of infants. *J Health Popul Nutr* 2011; 29:593–604.
27. Taljaard C, Covic NM, van Graan AE. Effects of a multimicronutrient-fortified beverage, with and without sugar, on growth and cognition in South African schoolchildren: a randomised, double-blind, controlled intervention. *Br J Nutr* 2013; 110:2271–2284.
28. Baumgartner J, Smuts CM, Malan L. Effects of iron and n-3 fatty acid supplementation, alone and in combination, on cognition in school children: a randomized, double-blind, placebo-controlled intervention in South Africa. *Am J Clin Nutr* 2012; 96:1327–1338.
29. Youdim MB, Yehuda S. The neurochemical basis of cognitive deficits induced by brain iron deficiency: involvement of dopamine-opiate system. *Cell Mol Biol* 2000; 46:491–500.
30. Agaoglu L, Torun O, Unuvar E. Effects of iron deficiency anemia on cognitive function in children. *Arzneimittel-Forschung* 2007; 57:426–430.
31. Lozoff B, Armony-Sivan R, Kaciroti N. Eyeblinking rates are slower in infants with iron-deficiency anemia than in nonanemic iron-deficient or iron-sufficient infants. *J Nutr* 2010; 140:1057–1061.
32. Yehuda S, Yehuda M. Long-lasting effects of infancy iron deficiency-preliminary results. *J Neurol Trans Suppl* 2006; 71:197–200.
33. Domellof M. Benefits and harms of iron supplementation in iron-deficient and iron-sufficient children. *Nestle Nutr Workshop SerPediatr Program* 2010; 65:153–165.
34. Logan S, Martins S, Gilbert R. Iron therapy for improving psychomotor development and cognitive function in children under the age of three with iron deficiency anaemia. *Cochrane Database Syst Rev* 2001;(2):CD001444.
35. Falkingham M, Abdelhamid A, Curtis P. The effects of oral iron supplementation on cognition in older children and adults: a systematic review and meta-analysis. *Nutr J* 2010; 9:4.
36. Krebs NF. Dietary zinc and iron sources, physical growth, and cognitive development of breastfed infants. *J Nutr* 2000; 130:358S–360S.

All correspondence to:  
**Professor Dr. Md Belal Uddin**,  
 Department of Pediatrics,  
 Rajshahi Medical College, Rajshahi.  
 Email: drmbuddin@yahoo.com